Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CLASSICAL THEOREMS ON QUADRATIC RESIDUES
Autor: Berndt, Bruce C.
Kapitel: 13. Sums of quadratic residues and nonresidues
DOI: https://doi.org/10.5169/seals-48188

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 21.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

THEOREM 12.5. Let χ be even, and let m be an arbitrary positive integer. Then

$$\sum_{j=0}^{m-1} S_{3m,3j+2} = -\frac{3^{1/2}\,G(\chi)}{\pi}\,\bar{\chi}(m)\,L(1,\bar{\chi}_{3k})\,.$$

The instances of Theorem 12.5 with $m = 1, 2, 4$ and 8 are consequences of Theorems 4.1, 6.1, 9.1 and 11.1, respectively.

THEOREM 12.6. Let χ be odd, and let m be an arbitrary positive integer. Then

$$S_{5m,2} - S_{5m,4} + S_{5m,7} - S_{5m,9} + - \cdots + S_{5m,5m-3} - S_{5m,5m-1}$$
$$= -\frac{i5^{1/2}\,G(\chi)}{\pi}\,\bar{\chi}(m)\,L(1,\bar{\chi}_{5k})\,.$$

The special cases of Theorem 12.6 for $m = 1$ and $m = 2$ follow immediately from Theorems 5.1 and 8.1, respectively.

THEOREM 12.7. Let χ be odd, and let m be an arbitrary positive integer. Then

$$S_{12m,2} + S_{12m,3} + S_{12m,4} + S_{12m,5} - S_{12m,8} - S_{12m,9} - S_{12m,10} - S_{12m,11}$$
$$+ + + + - - - - - \cdots - S_{12m,12m-4} - S_{12m,12m-3} - S_{12m,12m-2} - S_{12m,12m-1}$$
$$= -\frac{i(12)^{1/2}\,G(\chi)}{\pi}\,\bar{\chi}(m)\,L(1,\bar{\chi}_{12k})\,.$$

The special instances of $m = 1$ and $m = 2$ of Theorem 12.7 yield results that are easily deduced from Theorems 9.1 and 11.1, respectively.

The class number formula arising from Theorem 12.1 was first proved by Holden [39]. A less general form of Theorem 12.2 was also established by Holden [36] who in another paper [37] used his result to derive formulas for sums of the Legendre-Jacobi symbol over various residue classes. The special case $m = 1$ of the class number formula deducible from Theorem 12.7 is due to Lerch [44, p. 407]. Otherwise, the results of this section appear to be new.

13. SUMS OF QUADRATIC RESIDUES AND NONRESIDUES

We mentioned in the Introduction the two equivalent formulations of Dirichlet's theorem for primes that are congruent to 3 modulo 4. In this section, we state and prove as many theorems as we can that are of the same

nature as (1.3). In the case that $\chi(n)$ is the Legendre symbol, we stated our results in [7, Section 4]. For convenience, we put

$$S_{ji}(\chi, r) = \sum_{(i-1)k/j<n<ik/j} \chi(n)\, n^r ,$$

where i, j, and r are natural numbers. Again, χ is primitive throughout the section.

THEOREM 13.1. Let χ be even. Then

$$S_{21}(\chi, 1) = -\frac{G(\chi)\,k}{\pi^2}\left\{1 - \frac{1}{4}\bar{\chi}(2)\right\} L(2, \bar{\chi}) .$$

Proof. In (12.1), put $f(x) = x$, $c = 0$, and $d = k/2$. Integrating by parts, we find that

$$S_{21}(\chi, 1) = \frac{G(\chi)\,k}{2\pi^2} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n^2} \{\cos(\pi n) - 1\} ,$$

and the desired result readily follows.

COROLLARY 13.2. For any even, real character χ, we have $S_{21}(\chi, 1) < 0$.

In view of Corollary 3.8 and the fact that $S_{21} = 0$ for even χ, Corollary 13.2 is certainly not surprising.

THEOREM 13.3. Let χ be odd. Then

$$S_{21}(\chi, 1) = \frac{iG(\chi)\,k}{2\pi} \{\bar{\chi}(2) - 1\} L(1, \bar{\chi}) .$$

Proof. In (12.2), put $f(x) = x$, $c = 0$, and $d = k/2$. Thus, upon integrating by parts, we get

$$S_{21}(\chi, 1) = \frac{iG(\chi)\,k}{2\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \cos(\pi n) ,$$

from which the desired result readily follows.

COROLLARY 13.4. If χ is real and odd, then $S_{21}(\chi, 1) > 0$, if $\chi(2) \neq 1$, and $S_{21}(\chi, 1) = 0$, otherwise.

In view of Corollary 3.3 and the elementary fact that $S_{41} = 0$ if $\chi(2) = -1$ [8], at least part of Corollary 13.4 is expected. If p is a prime, Corollary 13.4 shows that the sum of the quadratic residues modulo p exceeds the sum of the non-residues on $(0, p/2)$ if $p \equiv 3 \pmod 8$, while the two sums are equal if $p \equiv 7 \pmod 8$.

THEOREM 13.5. Let χ be odd. Then

$$S_{31}(\chi, 1) = -\frac{iG(\chi)k}{\pi}\left\{\frac{1}{6}[1-\bar{\chi}(3)]\,L(1,\bar{\chi}) + \frac{3^{1/2}}{4\pi}L(2,\bar{\chi}_{3k})\right\}.$$

Proof. In (12.2), put $f(x) = x$, $c = 0$, and $d = k/3$. The result follows from the same type of calculation as above.

COROLLARY 13.6. If χ is real and odd, then $S_{31}(\chi, 1) > 0$.

The following theorems are proved in the same manner as above.

THEOREM 13.7. Let χ be odd. Then

$$S_{32}(\chi, 1) = \frac{iG(\chi)k}{\pi}\left\{\frac{1}{6}[\bar{\chi}(3)-1]\,L(1,\bar{\chi}) + \frac{3^{1/2}}{2\pi}L(2,\bar{\chi}_{3k})\right\}.$$

COROLLARY 13.8. If χ is real and odd and if $\chi(3) = 1$, then $S_{32}(\chi, 1) < 0$.

THEOREM 13.9. Let χ be even. Then

$$S_{42}(\chi, 1) = -\frac{kG(\chi)}{4\pi}\left\{L(1,\bar{\chi}_{4k}) + \frac{1}{\pi}[2-\bar{\chi}(2)]\left[1-\frac{1}{4}\bar{\chi}(2)\right]L(2,\bar{\chi})\right\}.$$

COROLLARY 13.10. For χ real and even, we have $S_{42}(\chi, 1) < 0$.

THEOREM 13.11. Let χ be odd. Then

$$S_{41}(\chi, 1) = -\frac{iG(\chi)k}{2\pi}\left\{\frac{1}{4}\bar{\chi}(2)[1-\bar{\chi}(2)]\,L(1,\bar{\chi}) + \frac{1}{\pi}L(2,\bar{\chi}_{4k})\right\}$$

and

$$S_{43}(\chi, 1) = \frac{iG(\chi)k}{2\pi}\left\{[\bar{\chi}(2)-1]\left[\frac{3}{4}\bar{\chi}(2)-1\right]L(1,\bar{\chi}) + \frac{1}{\pi}L(2,\bar{\chi}_{4k})\right\}.$$

COROLLARY 13.12. Let χ be real and odd. If $\chi(2) \neq -1$, then $S_{41}(\chi, 1) > 0$; in any case, $S_{43}(\chi, 1) < 0$.

THEOREM 13.13. Let χ be even. Then

$$S_{11}(\chi, 2) = \frac{G(\chi)k^2}{\pi^2}L(2,\bar{\chi})$$

and

$$S_{21}(\chi, 2) = \frac{G(\chi)k^2}{4\pi^2}\{\bar{\chi}(2) - 2\}\,L(2,\bar{\chi}).$$

COROLLARY 13.14. If χ is even and real, then $S_{11}(\chi, 2) > 0$ and $S_{21}(\chi, 2) < 0$.

THEOREM 13.15. Let χ be odd. Then

$$S_{11}(\chi, 2) = \frac{iG(\chi)k^2}{\pi} L(1, \bar{\chi}) \tag{13.1}$$

and

$$S_{21}(\chi, 2) = \frac{iG(\chi)k^2}{\pi}\left\{\frac{1}{4}[\bar{\chi}(2)-1]L(1, \bar{\chi}) + \frac{1}{\pi^2}\left[1 - \frac{1}{8}\bar{\chi}(2)\right]L(3, \bar{\chi})\right\}.$$

COROLLARY 13.16. Let χ be odd and real. Then in all cases, $S_{11}(\chi, 2) < 0$; if $\chi(2) = 1$, then $S_{21}(\chi, 2) < 0$.

If χ is real, the class number formula corresponding to (13.1) is due to Cauchy [17]. Pepin [51, p. 205], Lerch [44, p. 395], and Ayoub, Chowla, and Walum [3] have also given proofs of (13.1). Of course, any number of formulas could be proven for $\sum_{a \leq n \leq b} \chi(n) n^r$, where r is a positive integer and a and b are rational multiples of k. However we are unable to make any more non trivial deductions about the positivity (or negativity) of such character sums. In this connection, see [3] and [25].

14. SOME QUESTIONS AND PROBLEMS

In the foregoing work, in order to determine if S_{ji} is of constant sign for classes of real, primitive characters, we expressed S_{ji} as a linear combination of L-functions of real characters evaluated as $s = 1$, and then we inspected the coefficients in this linear combination to determine if all were either non-negative or non-positive. In fact, S_{ji} may always be expressed as a linear combination of L-functions evaluated at $s = 1$. However, in the general situation, the L-functions are associated with complex characters. When non-real characters arise in the representation of S_{ji}, we are unable to say anything about the sign of S_{ji}. We have attempted to find all instances when S_{ji} can be expressed in terms of L-functions of real characters. It is natural to ask if these cases are the only instances when theorems about the non-negativity or non-positivity of S_{ji} are possible. Results of P. D. T. A. Elliott (written communication) appear to indicate that this, indeed, is the case. For example, he has proved the following result. Consider the set of primes p in any residue class, e.g., $p \equiv 1 \pmod 8$, and the as-