

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE LIE BRACKET AND THE CURVATURE TENSOR
Autor: Faber, Richard L.
Kapitel: 3. A Particular Case
DOI: <https://doi.org/10.5169/seals-48173>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$(3) \quad f_3 - f_2 = tXf_2 + \frac{t^2}{2} X^2 f_2 + O(3)$$

$$(4) \quad f_2 - f_0 = tYf_0 + \frac{t^2}{2} Y^2 f_0 + O(3)$$

Subtracting (3) and (4) from the sum of (1) and (2), and applying Lemma 1 again (up to $O(2)$ only), we obtain

$$f_4 - f_3 = t^2 (XYf - YXf)_0 + \frac{t^3}{2} (XY^2f - YX^2f)_0 + O(3),$$

or

$$(5) \quad f_4 - f_3 = t^2 [X, Y]_0 f + O(3)$$

The meaning of this is that $[X, Y]$ measures the degree to which the circuit $p_3 \rightarrow p_2 \rightarrow p_0 \rightarrow p_1 \rightarrow p_4$ fails to be closed. Indeed, if $[X, Y] = 0$, then $p_3 = p_4$ (cf. [1, pp. 134-135]).

If we think of $p = p_3$ as the starting point, and (see figure) define $\sigma(u) = Y_u X_u Y_{-u} X_{-u} p$ (so that $p_4 = \sigma(t)$), we may re-express (5) as

$$f(\sigma(t)) - f(\sigma(0)) = t^2 [X, Y]_0 f + O(3) = t^2 [X, Y]_p f + O(3),$$

since switching to p changes $[X, Y]f$ by an amount which is only of order $O(1)$.

3. A PARTICULAR CASE

As a special case, assume X and Y are left invariant vector fields on a Lie group G , i.e., elements of $L(G)$, the Lie algebra of G ; and take p to be e , the identity element of the group. Since, in this context, $X_t(p) = p \exp(tX)$, for p in G , we have

$$\sigma(t) = \exp(-tX) \exp(-tY) \exp(tX) \exp(tY).$$

If we assume $f(e) = 0$, Theorem 1 yields

$$\begin{aligned} & f(\exp(-tX) \exp(-tY) \exp(tX) \exp(tY)) \\ &= t^2 [X, Y]_e f + O(3) \\ &= f(\exp\{t^2 [X, Y] + O(3)\}) \end{aligned}$$

and so

$$\exp(-tX) \exp(-tY) \exp(tX) \exp(tY) = \exp(t^2 [X, Y] + O(3)).$$

This formula is involved in proving that if H is (algebraically) a subgroup of a Lie group G and if H is a closed subset of G , then H is a topological Lie subgroup of G ([3, pp. 96, 105]). Specifically, it implies that $\{V \text{ in } L(G) \mid \exp(tV) \text{ is in } H, \text{ for all } t \text{ real}\}$ is closed under the bracket. The formula also provides the following geometric interpretation of the bracket $[X, Y]$ on the Lie algebra $L(G)$ of a Lie group G .

COROLLARY 1. If X and Y belong to $L(G)$, then the curve

$$t \rightarrow \exp(-\sqrt{t}X) \exp(-\sqrt{t}Y) \exp(\sqrt{t}X) \exp(\sqrt{t}Y)$$

has velocity vector $[X, Y]$ at $t = 0$.

4. THE CURVATURE TENSOR

Now assume M is furnished with an affine connection (covariant differentiation operator) ∇ .

The *curvature tensor* R on M is the $\binom{1}{3}$ -tensor (equivalently, the linear-transformation-valued bilinear mapping) R defined by

$$\begin{aligned} R(X, Y)A &= \nabla_X \nabla_Y A - \nabla_Y \nabla_X A - \nabla_{[X, Y]} A \\ &= ([\nabla_X, \nabla_Y] - \nabla_{[X, Y]}) A, \end{aligned}$$

for X, Y , and A vector fields on M . The relationship between this tensor and the Riemann curvature (in a Riemannian manifold) may be found in [4, pp. 72-73], [2, Chapter 9], and [5, pp. 125-127]. Here we shall show its relationship to parallel translation.

Consider the figure again, and let A be any vector field on M . We shall compare parallel translation along $p_0 \rightarrow p_1 \rightarrow p_4$ with that along $p_0 \rightarrow p_2 \rightarrow p_3$. Then, by adding the curve $\sigma(u) = Y_u X_u Y_{-u} X_{-u} p_3$ defined previously (the dotted curve in the figure), we obtain a closed circuit. We shall need the following.

LEMMA 2. (Taylor's Theorem for parallel translation). Let X be a vector field defined in a neighborhood of a curve γ , let $T = \gamma'(0)$, and for any t in domain (γ) , let τ_t denote parallel translation along γ to $\gamma(t)$. Then

$$\tau_0 X(\gamma(t)) - X(\gamma(0)) = \sum_{k=1}^n \frac{t^k}{k!} \nabla_T^k X + O(n+1).$$