

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE VALUES AT NEGATIVE INTEGERS OF THE ZETA-FUNCTION OF A REAL QUADRATIC FIELD
Autor: Zagier, Don
Kapitel: §3. The Siegel Formula for Quadratic Fields
DOI: <https://doi.org/10.5169/seals-48176>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

while for $K = \mathbf{Q}(\sqrt{13})$

$$\begin{aligned} & \zeta_K(-1) \\ &= \frac{1}{24 \times 13} [1^2 - 2^2 + 3^2 + 4^2 - 5^2 - 6^2 - 7^2 - 8^2 + 9^2 + 10^2 - 11^2 + 12^2] \\ &= \frac{1}{6}. \end{aligned} \tag{18}$$

For a more complete discussion of the formulas treated in this section, see Siegel [8].

§3. THE SIEGEL FORMULA FOR QUADRATIC FIELDS

In this section we shall exploit the simple arithmetic of quadratic fields to evaluate in elementary form the various terms entering into Siegel's formula, thus arriving at an expression for $\zeta_K(1-2m)$ which is elementary in the sense that it involves only rational integers and not algebraic numbers or ideals.

We have to evaluate $s_i^K(2m)$, and to do so we must first know how to compute $\sigma_r(\mathfrak{A})$ for any ideal \mathfrak{A} .

LEMMA. *Let \mathfrak{A} be any ideal of the ring of integers \mathcal{O} of a quadratic field K . Let D be the discriminant of K and $\chi(j) = \left(\frac{D}{j}\right)$ the associated character (as in §2). Then, for any $r \geq 0$,*

$$\sigma_r(\mathfrak{A}) = \sum_{j|\mathfrak{A}} \chi(j) j^r \sigma_r(N/j^2), \tag{1}$$

where $N = N(\mathfrak{A})$ is the norm of \mathfrak{A} , the function σ_r on the right-hand side is the arithmetic function of 1 (12), and the sum is over all positive integers j dividing \mathfrak{A} (i.e. $v/j \in \mathcal{O}$ for every $v \in \mathfrak{A}$; clearly this implies $j^2 \mid N$, so equation (1) makes sense).

Proof: It is very easy to check that both sides of (1) are multiplicative functions, i.e. $\sigma_r(\mathfrak{A}\mathfrak{B}) = \sigma_r(\mathfrak{A})\sigma_r(\mathfrak{B})$ for relatively prime ideals \mathfrak{A} and \mathfrak{B} , and similarly for the expression on the right-hand side of (1). It therefore suffices to take \mathfrak{A} to be a power \mathfrak{P}^m of a prime ideal \mathfrak{P} . Write $N(\mathfrak{P}) = p^i$

where p is a rational prime and $i = 1$ or 2 . Then we can evaluate the left-hand side of (1):

$$\begin{aligned} \sigma_r(\mathfrak{A}) &= \sigma_r(\mathfrak{P}^m) = \sum_{\mathfrak{B} | \mathfrak{P}^m} N(\mathfrak{B})^r \\ &= \sum_{n=0}^m N(\mathfrak{P}^n)^r = \sum_{n=0}^m p^{inr} = \sigma_{ir}(p^m). \end{aligned} \quad (2)$$

To evaluate the right-hand side of (1), we must distinguish three cases, according to the value of $\chi(p)$.

Case 1. $\chi(p) = 1$, $(p) = \mathfrak{P}\mathfrak{P}'$ ($\mathfrak{P}' =$ conjugate of \mathfrak{P}). Then $N(\mathfrak{A}) = N(\mathfrak{P})^m = p^m$. Clearly $j | \mathfrak{A} \Rightarrow j = 1$, for j can only be a power of p (since $j | N(\mathfrak{A})$) and cannot be divisible by p (because $\mathfrak{P}' \nmid p$, $\mathfrak{P}' \nmid \mathfrak{A}$). Hence the sum in (1) has only one term $\sigma_r(N) = \sigma_r(p^m)$, in agreement with (2).

Case 2. $\chi(p) = 0$, $(p) = \mathfrak{P}^2$. Again j can only be a power of p , and since $\chi(p) = 0$, the only term in (1) that does not vanish is the term $j = 1$, namely $\sigma_r(N)$. Since $N = N(\mathfrak{P})^m = p^m$ and $i = 1$, this again agrees with (2).

Case 3. $\chi(p) = -1$, $(p) = \mathfrak{P}$. Now $\mathfrak{A} = \mathfrak{P}^m = (p^m)$, so j can take on the values $1, p, p^2, \dots, p^m$, with $\chi(p^n) = (-1)^n$. Here $i = 2$ and $N = N(\mathfrak{P})^m = p^{2m}$, so we must prove

$$\sigma_{2r}(p^m) = \sum_{n=0}^m (-1)^n p^{nr} \sigma_r(p^{2m-2n}). \quad (3)$$

This is just an exercise in summing geometric series.

The lemma enables us to calculate the generalized sums-of-powers functions $\sigma_r(\mathfrak{A})$ in terms of the ordinary function $\sigma_r(m)$. It remains to see what ideals \mathfrak{A} occur in Siegel's formula. Recall that

$$s_i^k(2m) = \sum_{\substack{v \in \mathfrak{d}^{-1} \\ v \geq 0 \\ \text{tr}(v) = 1}} \sigma_{2m-1}((v)\mathfrak{d}), \quad (4)$$

and that

$$\mathfrak{d} = (\sqrt{D}) \quad (5)$$

for a quadratic field. Furthermore, the ring of integers of K is

$$\mathcal{O} = \left\{ \frac{x + y\sqrt{D}}{2} \mid x, y \in \mathbf{Z}, x^2 \equiv y^2 D \pmod{4} \right\}. \quad (6)$$

We can now describe explicitly the v occurring in the sum (4). Write such a v as $\alpha + \beta\sqrt{D}$ with α and β rational. Then

$$v \in \mathfrak{d}^{-1} \Leftrightarrow v\sqrt{D} \in \mathcal{O}, \quad (7)$$

$$v \geq 0 \Leftrightarrow \alpha > |\beta| \sqrt{D}, \quad (8)$$

$$\text{tr}(v) = l \Leftrightarrow \alpha = l/2. \quad (9)$$

From (6), (7) and (9) we then get $\beta = b/2D$, where b is a rational integer satisfying

$$b^2 \equiv l^2 D \pmod{4} \quad (10)$$

and (because of (8)) also

$$b^2 < l^2 D. \quad (11)$$

Then $(v)\delta$ is the principal ideal

$$(v)\delta = (v\sqrt{D}) = \left(\frac{b}{2} + \frac{l}{2} \sqrt{D} \right). \quad (12)$$

An integer j can divide this only if $j|b$ and $j|l$ and $(b/j)^2 \equiv (l/j)^2 D \pmod{4}$, so by the lemma

$$\sigma_r((v)\delta) = \sum_{\substack{l=jl' \\ b=jb' \\ b'^2 \equiv l'^2 D \pmod{4}}} \chi(j) j^r \sigma_r \left(\frac{l'^2 D - b'^2}{4} \right). \quad (13)$$

We now substitute this into (4), where the summation in (4) is now to be taken over all integers b satisfying (10) and (11), and obtain finally

$$s_l^K(2m) = \sum_{j|l} \chi(j) j^{2m-1} e_{2m-1}((l/j)^2 D), \quad (14)$$

where the arithmetic function $e_r(n)$ is defined by

$$e_r(n) = \sum_{\substack{x^2 \equiv n \pmod{4} \\ |x| \leq \sqrt{n}}} \sigma_r \left(\frac{n - x^2}{4} \right) \quad (15)$$

($r = 0, 1, 2, \dots$; n a positive integer, not a perfect square). Then (15) is a finite sum (empty, if $n \equiv 2$ or $3 \pmod{4}$), and so is (14), so that we have completely evaluated $s_l^K(2m)$ in elementary terms. Then Siegel's theorem states

$$\zeta_K(1-2m) = 4 \sum_{l=1}^r b_l(4m) s_l^K(2m), \quad (16)$$

with $r = [m/3]$ and the coefficients $b_l(4m)$ computable rational numbers tabulated on p. 60 for $1 \leq l \leq 10$.

Using the values of $b_l(4m)$ and equation (14), we can write out the first few cases to illustrate (16): $m = 1$. Here $r = 1$, $b_1(4) = 1/240$, and so (16) reduces to

$$\zeta_K(-1) = \frac{1}{60} s_1^K(2) = \frac{1}{60} e_1(D). \quad (17)$$

Thus for $K = \mathbf{Q}(\sqrt{5})$ we find

$$\begin{aligned} \zeta_K(-1) &= \frac{1}{60} e_1(5) = \frac{1}{60} \left\{ \sigma_1\left(\frac{5-1^2}{4}\right) + \sigma_1\left(\frac{5-(-1)^2}{4}\right) \right\} \\ &= 2\sigma_1(1)/60 = 1/30, \end{aligned} \quad (18)$$

in agreement with 2 (17), and similarly for $K = \mathbf{Q}(\sqrt{13})$

$$\begin{aligned} \zeta_K(-1) &= \frac{1}{60} e_1(13) = \frac{2}{60} \left\{ \sigma_1\left(\frac{13-1^2}{4}\right) + \sigma_1\left(\frac{13-3^2}{4}\right) \right\} \\ &= \frac{2}{60} (3+1+1) = 1/6, \end{aligned} \quad (19)$$

in agreement with 2 (18) (but notice how many fewer terms!). $m = 2$. Here again $r = 1$, and the formula is just as simple:

$$\zeta_K(-3) = \frac{1}{120} s_1^K(4) = \frac{1}{120} e_3(D). \quad (20)$$

Thus with $K = \mathbf{Q}(\sqrt{13})$ we find

$$\zeta_K(-3) = \frac{2}{120} (3^3 + 1^3 + 1^3) = \frac{29}{60}. \quad (21)$$

$m = 3$. Here $r = 2$ and the formula is more complicated:

$$\begin{aligned} \zeta_K(-5) &= \frac{4}{196560} (s_2^K(6) - 24 s_1^K(6)) \\ &= \frac{1}{49140} \{ e_5(4D) + 32 \chi(2) e_5(D) - 24 e_5(D) \}. \end{aligned} \quad (22)$$

Here for $K = \mathbf{Q}(\sqrt{13})$ we get

$$\begin{aligned} \zeta_K(-5) &= (e_5(52) - 56e_5(13))/49140 \\ &= (\sigma_5(13) + 2\sigma_5(12) + 2\sigma_5(9) + 2\sigma_5(4) \\ &\quad - 112\sigma_5(3) - 112\sigma_5(1))/49140 \\ &= 980370/49140 = 3631/182. \end{aligned} \quad (23)$$

TABLE 2.

The Siegel formulas for quadratic fields

$$K = \mathbf{Q}(\sqrt{D}), \quad D = \text{discriminant}, \quad \chi(m) = \left(\frac{D}{m}\right),$$

$$e_r(n) = \sum_{\substack{b^2+4ac=n \\ a,c>0}} a^r.$$

$$60 \zeta_K(-1) = e_1(D)$$

$$120 \zeta_K(-3) = e_3(D)$$

$$49140 \zeta_K(-5) = e_5(4D) + [32 \chi(2) + 24] e_5(D)$$

$$36720 \zeta_K(-7) = e_7(4D) + [128 \chi(2) - 216] e_7(D)$$

$$9900 \zeta_K(-9) = e_9(4D) + [512 \chi(2) - 456] e_9(D)$$

$$13104000 \zeta_K(-11) = e_{11}(9D) + 48e_{11}(4D) + [177147 \chi(3) \\ + 98304 \chi(2) - 195660] e_{11}(D)$$

$$3897600 \zeta_K(-13) = e_{13}(9D) - 192e_{13}(4D) + [1594323 \chi(3) \\ - 1572864 \chi(2) - 151740] e_{13}(D)$$

$$652800 \zeta_K(-15) = e_{15}(9D) - 432e_{15}(4D) + [14348907 \chi(3) \\ - 14155776 \chi(2) - 50220] e_{15}(D)$$

$$1554543900 \zeta_K(-17) = e_{17}(16D) + 72e_{17}(9D) + [131072 \chi(2) \\ - 194184] e_{17}(4D) + [17179869184 \chi(4) + 9298091736 \chi(3) \\ - 25452085248 \chi(2) - 57093088] e_{17}(D)$$

$$312543000 \zeta_K(-19) = e_{19}(16D) - 168e_{19}(9D) + [524288 \chi(2) \\ - 156024] e_{19}(4D) + [274877906944 \chi(4) - 195259926456 \chi(3) \\ - 81801510912 \chi(2) - 19291168] e_{19}(D)$$

$$42124500 \zeta_K(-21) = e_{21}(16D) - 408 e_{21}(9D) + [2097152 \chi(2) \\ - 60264] e_{21}(4D) + [4398046511104 \chi(4) - 4267824106824 \chi(3) \\ - 126382768128 \chi(2) - 3953248] e_{21}(D)$$

TABLE 3.

Values of $\zeta_K(1-2m)$ for quadratic fields

$$Z_1 = 60\zeta_K(-1). \quad Z_3 = 120\zeta_K(-3). \quad Z_5 = 49140\zeta_K(-5). \quad Z_7 = 36720\zeta_K(-7).$$

$$Z_9 = 9900\zeta_K(-9). \quad Z_{11} = 13104000\zeta_K(-11). \quad (K = \mathbf{Q}(\sqrt{D}), D = \text{discriminant})$$

| D | Z_1 | Z_3 | Z_5 | Z_7 | Z_9 | Z_{11} |
|-----|-------|-------|-----------|---------------|------------------|-------------------------|
| 5 | 2 | 2 | 5226 | 110466 | 2476506 | 636229128800 |
| 8 | 5 | 11 | 70395 | 3765483 | 215478075 | 141611774080400 |
| 12 | 10 | 46 | 655590 | 78808158 | 10145592150 | 15002017227306400 |
| 13 | 10 | 58 | 1003890 | 143106714 | 21682075650 | 37653788862335200 |
| 17 | 20 | 164 | 4516980 | 1078232292 | 277803225300 | 823821554778449600 |
| 21 | 20 | 308 | 14017380 | 5219942004 | 2064025431300 | 9353651984246859200 |
| 24 | 30 | 522 | 29672370 | 14265873306 | 7346194920450 | 43450483506376984800 |
| 28 | 40 | 904 | 69359160 | 45338101992 | 31773438504600 | 255789968221174153600 |
| 29 | 30 | 942 | 82614870 | 58740797646 | 44300167762950 | 382856016709462960800 |
| 33 | 60 | 1692 | 173700540 | 156050858556 | 151482447747900 | 1692706573508047636800 |
| 37 | 50 | 2258 | 316311450 | 365256498834 | 448286221058250 | 6306377416787885007200 |
| 40 | 70 | 3154 | 493274730 | 658004816322 | 941093728561050 | 15461657528842738261600 |
| 41 | 80 | 3584 | 572460720 | 794742744672 | 1191020559229200 | 20543995478169063449600 |
| 44 | 70 | 4306 | 830983530 | 1344445147458 | 2327280476401050 | 46266888778260351522400 |

In Table 2 we write out in full the formula for $\zeta_K(1-2m)$ ($1 \leq m \leq 6$) in terms of the arithmetical functions $e_r(n)$. In Table 3 we give the values of $\zeta_K(1-2m)$ for $1 \leq m \leq 6$ and K a quadratic field with discriminant at most 50. Since it is more convenient to tabulate integers, we in fact give the values of

$$Z_{2m-1} = t(m) \zeta_K(1-2m), \quad (24)$$

where $t(m)$ is the bound implied by (16) for the denominator of $\zeta_K(1-2m)$, namely

$$t(m) = L.C.M. \{ \text{denom } 4b_l(4m), 1 \leq l \leq r \}. \quad (25)$$

Because the question of the denominator of $\zeta_K(1-2m)$ is important (namely, a prime p divides this denominator whenever the p -adic analogue of $\zeta_K(s)$ has a pole at $s = 1 - 2m$), it is worthwhile to try to sharpen (25). To do this, we use the result of §2, namely

$$\zeta_K(1-2m) = (B_{2m}/4m^2) \sum_{r=0}^{2m} B_r D^{r-1} \beta_{2m-r}(D), \quad (26)$$

where B_r is the r -th Bernoulli number and

$$\beta_r(D) = \sum_{j=1}^D \chi_D(j) j^r. \quad (27)$$

Set

$$a(m) = \prod_{\substack{3 \leq p \leq 2m+1 \\ p \text{ prime}}} p. \quad (28)$$

For $0 \leq r \leq 2m$, $2a(m) B_r$ is an integer, by von Staudt's theorem, and since $\beta_r(D) \equiv 0 \pmod{4}$, $\frac{1}{2} a(m) B_r D^{r-1} \beta_{2m-r}(D)$ is an integer for $r \geq 1$.

There remains the term $r = 0$ of (26). If D is divisible by an odd prime p but $D \neq p$, then (writing $D = pD'$, with $p \nmid D'$)

$$\beta_{2m}(D) \equiv \sum_{k=1}^p \chi_p(k) k^{2m} \sum_{\substack{j=1 \\ j \equiv k \pmod{p}}}^D \chi_{D'}(j) \pmod{p}, \quad (29)$$

and the inner sum is 0 for $D' > 1$. One also checks easily that $\beta_{2m}(D)$ is always even, is divisible by 8 if $D \equiv 0 \pmod{4}$ and is divisible by 16 if $D \equiv 0 \pmod{8}$. Therefore $\beta_{2m}(D)/D$ is an even integer, unless $D = p$ is a prime ($\equiv 1 \pmod{4}$). In that case,

$$\beta_{2m}(p) = \sum_{k=1}^{p-1} \binom{k}{p} k^{2m} \equiv \sum_{k=1}^{p-1} k^{2m+(p-1)/2} \equiv 0 \pmod{p} \quad (30)$$

if $2m + \frac{p-1}{2}$ is not divisible by $p-1$. Finally, if $2m + \frac{p-1}{2}$ is divisible by $p-1$, then $(p-1) \mid 4m$ and hence $p = 4m + 1$ or $p \leq 2m + 1$. Therefore $a(m)\beta_{2m}(D)/D$ is an even integer here also, except in the one case $D = 4m + 1 = \text{prime}$. Thus, if we set

$$s(m) = a(m) \cdot \text{denom} (B_{2m}/2m^2) \cdot \varepsilon_m, \quad (31)$$

$$\varepsilon_m = \begin{cases} 4m + 1 & \text{if } 4m + 1 \text{ is prime,} \\ 1 & \text{otherwise,} \end{cases} \quad (32)$$

then $s(m)\zeta_K(1-2m)$ will be an integer for all quadratic fields K , and indeed $(s(m)/\varepsilon_m)\zeta_K(1-2m)$ will be an integer for all fields except $\mathbf{Q}(\sqrt{4m+1})$. We have tabulated the two bounds $t(m)$ and $s(m)$ for $1 \leq m \leq 17$ in Table 4, putting the factor ε_m of $s(m)$ in brackets because it only occurs in the denominator of $\zeta_K(1-2m)$ for a single exceptional field K . It will be seen that in general neither of $s(m)$, $t(m)$ divides the other, so that

$$u(m) = G.C.D. \{s(m), t(m)\} \quad (33)$$

gives a better bound than is provided by either the Siegel or the elementary method alone. From the table of values of $u(m)$ one sees that, for instance,

$$3 \mid Z_7, \quad 20 \mid Z_{11} \quad (34)$$

and that

$$5 \mid Z_1 \text{ if } D \neq 5, \quad 13 \mid Z_5 \text{ if } D \neq 13, \quad 17 \mid Z_7 \text{ if } D \neq 17. \quad (35)$$

All of these congruences can be verified in Table 3. Indeed, Table 3 suggests that (34) can be improved to

$$3 \mid Z_5, \quad 9 \mid Z_7, \quad 3 \mid Z_9, \quad 400 \mid Z_{11} \quad (36)$$

and that, as well as the congruences (35), one has

$$5 \mid Z_5, \quad 25 \mid Z_9 \text{ if } D \neq 5. \quad (37)$$

All of these are special cases of the following

CONJECTURE ([6], p. 164). *For any totally real K ,*

$$w_m(K)\zeta_K(1-2m) \in \mathbf{Z}, \quad (38)$$

where the integer $w_m(K)$ is defined as

$$G.C.D. \{ (N\mathfrak{P})^i (N\mathfrak{P}^{2m} - 1), i \geq m, \mathfrak{P} \text{ a prime ideal} \}. \quad (39)$$

TABLE 4.
Bounds for the denominator of $\zeta_K(1-2m)$, K quadratic

| m | $t(m)$ (Siegel bound) | $s(m)$ (elementary bound) | $u(m) = (t(m), s(m))$ |
|-----|---|--|---|
| 1 | 60 = $2^2 3 \cdot 5$ | $2^3 3^2 (5)$ | $2^2 3 \cdot (5)$ |
| 2 | 120 = $2^3 3 \cdot 5$ | $2^5 3^2 5^2$ | $2^3 3 \cdot 5$ |
| 3 | 49140 = $2^2 3^3 5 \cdot 7 \cdot 13$ | $2^3 3^4 5 \cdot 7^2 (13)$ | $2^2 3^3 5 \cdot 7 \cdot (13)$ |
| 4 | 36720 = $2^4 3^3 5 \cdot 17$ | $2^7 3^2 5^2 7 \cdot (17)$ | $2^4 3^2 5 \cdot (17)$ |
| 5 | 9900 = $2^2 3^2 5^2 11$ | $2^3 3^2 5^2 7 \cdot 11^2$ | $2^2 3^2 5^2 11$ |
| 6 | 13104000 = $2^7 3^2 5^3 7 \cdot 13$ | $2^5 3^4 5^2 7^2 11 \cdot 13^2$ | $2^5 3^2 5^2 7 \cdot 13$ |
| 7 | 3897600 = $2^8 3 \cdot 5^2 7 \cdot 29$ | $2^3 3^2 5 \cdot 7^2 11 \cdot 13 \cdot (29)$ | $2^8 3 \cdot 5 \cdot 7 \cdot (29)$ |
| 8 | 652800 = $2^9 3 \cdot 5^2 17$ | $2^3 3^2 5^2 7 \cdot 11 \cdot 13 \cdot 17^2$ | $2^9 3 \cdot 5^2 17$ |
| 9 | 1554543900 = $2^2 3^5 5^2 7 \cdot 13 \cdot 19 \cdot 37$ | $2^3 3^5 \cdot 7^2 11 \cdot 13 \cdot 17 \cdot 19^2 (37)$ | $2^2 3^5 \cdot 7 \cdot 13 \cdot 19 \cdot (37)$ |
| 10 | 312543000 = $2^3 3^2 5^3 7 \cdot 11^2 41$ | $2^5 3^2 5^4 7 \cdot 11^2 13 \cdot 17 \cdot 19 \cdot (41)$ | $2^3 3^2 5^3 7 \cdot 11^2 (41)$ |
| 11 | 42124500 = $2^3 3^2 5^3 11 \cdot 23 \cdot 37$ | $2^3 3^2 5 \cdot 7 \cdot 11^2 13 \cdot 17 \cdot 19 \cdot 23^2$ | $2^3 3^2 5 \cdot 23$ |
| 12 | 141466590720 = $2^9 3^6 5 \cdot 7^3 13 \cdot 17$ | $2^7 3^4 5^2 7^2 11 \cdot 13^2 17 \cdot 19 \cdot 23$ | $2^7 3^4 5 \cdot 7^2 13 \cdot 17$ |
| 13 | 22877225280 = $2^6 3^5 5 \cdot 7 \cdot 13 \cdot 53 \cdot 61$ | $2^3 3^2 5 \cdot 7 \cdot 11 \cdot 13^2 17 \cdot 19 \cdot 23 \cdot (53)$ | $2^3 3^2 5 \cdot 7 \cdot 13 \cdot (53)$ |
| 14 | 2722083840 = $2^{10} 3^5 5 \cdot 7 \cdot 29 \cdot 97$ | $2^5 3^2 5^2 7^2 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 29^2$ | $2^5 3^2 5 \cdot 7 \cdot 29$ |
| 15 | 11448204768000 = $2^8 3^3 5^3 7^2 11 \cdot 13 \cdot 31 \cdot 61$ | $2^3 3^4 5^2 7^2 11^2 13 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31^2 (61)$ | $2^8 3^3 5^2 7^2 11 \cdot 13 \cdot 31 \cdot (61)$ |
| 16 | 1611414604800 = $2^{10} 3^3 5^2 7 \cdot 11 \cdot 13 \cdot 17 \cdot 137$ | $2^{11} 3^2 5^2 7 \cdot 11 \cdot 13 \cdot 17^2 19 \cdot 23 \cdot 29 \cdot 31$ | $2^{10} 3^2 5^2 7 \cdot 11 \cdot 13 \cdot 17$ |
| 17 | 176840092800 = $2^7 3^2 5^2 7 \cdot 17 \cdot 51599$ | $2^3 3^2 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17^2 19 \cdot 23 \cdot 29 \cdot 31$ | $2^3 3^2 5 \cdot 7 \cdot 17$ |

Define an integer $j(m)$ for $m = 1, 2, \dots$ by

$$j(m) = G.C.D. \{ n^{m+2} (n^{2m} - 1), n \in \mathbf{Z} \}. \quad (40)$$

Thus

$$j(1) = 24, j(2) = 240, j(3) = 504, j(4) = 480, \dots$$

Then it is easy to check that, for K a quadratic field, $w_m(K) = j(m)$ (independent of K !) unless K is one of the finitely many fields $\mathbf{Q}(\sqrt{p})$ with p a prime such that $(p-1) \mid 4m$, $(p-1) \nmid 2m$, in which case $w_m(K) = p^{v+1} j(m)$, where p^v is the largest power of p dividing m . This is interesting because the numbers $j(m)$ occur in topology: it is known (now that the Adams conjecture has been proved) that $j(m)$ is precisely the order of the group $J(S^{4m})$. This may be just a coincidence, of course, but could conceivably reflect some deeper connection between the values of zeta-functions and topological K -theory (the conjectured connection between these values and algebraic K -theory was mentioned in the introduction).

§4. THE CIRCLE METHOD AND THE NUMBERS $e_{2m-1}(n)$

In §3 we defined

$$e_r(n) = \sum_{\substack{k^2 \equiv n \pmod{4} \\ |k| \leq \sqrt{n}}} \sigma_r \left(\frac{n - k^2}{4} \right), \quad (1)$$

where r and n are positive integers and, for b a positive integer, $\sigma_r(b)$ is defined as the sum of the r -th powers of the positive divisors of b . Since (1) was only needed for n not a perfect square, we are still at liberty to define $\sigma_r(0)$; we set

$$\sigma_r(0) = \frac{1}{2} \zeta(-r) = -\frac{1}{2} \frac{B_{r+1}}{r+1}. \quad (2)$$

This defines $\sigma_r(b)$ for $b = 0, 1, 2, \dots$; we extend the definition to all real b by setting $\sigma_r(b) = 0$ if $b < 0$ or $b \notin \mathbf{Z}$. Then (1) can be rewritten

$$e_r(n) = \sum_{k=-\infty}^{\infty} \sigma_r \left(\frac{n - k^2}{4} \right). \quad (3)$$

We were led to consider these numbers by Siegel's theorem, which, for real quadratic fields K , expresses the value of $\zeta_K(2m)$ or $\zeta_K(1-2m)$ in terms of the numbers $e_{2m-1}(n)$ with $K = \mathbf{Q}(\sqrt{n})$. In this section we