۱ \$
(·

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A BOUNDARY VALUE CHARACTERIZATION OF WEIGHTED H^1

by Richard L. WHEEDEN 1)

Abstract

We give a proof of the elementary result that for certain weight functions w, the Hardy space H_w^1 can be identified with the class of functions f such that f and all its Riesz transforms $R_j f$ belong to L_w^1 . An important ingredient of the proof is that there exist positive constants c and μ , $0 < \mu < 1$, depending only on the dimension n such that if f belongs to L_w^1 , then

$$N(f)(x) \leq c \left[M_{\mu}(f)(x) + \sum_{j=1}^{n} M_{\mu}(R_{j}f)(x) \right],$$

where N(f) denotes the non-tangential maximal function of the Poisson (or any conjugate Poisson) integral of f, and M_{μ} denotes the Hardy-Littlewood maximal operator of order μ :

$$M_{\mu}(g)(x) = \left(\sup_{h>0} h^{-n} \int_{|y| < h} |g(x+y)|^{\mu} dy\right)^{1/\mu}.$$

§1. INTRODUCTION

Let $F(x, t) = (u(x, t), v_1(x, t), ..., v_n(x, t)), x = (x_1, ..., x_n) \in \mathbb{R}^n, t > 0$, satisfy the Cauchy-Riemann equations in the sense of Stein and Weiss [9]: i.e., $u, v_1, ..., v_n$ are harmonic in

$$R_{+}^{n+1} = \left\{ (x,t) \colon x \in \mathbb{R}^{n}, t > 0 \right\}, \frac{\partial u}{\partial t} + \sum_{j=1}^{n} \frac{\partial v_{j}}{\partial x_{j}} = 0$$

and

$$\frac{\partial v_j}{\partial x_i} = \frac{\partial v_i}{\partial x_j}, \frac{\partial v_j}{\partial t} = \frac{\partial u}{\partial x_j}$$

¹ Supported in part by NSF-MPS75-07596.

— 122 —

there. F is said to belong to H_w^1 , where w is a non-negative measurable function on \mathbb{R}^n , if

$$|||F||| = \sup_{t>0} \int_{\overline{R^n}} |F(x,t)| w(x) dx < +\infty.$$

Letting

$$L_{w}^{1} = \left\{ f \colon ||f||_{1,w} = \int_{\mathbb{R}^{n}} |f(x)| w(x) dx < +\infty \right\},$$

it is immediate that a Cauchy-Riemann system belongs to H_w^1 if and only if the L_w^1 norms of its components are uniformly bounded for t > 0. See also [4], p. 118.

We consider primarily weight functions w satisfying

$$(A_1) \qquad \qquad \frac{1}{|I|} \int_{I}^{\infty} w(x) \, dx \leqslant c \text{ ess inf } w,$$

where I is a "cube" in \mathbb{R}^n , and c is a constant independent of I. (See [7], [3].) If $w \in A_1$ and $f \in L^1_w$, the Riesz transforms of f, defined as the *pointwise* limits

(1)

$$(R_{j}f)(x) = \lim_{\varepsilon \to 0} (R_{j,\varepsilon}f)(x), \ j = 1, ..., n, \text{ where}$$
(1)

$$(R_{j,\varepsilon}f)(x) = c_{n} \int_{|y| > \varepsilon} f(x-y) \frac{y_{j}}{|y|^{n+1}} dy, \ c_{n} = \Gamma\left(\frac{n+1}{2}\right) / \pi^{\frac{n+1}{2}},$$

exist a.e. (See [2].) Moreover, as we shall see, the Poisson and conjugate Poisson integrals of f exist and are finite if $f \in L_w^1$, $w \in A_1$. These will be denoted respectively by

$$(Pf)(x,t) = \int_{R^n} f(x-y) P(y,t) \, dy \,,$$
$$(Q_j f)(x,t) = \int_{R^n} f(x-y) Q_j(y,t) \, dy \,,$$

where

$$P(y, t) = c_n t / (t^2 + |y|^2)^{\frac{n+1}{2}}$$

and

$$Q_j(y,t) = c_n y_j / (t^2 + |y|^2)^{\frac{n+1}{2}}$$

are the Poisson and conjugate Poisson kernels. The vector $(Pf, Q_1 f, ..., Q_n f)$ is of course a Cauchy-Riemann system, and the formulas

 $\lim_{t \to 0} (Pf)(x,t) = f(x), \lim_{t \to 0} (Q_j f)(x,t) = (R_j f)(x)$

hold a.e. if $f \in L_w^1$, $w \in A_1$.

THEOREM 1. Let $w \in A_1$.

(i) If $F = (u, v_1, ..., v_n)$ belongs to H_w^1 , there exists $f \in L_w^1$ such that $R_j f \in L_w^1$, u = Pf and $v_j = P(R_j f) = Q_j f$ for each j. Moreover, there are positive constants c_1 and c_2 , independent of F, such that

(2)
$$c_1 |||F||| \leq ||f||_{1,w} + \sum_{j=1}^{n} ||R_jf||_{1,w} \leq c_2 |||F|||.$$

(ii) Let $f \in L_w^1$. If each $R_i f \in L_w^1$, then the vector

$$F = (Pf, Q_1f, \dots, Q_nf)$$

belongs to H_w^1 . Moreover, $Q_j f = P(R_j f)$ and (2) holds.

Thus, if $w \in A_1$, H_w^1 can be identified with

$$\{f: ||f|| = ||f||_{1,w} + \sum_{1}^{n} ||R_{j}f||_{1,w} < +\infty\},\$$

with equivalence of norms. This result, which is very natural, seems to be generally taken for granted, although there appear to be no proofs (at least of (ii)) in the literature, even when $w \equiv 1$. In the one-dimensional periodic case with $w \equiv 1$, two proofs of (ii) are given in [11], vol. 1: see (4.4), p. 263, and the remark at the bottom of p. 285. Our proof is modelled after the second of these. It is largely technical and contains little that is new; a simpler proof would be interesting. The proof of (i) is fairly standard and included only for completeness.

A weight w is said to belong to A_p , 1 , if there is a constant c such that

$$(A_p) \qquad \left(\frac{1}{|I|}\int_{I} w(x) dx\right) \left(\frac{1}{|I|}\int_{I} w(x)^{-\frac{1}{p-1}} dx\right)^{p-1} \leq c$$

for all cubes I. (See [7]). For 0 , let

$$L_{w}^{p} = \left\{ f: ||f||_{p,w} = \left(\sum_{R^{n}} |f|^{p} w \, dx \right)^{1/p} < + \infty \right\}.$$

In the course of proving (ii), we will derive the following result, analogous to Theorem D of [9], about boundary values of Cauchy-Riemann systems.

THEOREM 2. Let F be a Cauchy-Riemann system for which

$$\sup_{t>0} \int_{\mathbb{R}^{n}} |F(x,t)|^{p} w_{1}(x) dx < +\infty,$$

where $\frac{n-1}{n} and <math>w_1 \in A_{pn/(n-1)}$. Then F(x, t) has a limit F(x, 0) a.e. (and in $L_{w_1}^p$) as $t \to 0$. If $|F(x, 0)| \in L_{w_2}^r$ for $\frac{n-1}{n} < r < \infty$ and $w_2 \in A_{rn/(n-1)}$, there is a constant c, depending only on n and w_1 , such that

(3)
$$\sup_{t>0} \int_{\mathbb{R}^n} |F(x,t)|^r w_2(x) \, dx \leqslant c \, ||F(x,0)||_{r,w_2}^r \, .$$

The case $w_1 = w_2 = 1$ is proved in [9].

It follows (see (10) below) that for $F \in H_w^1$, ||| F ||| and $|| F(x, 0) ||_{1,w}$ are equivalent if $w \in A_{n/(n-1)}$. This is an interesting contrast to Theorem 1, which gives more boundary information, but requires the stronger condition $w \in A_1$.

The method used to prove Theorems 1 and 2 leads to the following result, in which we use the notation

$$N(F)(x) = \sup \{ |F(y,t)| : (y,t) \text{ satisfies } |x-y| < t \},\$$
$$(M_{\mu}f)(x) = \left(\sup_{h>0} h^{-n} \int_{|y| < h} |f(x+y)|^{\mu} dy \right)^{1/\mu}, \mu > 0.$$

THEOREM 3. Let f belong to L_w^1 , and let w satisfy A_1 . Let $F = (Pf, Q_1 f, ..., Q_n f)$. There exist positive constants c and μ depending only on n such that $0 < \mu < 1$ and

$$N(F)(x) \leq c \left[M_{\mu}(f)(x) + \sum_{j=1}^{n} M_{\mu}(R_{j}f)(x) \right].$$

The constant μ above can be taken to be (n-1)/n. It follows easily from this and the results of [7] that if $f \in L^1_{w_1}$ for any w_1 satisfying A_1

— 125 —

and if $f, R_1 f, \ldots, R_n f \in L_{w_2}^p$ for some $p > \mu$ and w_2 satisfying $A_{p/\mu} (=A_{pn/(n-1)})$, then $N(F) \in L_{w_2}^p$ and

$$||N(F)||_{p,w_2} \leq c[||f||_{p,w_2} + \sum_{j=1}^{n} ||R_jf||_{p,w_2}].$$

Finally, as a corollary of Theorem 1, we will show that if $f, R_j f \in L^1$ ($w \equiv 1$) for all *j*, then the Fourier transforms satisfy the standard formula

$$(R_j f)^{\wedge}(x) = i \frac{x_j}{|x|} \hat{f}(x)$$

for $x \neq 0$, and, by continuity, $(R_j f)^{\uparrow}(0) = \hat{f}(0) = 0$. The simple proof is given at the end of §3.

§2. Preliminary results

In this section, we prove some facts, including Theorem 2, which will be useful later.

First, we need several observations about condition A_1 . If g^* denotes the Hardy-Littlewood maximal function of a function g, it is not hard to see that $w \in A_1$ if and only if there is a constant c such that

(4)
$$w^*(x) \leqslant c w(x) \quad \text{a.e.}$$

It is also easy to check that if $w \in A_1$ and I and J are cubes with $I \subset J$, then

(5)
$$\int_{J} w dx \leqslant c \, \frac{|J|}{|I|} \int_{I} w dx \, .$$

Since for any w that is not identically zero, there is a constant c > 0such that $w^*(x) \ge c (1+|x|)^{-n}$, we obtain that $w(x) \ge c (1+|x|)^{-n}$ a.e. if $w \in A_1$. Actually, if $w \in A_1$, there exists δ , $0 < \delta < 1$, such that $w^{1/\delta} \in A_1$ (see [7]), so that $w(x) \ge c (1+|x|)^{-n\delta}$ a.e. This shows that if $f \in L^1_w$, $w \in A_1$, then Pf(x, t) and $Q_j f(x, t)$ are finite and tend to zero as $t \to +\infty$ (for fixed x). In fact, the estimate implies that

(6)
$$\sup_{y} \frac{w(y)^{-1}}{(t+|x-y|)^{n}} \quad ((x,t) \quad \text{fixed}, \quad t > 0)$$

is finite and tends to zero as $t \to +\infty$. Thus, since P(x-y, t) and $Q_j(x-y, t)$ are bounded in absolute value by a multiple of $(t+|x-y|)^{-n}$, it follows that |Pf(x, t)| and $|(Q_j f)(x, t)|$ are bounded by