

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	22 (1976)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	BOUNDARY VALUE CHARACTERIZATION OF WEIGHTED H^1
Autor:	Wheeden, Richard L.
Kapitel:	§2. Preliminary results
DOI:	https://doi.org/10.5169/seals-48178

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

and if $f, R_1 f, \dots, R_n f \in L_{w_2}^p$ for some $p > \mu$ and w_2 satisfying $A_{p/\mu}$ ($= A_{pn/(n-1)}$), then $N(F) \in L_{w_2}^p$ and

$$\|N(F)\|_{p,w_2} \leq c[\|f\|_{p,w_2} + \sum_{j=1}^n \|R_j f\|_{p,w_2}].$$

Finally, as a corollary of Theorem 1, we will show that if $f, R_j f \in L^1$ ($w \equiv 1$) for all j , then the Fourier transforms satisfy the standard formula

$$(R_j f) \hat{\ } (x) = i \frac{x_j}{|x|} \hat{f}(x)$$

for $x \neq 0$, and, by continuity, $(R_j f) \hat{\ } (0) = \hat{f}(0) = 0$. The simple proof is given at the end of §3.

§2. PRELIMINARY RESULTS

In this section, we prove some facts, including Theorem 2, which will be useful later.

First, we need several observations about condition A_1 . If g^* denotes the Hardy-Littlewood maximal function of a function g , it is not hard to see that $w \in A_1$ if and only if there is a constant c such that

$$(4) \quad w^*(x) \leq c w(x) \text{ a.e.}$$

It is also easy to check that if $w \in A_1$ and I and J are cubes with $I \subset J$, then

$$(5) \quad \int_J w dx \leq c \frac{|J|}{|I|} \int_I w dx.$$

Since for any w that is not identically zero, there is a constant $c > 0$ such that $w^*(x) \geq c(1+|x|)^{-n}$, we obtain that $w(x) \geq c(1+|x|)^{-n}$ a.e. if $w \in A_1$. Actually, if $w \in A_1$, there exists δ , $0 < \delta < 1$, such that $w^{1/\delta} \in A_1$ (see [7]), so that $w(x) \geq c(1+|x|)^{-n\delta}$ a.e. This shows that if $f \in L_w^1$, $w \in A_1$, then $Pf(x, t)$ and $Q_j f(x, t)$ are finite and tend to zero as $t \rightarrow +\infty$ (for fixed x). In fact, the estimate implies that

$$(6) \quad \sup_y \frac{w(y)^{-1}}{(t+|x-y|)^n} \quad ((x, t) \text{ fixed}, \quad t > 0)$$

is finite and tends to zero as $t \rightarrow +\infty$. Thus, since $P(x-y, t)$ and $Q_j(x-y, t)$ are bounded in absolute value by a multiple of $(t+|x-y|)^{-n}$, it follows that $|Pf(x, t)|$ and $|Q_j f(x, t)|$ are bounded by

$$c \int_{R^n} |f(y)| \frac{dy}{(t + |x - y|)^n} \leq c \|f\|_{1,w} \left\{ \sup_y \frac{w(y)^{-1}}{(t + |x - y|)^n} \right\},$$

which is finite and tends to zero as $t \rightarrow +\infty$.

In addition to the pointwise existence of $R_j f$ for $f \in L_w^1$, $w \in A_1$, there is also a weak-type estimate: if $m_w(E)$ denotes the w -measure of a set E (i.e., $m_w(E) = \int_E w dx$) and if $R_j^* f$ is defined by

$$(R_j^* f)(x) = \sup_{\varepsilon > 0} |(R_{j,\varepsilon} f)(x)|,$$

then

$$m_w \{ x : (R_j^* f)(x) > \lambda \} \leq c \lambda^{-1} \|f\|_{1,w}, \quad \lambda > 0,$$

with c independent of f and λ . A similar estimate holds for f^* . (See [2], [7].)

We need several facts about condition A_p , $p > 1$, all of which can be found in [2], [6], and [7]. Here we note only that if $w \in A_p$, $p > 1$, there is a constant c such that

$$(B_p) \quad \int_{R^n} \frac{w(y)}{(t + |x - y|)^{np}} dy \leq c t^{-np} \int_{|x-y| < t} w(y) dy, \quad t > 0.$$

(Cf. lemma 1 of [6].) In particular, $w(y)/(1 + |y|)^{np}$ is integrable over R^n if $w \in A_p$. This shows that Pf and $Q_j f$ are finite if $f \in L_w^p$, $w \in A_p$, $p > 1$. In fact, by Hölder's inequality,

$$\int_{R^n} |f(y)| \frac{dy}{(t + |x - y|)^n} \leq \|f\|_{p,w} \left(\int_{R^n} \frac{w(y)^{-p'/p}}{(t + |x - y|)^{np'}} dy \right)^{1/p'},$$

$p' = p/(p-1)$. Since $w \in A_p$, we have $w^{-p'/p} \in A_{p'}$, so that

$$w(y)^{-p'/p}/(1 + |y|)^{np'}$$

is integrable and the last expression is finite.

We need the following lemma about harmonic majorization.

LEMMA 1. Let $s(x, t)$ be subharmonic in R_+^{n+1} and satisfy

$$\sup_{t > 0} \int_{R^n} |s(x, t)|^p w(x) dx < +\infty$$

for some p , $1 \leq p < +\infty$, with $w \in A_p$. Then for $a > 0$,

$$(7) \quad s(x, t+a) \leq P(s(\cdot, a))(x, t).$$

If s is harmonic, equality holds in (7).

Proof. First note by the remarks above that $P(s(., a))(x, t)$ is finite, since $s(., a) \in L_w^p$, $w \in A_p$. Inequality (7) is a corollary of Theorem 2 of [8], provided that we show

$$(a) \quad \sup_{t>0} \int_{R^n} \frac{|s(x, t)|}{(1+t+|x|)^{n+1}} dx < +\infty,$$

$$(b) \quad \lim_{t \rightarrow +\infty} \int_{R^n} \frac{|s(x, t)|}{(1+t+|x|)^{n+1}} dx = 0.$$

If $p > 1$,

$$\begin{aligned} & \int_{R^n} \frac{|s(x, t)|}{(1+t+|x|)^{n+1}} dx \\ & \leq \left(\int_{R^n} |s(x, t)|^p w(x) dx \right)^{1/p} \left(\int_{R^n} \frac{w(x)^{-p'/p}}{(1+t+|x|)^{(n+1)p'}} dx \right)^{1/p'} \\ & \leq c \left(\int_{R^n} \frac{w(x)^{-p'/p}}{(1+t+|x|)^{(n+1)p'}} dx \right)^{1/p'}. \end{aligned}$$

Since $(1+t+|x|)^{(n+1)p'} \geq (1+t)^{p'} (1+|x|)^{np'}$ and $w^{-p'/p}$ satisfies $B_{p'}$, the last expression is at most

$$\frac{c}{1+t} \left(\int_{R^n} \frac{w(x)^{-p'/p}}{(1+|x|)^{np'}} dx \right)^{1/p'} \leq \frac{c}{1+t} \left(\int_{|x|<1} w(x)^{-p'/p} dx \right)^{1/p'},$$

from which (a) and (b) follow. The argument for $p = 1$ is similar, using for example the simple estimate $w(x)^{-1} \leq c(1+|x|)^n$. Finally, if s is harmonic then $s(x, t+a) = P(s(., a))(x, t)$, by applying (7) to both s and $-s$.

LEMMA 2. Let F be a Cauchy-Riemann system for which

$$\sup_{t>0} \int_{R^n} |F(x, t)|^p w(x) dx < +\infty,$$

where $\frac{n-1}{n} < p < \infty$ and $w \in A_{pn/(n-1)}$. Then $F(x, t)$ converges a.e. to a limit $F(x, 0)$ as $t \rightarrow 0$. Moreover, $\|F(x, t) - F(x, 0)\|_{p,w} \rightarrow 0$ as $t \rightarrow 0$, and there is a constant c depending only on n such that

$$(8) \quad N(F)(x) \leq c(|F(x, 0)|^{\frac{n-1}{n}})^*^{\frac{n}{n-1}},$$

where $*$ denotes the Hardy-Littlewood maximal function.

Proof. Except for the last estimate, this lemma is proved in [4]. The method is standard. Let $q = pn/(n-1)$ and $s(y, t) = |F(y, t)|^{\frac{n-1}{n}}$ $= |F(y, t)|^{\frac{p}{q}}$. Then s is non-negative, continuous and (by [9]) subharmonic in R_+^{n+1} . Also,

$$\int_{R^n} s(y, t)^q w(y) dy = \int_{R^n} |F(y, t)|^p w(y) dy \leq c_1, \quad t > 0.$$

Since $q > 1$, there exist $\{t_k\} \rightarrow 0$ and $h \in L_w^q$ such that $\|h\|_{q,w}^q \leq c_1$ and $s(., t_k)$ converges weakly in L_w^q to h —i.e.,

$$\int_{R^n} s(y, t_k) g(y) w(y) dy \rightarrow \int_{R^n} h(y) g(y) w(y) dy$$

if $g \in L_w^{q'}$, $q' = q/(q-1)$. For fixed (x, t) , choose $g(y) = P(x-y, t) w(y)^{-1}$. Since $w \in A_q$, we have $w^{-q'/q}$ ($= w^{1-q'}$) $\in B_{q'}$, and therefore $g \in L_w^{q'}$. For this g , the integral on the left above equals $P(s(., t_k))(x, t)$, which majorizes $s(x, t+t_k)$ by Lemma 1, and the integral on the right equals $(Ph)(x, t)$. Hence,

$$s(x, t) = \lim_{t_k \rightarrow 0} s(x, t+t_k) \leq (Ph)(x, t).$$

Therefore, $|F(x, t)| \leq (Ph)(x, t)^{\frac{q}{p}}$, so that

$$(9) \quad N(F)(x) \leq N(h)(x)^{\frac{q}{p}} \leq ch^*(x)^{\frac{q}{p}}.$$

We have

$$\int_{R^n} h^{*q} w dx \leq c \int_{R^n} |h|^q w dx$$

by [7]. Hence, h^* , and so $N(F)$, is finite a.e., and it follows from [1] that F has non-tangential boundary values $F(x, 0)$ a.e. Moreover,

$$\int_{R^n} |F(x, t) - F(x, 0)|^p w(x) dx \rightarrow 0 \quad \text{as } t \rightarrow 0$$

by dominated convergence:

$$|F(x, t)| \leq N(F)(x) \leq ch^*(x)^{q/p} \in L_w^p.$$

Since $s(x, t) = |F(x, t)|^{\frac{n-1}{n}}$, we have $s(x, t) \rightarrow |F(x, 0)|^{\frac{n-1}{n}}$ a.e. This convergence is also in L_w^q norm since

$$|s(x, t)| \leq N(F)(x)^{\frac{p}{q}} \in L_w^q.$$

Since $s(\cdot, t_k)$ also converges weakly in L_w^q to h , it follows that $h(x) = |F(x, 0)|^{\frac{n-1}{n}}$ a.e. Inequality (8) now follows immediately from (9).

Proof of Theorem 2. Let F be a Cauchy-Riemann system satisfying

$$\sup_{t > 0} \int_{R^n} |F(x, t)|^p w_1(x) dx < +\infty,$$

where

$$\frac{n-1}{n} < p < \infty, \quad w_1 \in A_{pn/(n-1)}.$$

Then F has boundary value $F(x, 0)$ a.e. and in L_w^p by Lemma 2; moreover,

$$N(F)(x) \leq c(|F(x, 0)|^{\frac{n-1}{n}})^{* \frac{n}{n-1}}.$$

If we now assume that

$$|F(x, 0)| \in L_{w_2}^r, \quad \frac{n-1}{n} < r < \infty, \quad w_2 \in A_{rn/(n-1)},$$

then

$$\begin{aligned} \int_{R^n} N(F)(x)^r w_2(x) dx &\leq c \int_{R^n} (|F(x, 0)|^{\frac{n-1}{n}})^{* \frac{nr}{n-1}} w_2(x) dx \\ &\leq c \int_{R^n} |F(x, 0)|^r w_2(x) dx \end{aligned}$$

by [7]. This gives (3) immediately.

Remark. We note in passing that if

$$\sup_{t > 0} \int_{R^n} |F(x, t)|^p w(x) dx < +\infty, \quad \frac{n-1}{n} < p < \infty, \quad w \in A_{pn/(n-1)},$$

then

$$(10) \quad \sup_{t > 0} \int_{R^n} |F(x, t)|^p w(x) dx \approx \|F(\cdot, 0)\|_{p,w}^p.$$

This follows from Theorem 2: the right-hand side is at most a multiple of the left since $F(x, t) \rightarrow F(x, 0)$ in L_w^p ; the converse inequality is just (3) with w_2 and r chosen to be w and p , resp.

§3. PROOF OF THEOREMS 1 AND 3

We will prove Theorem 1 first, beginning with part (i). Let $F \in H_w^1$, $F = (u, v_1, \dots, v_n)$, $w \in A_1$. By Theorem 2, F has boundary values $F(x, 0) = (f(x), g_1(x), \dots, g_n(x))$ pointwise a.e. and in L_w^1 . In particular, $f, g_1, \dots, g_n \in L_w^1$. We will show that $u = P(f)$ and $v_j = P(g_j)$. Since $u(x, s)$ converges to $f(x)$ in L_w^1 , $P(u(\cdot, s))(x, t) \rightarrow (Pf)(x, t)$ as $s \rightarrow 0$:

$$\begin{aligned} |P(u(\cdot, s))(x, t) - (Pf)(x, t)| &= \left| \int_{R^n} [u(y, s) - f(y)] P(x-y, t) dy \right| \\ &\leq \|u(\cdot, s) - f\|_{1,w} \left\{ \sup_y w(y)^{-1} P(x-y, t) \right\}, \end{aligned}$$

where the expression in curly brackets is finite for each (x, t) (see (6)). By Lemma 1, $u(x, s+t) = P(u(\cdot, s))(x, t)$ since u is harmonic. Hence, letting $s \rightarrow 0$, we obtain $u(x, t) = (Pf)(x, t)$, as desired. The argument proving that $v_j = P(g_j)$ is similar.

Now let $G = (Pf, Q_1 f, \dots, Q_n f)$. Then G is a Cauchy-Riemann system with the same first component as F . This implies that the first component of $F-G$ is zero, and so that the others are independent of t ; that is, $v_j - Q_j f$ is independent of t . Thus, $v_j = Q_j f$ if both $v_j(x, t)$ and $(Q_j f)(x, t)$ tend to zero as $t \rightarrow +\infty$ (x fixed). We have already observed this for $Q_j f$. For v_j , the mean-value property of harmonic functions gives

$$\begin{aligned} |v_j(x, t)| &\leq ct^{-n-1} \iint_{|\xi-x|^2 + |t-\eta|^2 < t^2} |v_j(\xi, \eta)| d\xi d\eta \\ &\leq ct^{-n} \sup_{\eta > 0} \int_{|\xi-x| < t} |v_j(\xi, \eta)| d\xi \\ &\leq ct^{-n} \left(\sup_{\eta > 0} \int_{R^n} |v_j(\xi, \eta)| w(\xi) d\xi \right) \left(\sup_{\xi: |\xi-x| < t} w(\xi)^{-1} \right) \\ &\leq ct^{-n} \sup_{\xi: |\xi-x| < t} w(\xi)^{-1}. \end{aligned}$$