

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: BOUNDARY VALUE CHARACTERIZATION OF WEIGHTED H^1
Autor: Wheeden, Richard L.

Bibliographie
DOI: <https://doi.org/10.5169/seals-48178>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$\{(1 + |x|)^{-\alpha}\}_r(\lambda) \leq c(1 + \lambda)^{-\alpha/n}, \lambda > 0.$$

Combining estimates, we obtain

$$\int_{R^n} \phi^p w_1 dx \leq c \int_0^\infty \lambda^{-p} (1 + \lambda)^{-\alpha/n} d\lambda < +\infty$$

if $1 - \frac{\alpha}{n} < p < 1$, as desired. This completes the proof of (ii).

To prove Theorem 3, let $f \in L_w^1$ and $w \in A_1$. Then (11) holds for F , p and w_1 as in the proof of Theorem 1 (ii). (The proof of (11) does not require $R_j f \in L_w^1$.) Hence, by Lemma 2 (see (8)),

$$N(F)(x) \leq c(|F(x, 0)|^{\frac{n-1}{n}})^{*}_{n-1}.$$

Since $F(x, 0) = (f(x), (R_1 f)(x), \dots, (R_n f)(x))$, the conclusion of Theorem 3 follows immediately with $\mu = (n-1)/n$.

To prove the fact stated at the end of the introduction, let

$$f, R_1 f, \dots, R_n f \in L^1.$$

Clearly,

$$P(R_j f)^\wedge(x, t) = \hat{P}(x, t)(R_j f)^\wedge(x) = e^{-2\pi t|x|}(R_j f)^\wedge(x),$$

$$(Q_j f)^\wedge(x, t) = \hat{Q}_j(x, t)f^\wedge(x) = i \frac{x_j}{|x|} e^{-2\pi t|x|} f^\wedge(x) \text{ a.e.},$$

where the Fourier transform is taken in the x variable with t fixed. (Note that for fixed t , $P(x, t)$ belongs to L^1 and $Q_j(x, t)$ belongs to L^2 .) However, these expressions are all equal everywhere since $P(R_j f) = Q_j f$ by Theorem 1 and $P(R_j f) \in L^1$. Therefore, $(R_j f)^\wedge(x) = ix_j |x|^{-1} f^\wedge(x)$, as claimed.

REFERENCES

- [1] CALDERÓN, A.P. On the behavior of harmonic functions at the boundary. *Trans. Amer. Math. Soc.* 68 (1950), pp. 47-54.
- [2] COIFMAN, R.R. and C.L. FEFFERMAN. Weighted norm inequalities for maximal functions and singular integrals. *Studia Math.* 51 (1974), pp. 241-250.
- [3] FEFFERMAN, C.L. and E.M. STEIN. Some maximal inequalities. *Amer. J. Math.* 93 (1971), pp. 107-115.
- [4] GUNDY R.F. and R.L. WHEEDEN. Weighted integral inequalities for the nontangential maximal function, Lusin area integral, and Walsh-Paley series. *Studia Math.* 49 (1973), pp. 101-118.
- [5] HUNT, R.A. On $L(p, q)$ spaces. *L'Ens. Math.* 12 (1966), pp. 249-275.

- [6] — B. Muckenhoupt and R.L. WHEEDEN. Weighted norm inequalities for the conjugate function and Hilbert transform. *Trans. Amer. Math. Soc.* 176 (1973), pp. 227-251.
- [7] MUCKENHOUP, B. Weighted norm inequalities for the Hardy maximal function. *Trans. Amer. Math. Soc.* 165, (1972), pp. 207-226.
- [8] NUALTARANEE, S. On least harmonic majorants in half-spaces. *Proc. London Math. Soc.* 27 (1973), pp. 243-260.
- [9] STEIN, E.M. and G. WEISS, On the theory of harmonic functions of several variables, I. The theory of H^p spaces. *Acta Math.* 103 (1960), pp. 25-62.
- [10] WHEEDEN, R. On the dual of weighted $H^1(|z| < 1)$. *To appear in Studia Math.*
- [11] ZYGMUND, A. *Trigonometric Series*. Vol. 1, 2nd edition. Cambridge Univ. Press, New York, 1959.

(Reçu le 28 août 1975)

Richard L. Wheeden

Rutgers University
New Brunswick, N.J. 08903
USA