

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SUMMABILITY OF SINGULAR VALUES OF L^2 KERNELS.
ANALOGIES WITH FOURIER SERIES
Autor: Cochran, James Alan
Kapitel: 3. Preliminaries
DOI: <https://doi.org/10.5169/seals-48180>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. PRELIMINARIES

Since in this paper we are concerned for the most part with square-integrable kernels $K(x, y)$, and the Lebesgue integral is employed throughout, equalities and inequalities between functions, therefore, are generally to be understood as holding “almost everywhere” (a.e.).

For convenience we take $a = 0, b = \pi$, which we may do without any loss of generality, and consider the class of L^2 kernels $K(x, y)$ with $0 \leq x, y \leq \pi$. In our later work we will need to direct our attention primarily to one of the two variables x, y . Let us choose this to be the first and extend K to be periodic in this variable. There are many ways, of course, to accomplish this task, but a not unreasonable procedure is to first define

$$(3.1) \quad K^{(r)}(x, y) \equiv \frac{\partial^r K(x, y)}{\partial x^r} \quad (r = 0, 1, \dots, s)$$

for some preselected nonnegative integer s , and then assume that $K(x, y)$ is extended, as an even function of x if s is even, and as an odd function of x if s is odd, into the domain $-\pi \leq x \leq 0$, and thence as a periodic function of x with period 2π . This approach ensures that, under suitable restrictions, the classical Fourier series for $K^{(s)}(x, y)$, viewed as a function of its first variable, consists solely of sine terms.

In order to enhance the character of the analogies in which we are interested, we shall say that $K^{(s)}(x, y)$ is in $\text{Lip } \alpha$ if

$$|K^{(s)}(x+h, y) - K^{(s)}(x-h, y)| < |h|^\alpha A(y) \quad (0 < \alpha \leq 1)$$

where $A(y)$ is nonnegative and square-integrable. More generally, for $p \geq 1$, $K^{(s)}(x, y)$ will be said to be in $\text{Lip } (\alpha, p)$ if

$$\int_0^\pi |K^{(s)}(x+h, y) - K^{(s)}(x-h, y)|^p dx < |h|^{\alpha p} A^p(y) \quad (0 < \alpha \leq 1)$$

with $L^2 A \geq 0$. In similar fashion, $K^{(s)}(x, y)$ will be said to be relatively uniformly of bounded variation if for all $N \geq 1$ and arbitrary choice of partition $0 \leq x_0 \leq x_1 \leq \dots \leq x_N \leq \pi$,

$$\sum_{n=1}^N |K^{(s)}(x_n, y) - K^{(s)}(x_{n-1}, y)| < B(y)$$

where $L^2 B \geq 0$. The comparable definitions appropriate whenever the roles of x and y are reversed should be obvious.

Two-variable kernels behave very much like their one-variable analogues as regards integrated Lipschitz conditions. Indeed, the following can be easily established:

PROPERTY 4. Kernels in $\text{Lip}(\alpha, p)$ also belong to $\text{Lip}(\alpha, q)$ for all $1 \leq q < p$. Kernels in $\text{Lip } \alpha$ are automatically in $\text{Lip}(\alpha, p)$ for all $p \geq 1$.

PROPERTY 5. Kernels which are relatively uniformly of bounded variation belong to $\text{Lip}(1, 1)$.

PROPERTY 6. If $K(x, y)$ is absolutely continuous in x , for almost all y , and

$$\int_0^\pi \left[\int_0^\pi |K^{(1)}(x, y)|^p dx \right]^{2/p} dy < \infty,$$

$p > 1$, then $K(x, y)$ is in $\text{Lip}(1, p)$.

PROPERTY 7. If a kernel belongs both to $\text{Lip}(\alpha, p)$ and to $\text{Lip}(\beta, q)$ with $1 \leq p < q$, then it belongs to $\text{Lip}(\gamma, r)$ for all $p \leq r \leq q$, where

$$\gamma = \alpha \frac{p(q-r)}{r(q-p)} + \beta \frac{q(r-p)}{r(q-p)}.$$

A somewhat deeper result is

PROPERTY 8. Whenever $1 \leq p \leq q$, $pq(\alpha-\beta) \geq q-p$, kernels in $\text{Lip}(\alpha, p)$ are automatically also in $\text{Lip}(\beta, q)$.

4. GROWTH ESTIMATES FOR SINGULAR VALUES

We come now to the main thrust of our narrative. The characteristic values associated with a given L^2 kernel $K(x, y)$, $0 \leq x, y \leq \pi$, are those special values of λ for which there exist nontrivial solutions of the homogeneous Fredholm integral equation

$$\phi(x) = \lambda \int_0^\pi K(x, y) \phi(y) dy.$$

The singular values are those positive values μ for which there exist nontrivial $\phi(x)$, $\Psi(x)$ satisfying the coupled equations