

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HOW TO USE RUNGE'S THEOREM
Autor: Rubel, L. A.
DOI: <https://doi.org/10.5169/seals-48184>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

HOW TO USE RUNGE'S THEOREM

by L. A. RUBEL

Runge's Theorem on approximation to analytic functions by polynomials is a powerful tool, and belongs in every analyst's bag of tricks. We illustrate how it can be used by giving three applications of it here. The first two answer problems put to us by D. J. Newman, although we don't believe they originated with him. The third concerns a problem that seems to be part of the folklore. Rather than attempt a detailed history, we merely cite [1], [2], [3, p. 221], [4], [5, §4], and [7]. We give fully detailed proofs, although there are simple geometrical ideas that underlie them. It is our hope that this expository paper will help equip the reader with Runge's Theorem as a versatile tool.

RUNGE'S THEOREM [1, p. 177]. If G is an open set in the complex plane whose complement on the Riemann sphere is connected, if f is an analytic function on G , if K is a compact subset of G and if $\varepsilon > 0$, then there exists a polynomial P such that $|P(z) - f(z)| < \varepsilon$ for all $z \in K$.

PROBLEM 1. Let H be the class of real-valued functions that are harmonic in the open unit disc $D = \{z: |z| < 1\}$ and let H_0 consist of those functions $u \in H$ for which $u(0) = 0$. Let $N_u = \{z \in D: u(z) \leq 0\}$ and $m(u)$ be the area of N_u . Does there exist an $\varepsilon > 0$ such that $m(u) \geq \varepsilon$ for all $u \in H_0$?

Solution. It is plausible to some people that such an ε exists, since by the Gauss mean value theorem, $\iint_{D_r} u \, dx \, dy = 0$ for $r < 1$, where $D_r = \{z: |z| < r\}$, so that in one sense, the negative values of u balance out the positive values. Nevertheless, there is no such ε .

Proof. Take δ with $0 < \delta < 1/2$ and consider the open set $G_\delta = D_{\delta/2} \cup J_\delta$, where J_δ is the keyhole region

$$J_\delta = D_{1-\delta} \setminus (D_\delta^- \cup A_\delta),$$

where

$$A_\delta = \{z = re^{i\theta}: -\delta \leq \theta \leq \delta\}$$

and V^- denotes the topological closure of V . Let f be defined in G_δ by $f(z) = 10$ for $z \in J_\delta$ and $f(z) = 0$ for $z \in D_{\delta/2}$. Notice that f is analytic on G_δ and that G_δ has a connected complement. Let

$$K_\delta = D_{\delta/4}^- \cup J_{2\delta}^-,$$

so that K_δ is a compact subset of G_δ . By Runge's theorem, there is a polynomial $P = P_\delta$ such that

$$\sup \{|P(z) - f(z)| : z \in K_\delta\} < 1.$$

Let

$$u(z) = \operatorname{Re}(P(z) - P(0))$$

so that $u(0) = 0$ and $u(z) \geq 8$ for $z \in J_{2\delta}^-$. Hence $u = u_\delta \in H_0$ and $N_u \subseteq D \setminus J_{2\delta}^-$. But it is easy to verify that the limit of the area of $J_{2\delta}^-$ as $\delta \rightarrow 0$ is the area, π , of D , so that $\lim_{\delta \rightarrow 0} m(u_\delta) = 0$, and there cannot therefore exist an $\varepsilon > 0$ with the desired property.

PROBLEM 2. For $u \in H_0$, let L be the level line of u that passes through 0, i.e. L is that component of $\{z : u(z) = 0\}$ that passes through 0. Let

$$L_{1/2} = \{z \in L : |z| \geq 1/2\}.$$

Is there a finite number M such that if u is not identically 0, then the length of $L_{1/2}$ does not exceed M ?

Solution. The answer is no. We force L to wiggle past so many suitably placed barriers that the length of $L_{1/2}$ must be large.

Proof. Choose a positive integer n , and let

$$K_n = \{0\} \cup \left(\bigcup_{j=1}^{n-1} A_{2j}^n \right) \cup \left(\bigcup_{j=1}^n B_{2j-1}^n \right)$$

where for $v = 1, 2, \dots, 2n-1$,

$$A_v^n = \{z = re^{i\theta} : r = v/2n, -\pi/10 \leq \theta \leq \pi/10\}$$

$$B_v^n = \{z = re^{i\theta} : r = v/2n, -\pi + \pi/10 \leq \theta \leq \pi - \pi/10\}.$$

Further, let

$$G_v^n = \left\{ z = re^{i\theta} : \frac{v}{2n} - \frac{1}{10n} < r < \frac{v}{2n} + \frac{1}{10n}, -\frac{\pi}{20} < \theta < \frac{\pi}{20} \right\}$$

$$H_v^n = \left\{ z = re^{i\theta} : \frac{v}{2n} - \frac{1}{10n} < r < \frac{v}{2n} + \frac{1}{10n}, -\pi + \frac{\pi}{20} < \theta < \pi - \frac{\pi}{20} \right\},$$

and let

$$\Omega_n = D_{n/20} \cup \left(\bigcup_{j=1}^{n-1} G_{2j}^n \right) \cup \left(\bigcup_{j=1}^n H_{2j-1}^n \right).$$

Clearly, Ω_n is an open set in D whose complement is connected, and each component of K_n lies in exactly one component of Ω_n . Let f_n be defined by $f_n(z) = 0$ for $z \in D_{n/20}$, and $f_n(z) = 10$ for z in the other components of Ω_n , so that f_n is an analytic function on Ω_n , and we may apply Runge's theorem to find a polynomial $P = P_n$ so that

$$\sup \{ |P_n(z) - f_n(z)| : z \in K_n \} < 1.$$

In particular, $P_n(0) < 1$ and $P_n(z) > 9$ for $z \in \Gamma_n = K_n \setminus \{0\}$. Let

$$u_n(z) = \operatorname{Re} \{ P_n(z) - P_n(0) \}$$

so that $u_n \in H_0$ and $u_n(z) \geq 8$ for $z \in \Gamma_n$. Now by the maximum modulus theorem, the level line L^n of u_n through 0 must extend to the boundary of D . Yet it must avoid Γ_n . It is then easy to see that there is a positive constant c such that the length of $L_{1/2}^n$ exceeds cn , so that there is no upper bound on the length of $L_{1/2}$.

PROBLEM 3. Do there exist two analytic functions f_1 and f_2 in D such that

$$\liminf_{r \rightarrow 1} \{ |f_1(z)| + |f_2(z)| : |z| \geq r \} = \infty ?$$

Solution. We construct such a pair f_1, f_2 below, by using a gap series to define f_1 and then Runge's theorem to define f_2 . It is easy to see from the minimum modulus theorem that there does not exist a single analytic function f in D such that

$$\liminf_{r \rightarrow 1} \{ |f(z)| : |z| \geq r \} = \infty.$$

Proof. First, we construct f_1 . Choose $n_1 = 5$, and then choose r_1 with $0 < r_1 < 1$ so that $n_1 r_1^{n_1} > 3$. Then choose a positive integer m_1 so that

$$n_1 r_1^{n_1} \geq 3 + \sum_{j=m_1}^{\infty} j r_1^j.$$

Having constructed $n_1, \dots, n_k; r_1, \dots, r_k; m_1, \dots, m_k$, proceed as follows. Choose a positive integer $n_{k+1} > m_k$ so that

$$n_{k+1} > k + 3 + \sum_{j=1}^k n_j.$$

Now choose r_{k+1} , with $r_k < r_{k+1} < 1$, so that

$$n_{k+1} r_{k+1}^{n_{k+1}} > k + 1 + \sum_{j=1}^k n_j.$$

Hence

$$n_{k+1} r_{k+1}^{n_{k+1}} > k + 1 + \sum_{j=1}^k n_j r_{k+1}^j.$$

Then choose a positive integer m_{k+1} so that

$$n_{k+1} r_{k+1}^{n_{k+1}} \geq k + 1 + \sum_{j=1}^k n_j r_{k+1}^j + \sum_{j=m_{k+1}}^{\infty} j r_{k+1}^j.$$

Let

$$f_1(z) = \sum_{k=1}^{\infty} n_k z^{n_k}.$$

It is clear that

$$\lim_{x \rightarrow 1-} f_1(x) = +\infty.$$

We claim that $|f_1(z)| \geq k + 1$ for $|z| = r_{k+1}$. To see this, we write

$$\begin{aligned} |f(r_{k+1}e^{i\theta})| &\geq n_{k+1} r_{k+1}^{n_{k+1}} - \sum_{j=1}^k n_j r_{k+1}^{n_j} - \sum_{j=k+1}^{\infty} n_j r_{k+1}^{n_j} \\ &\geq n_{k+1} r_{k+1}^{n_{k+1}} - \sum_{j=1}^k n_j r_{k+1}^{n_j} - \sum_{j=m_{k+1}}^{\infty} j r_{k+1}^j \geq k + 1. \end{aligned}$$

So

$$\liminf_{r \rightarrow 1} \{|f_1(z)| : z \in E, |z| \geq r\} = \infty,$$

where

$$E = [0, 1) \cup \left(\bigcup_{k=1}^{\infty} \{z : |z| = r_k\} \right).$$

By continuity

$$(*) \quad \liminf_{r \rightarrow 1} \{|f_1(z)| : z \in G, |z| \geq r\} = \infty$$

where G is some open superset in D of the set E . Now to construct f_2 , we observe that the complement in D of G is contained in $\bigcup_{k=1}^{\infty} S_k$, where each S_k is a closed annular sector of the form

$$S_k = \{z = re^{i\theta} : s_k \leq r \leq t_k, \delta_k \leq \theta \leq 2\pi - \delta_k\}$$

and $s_k \uparrow 1$, $t_k \uparrow 1$ and $t_k < s_{k+1}$ for $k = 1, 2, 3, \dots$.

Define g_1 in D by $g_1(z) = 2$. Having defined g_1, \dots, g_n , consider a closed disc U^n with center at 0 that contains S_1, \dots, S_n and a slightly larger open disc D^n that excludes S_{n+1} . Let S'_{n+1} be an open superset of S_{n+1} that does not intersect D^n , and define φ_n in $D^n \cup S'_{n+1}$ by $\varphi_n(z) = n + 2 - \sum_{i=1}^n g_i(z)$ for $z \in S'_{n+1}$ and $\varphi_n(z) = 0$ in D^n . By Runge's theorem, there is a polynomial g_n such that

$$|g_n(z) - \varphi_n(z)| < 2^{-n-2}$$

for $z \in U^n \cup S_{n+1}$. Let $f_2 = \sum_{j=1}^{\infty} g_j$. It is easily verified that the series converges uniformly on compact subsets of D to a function f_2 that is analytic on D . On S_n ,

$$f_2(z) = g_n(z) + \sum_{i=1}^{n-1} g_i(z) + \sum_{i=n+1}^{\infty} g_i(z),$$

so that in S_n

$$|f_2(z)| \geq n + 1 - \sum_{i=n+1}^{\infty} 2^{-i-2} \geq n.$$

Hence

$$(**) \quad \liminf_{r \rightarrow 1} \{ |f_2(z)| : z \in \bigcup_{n=1}^{\infty} S_n, |z| \geq r \} = \infty$$

Since $G \cup \left(\bigcup_{n=1}^{\infty} S_n \right) = D$, we have the desired result on putting (*) and (**) together.

REFERENCES

- [1] CLUNIE, J. On a problem of Gauthier. *Mathematika* 18 (1971), pp. 126-129.
- [2] GAUTHIER, P. M. Une application de la théorie de l'approximation à l'étude des fonctions holomorphes. *Spline Functions and Approximation Theory*, pp. 113-118, Basel and Stuttgart, 1973.

- [3] GUNNING, R. and H. ROSSI. *Analytic Functions of Several Complex Variables*. Englewood Cliffs, 1965.
- [4] KASAHARA, K. and T. NISHIRO. *As announced in Math. Reviews* 38 (1969) #4721.
- [5] LAUFER, H. B. Imbedding annuli in C^2 . *J. D'Analyse Math.* 26 (1973), pp. 187-215.
- [6] SAKS, S. and A. ZYGMUND. *Analytic Functions*. Warsaw, 1952.
- [7] YANAGIHARA, N. A remark on imbedding of the unit disc into C^2 (Japanese). *J. College Arts Sci. Chiba Univ.* 5 (1967), No. 1, pp. 21-24.

(Reçu le 10 février 1976)

L. A. Rubel

University of Illinois
at Urbana-Champaign, Ill. 61801