

§1 Theta characteristics

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **22 (1976)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

where $0, I$ are respectively the zero, identity $g \times g$ matrix. The datum (J_o, e_o) is a symplectic pair, and is standard in the sense that for a fixed (J, e) , giving a symplectic basis for (J, e) amounts to the same thing as giving a linear isomorphism $J_o \simeq J$ compatible with e_o, e . By 0.4, this in turn defines an isomorphism $Q(J_o, e_o) \simeq Q(J, e)$ with the properties stated there.

Going back to the standard situation, there is an obvious identification $Q(J_o, e_o) \simeq (\mathbf{Z}/2\mathbf{Z})^{2g}$, obtained associating with every quadratic form q its values on the canonical basis of J_o . With this identification in mind, the action of J_o on $Q(J_o, e_o)$ defined at the end of 0.2 is the action of $(\mathbf{Z}/2\mathbf{Z})^{2g}$ on itself by translations, and the Arf invariant is given by the mapping $Q: (\varepsilon, \varepsilon') \mapsto \sum \varepsilon_i \varepsilon'_i$, where $\varepsilon, \varepsilon' \in (\mathbf{Z}/2\mathbf{Z})^g$.

We will use the following notation,

$$\begin{aligned} J_0(g) &= (\mathbf{Z}/2\mathbf{Z})^{2g} \\ S_0(g) &= Q(J_0, e_0) \\ S_0^+(g) &= \{s \in S_0(g) / Q(s) = 0\} \\ S_0^-(g) &= \{s \in S_0(g) / Q(s) = 1\} \end{aligned}$$

§ 1 THETA CHARACTERISTICS

1.1 *On an algebraic curve.* Let C be a non-singular projective algebraic curve over an algebraically closed base field k of characteristic different from 2. The set $S(C)$ of *theta characteristics* on C is the set of isomorphism classes of line bundles L on C whose tensor square is isomorphic to the canonical bundle. If $J_2(C)$ is the group of points of order two in $\text{Pic}(C)$, i.e. the multiplicative group of isomorphism classes of line bundles on C whose square is the trivial line bundle \mathbf{O}_C , then clearly $J_2(C)$ acts on the set $S(C)$, and this in a simply transitive way. In addition, there is a function

$$Q: S(C) \rightarrow \mathbf{Z}/2\mathbf{Z}$$

defined by

$$Q(L) = \dim \Gamma(C, L) \quad (2).$$

The following formula holds, where $x, y \in J_2(C)$, $s \in S(C)$, and we use additive notation both for the group law in $J_2(C)$ and the action of $J_2(C)$ on $S(C)$:

$$Q(s) + Q(x+s) + Q(y+s) + Q(x+y+s) = e(x, y).$$

Here, e stands for the intersection pairing on $J_2(C)$. If g is the genus of C , it is proved that $Q^{-1}(0)$ (resp. $Q^{-1}(1)$) has $2^{g-1}(2^g+1)$ (resp. $2^{g-1}(2^g-1)$) elements.

The proof of these assertions goes back to Riemann in the case $k = \mathbf{C}$, and in the general case it may be found in Mumford [5].

1.2 *On a principally polarized abelian variety.* Let X be an abelian variety over k , $\theta: X \xrightarrow{\sim} \hat{X}$ a principal polarization. The set $S(X, \theta)$ of *theta characteristics* on (X, θ) is the subset of $\text{Pic}^\theta(X)$ determined by the symmetric line bundles; i.e. the elements of $S(X, \theta)$ are the isomorphism classes of line bundles L on X belonging to θ and such that $L \simeq i^*(L)$, where $i: X \rightarrow X$ sends $x \in X$ into $-x$. Again, the group X_2 of points of order two in X acts on $S(X, \theta)$ through the induced isomorphism $\theta: X_2 \xrightarrow{\sim} \hat{X}_2$, and this in a simply transitive way. Now, for any symmetric line bundle L on X , there exists a unique isomorphism $\varphi: L \xrightarrow{\sim} i^*(L)$ such that over the zero of X , φ induces the identity on the fibers. Over any $x \in X_2$, the fibers of $L, i^*(L)$ identify naturally, and φ induces the multiplication by some scalar that will be denoted $e_*^L(x)$. It is proved that $e_*^L(x) = \pm 1$, and indeed that $e_*^L: X_2 \rightarrow \mathbf{Z}/2\mathbf{Z}$ is a quadratic form whose associated bilinear form is the intersection pairing e on X_2 . Now we define a mapping

$$Q: S(X, \theta) \rightarrow \mathbf{Z}/2\mathbf{Z}$$

by

$$Q(s) = \text{Arf invariant of } e_*^s.$$

The following formula is valid, where additive notation is used both for group law and group action, and where $s \in S(X, \theta)$, $x, y \in X_2$

$$Q(s) + Q(x+s) + Q(y+x) + Q(x+y+s) = e(x, y).$$

It is also true that, if $g = \dim X$, $Q^{-1}(0)$ (resp. $Q^{-1}(1)$) has $2^{g-1}(2^g+1)$ (resp. $2^{g-1}(2^g-1)$) elements.

All the preceding is proved in or follows easily from § 2 of Mumford [4] and from § 0 above. Note in addition that in $\text{Pic}^{2\theta}(X)$ there is a unique totally symmetric line bundle L_o (i.e. L_o is symmetric and $e_*^{L_o}(x) = 1$ for every $x \in X_2$), and that the symmetric line bundles in $\text{Pic}^\theta(X)$ are the line bundles L such that L^2 is isomorphic with L_o (cf. Mumford [4], *loc. cit.*).