Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	22 (1976)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	FINITE GEOMETRIES IN THE THEORY OF THETA CHARACTERISTICS
Autor:	Rivano, Neantro Saavedra
Kapitel:	§3 Some special cases
DOI:	https://doi.org/10.5169/seals-48185

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

§ 3 Some special cases

I present here some examples in order to motivate the general discussion in Part II. Proofs of most assertions are omitted and they may be found in or follow easily from Part II. The base field is C to simplify things.

3.1 Genus two. Let C be of genus two, and let P_C be the projective space of hyperplanes in $H^{1,0}(C)$. Then P_C is a projective line, and the natural map $C \to P_C$ presents C as a 2-sheeted covering of P_C ramified over a subset $R_C \subset P_C$ with $|R_C| = 6$. From the Riemann-Roch theorem it may be proved that the line bundles L in S(C) with Q(L) = 1, i.e. the odd theta characteristics, are those represented by effective divisors, and from here it follows easily that the set S(C) of odd theta characteristics identifies naturally with R_C . If s_1, s_2, s_3 are three different elements of $S^-(C)$ represented by line bundles L_1, L_2, L_3 , it is also easily proved that $L_1 \oplus L_2 \oplus L_3^{-1}$ is even. From this, and from II 2.4 it follows that there is a natural group isomorphism

$$\operatorname{Sp}(H_1(C, \mathbb{Z}/2\mathbb{Z})) \simeq \operatorname{Aut}(R_C).$$

It follows also from *loc. cit.* that it amounts to the same thing to give a symplectic basis for $H_1(C, \mathbb{Z}/2\mathbb{Z})$ or to give a bijection $S_0^-(2) \simeq R_C$, where $S_0^-(2)$ is the fixed 6-elements set defined in 0.5.

I will discuss $S^+(C)$ in a more general setting:

3.2 Even genus, hyperelliptic case. Let C be hyperelliptic. Then there is a projective line P_C and a map $C \rightarrow P_C$ defined up to unique isomorphisms such that $C \rightarrow P_C$ is a 2-sheeted covering. If R_C is the ramification locus, $|R_C| = 2g + 2$, and R_C identifies naturally with the set of Weierstrass points of C.

The group $H_1(C, \mathbb{Z}/2\mathbb{Z})$ can be reconstructed starting from R_C in the following way. If $\pi = {\pi', \pi''}$ is any partition of R_C into two even-order subsets, L_{π} is the line bundle defined by the divisor $\sum_{P' \in \pi_1'} P - \sum_{P \in \pi_2''} P$ where

 $|\pi'_1| = |\pi'_2|$ and $\{\pi'_1, \pi'_2\}$ partition π' . It is clear that L_{π} is of order two, thus defining an element of H_1 (C, Z/2Z). In this manner one gets a group isomorphism

$$P_2^+(R_c) \cong H_1(C, \mathbb{Z}/2\mathbb{Z})$$

where the group $P_2^+(R_c)$ is defined in II 3.5. It is easily verified that this isomorphism is compatible with the intersection pairing on H_1 and with the alternated bilinear form introduced in *loc. cit*.

All the preceding was valid for any genus g. Now if g is even, it follows from II 3.6 and II 1.4 that we have an isomorphism

$$P_2^-(R_C) \cong S(C)$$

compatible with the structures involved (i.e. an isomorphism of symplectic torsors, cf. II 1.1). The results of II, § 3 may thus be applied to the study of S(C).

Observe that if g is odd, there is a natural theta characteristic; namely, the line bundle of the divisor (g-1) P is independent of the Weierstrass point P (compare II 3.6b)).

3.3 Genus three. Two cases arise for C of genus three:

3.3.1 Chyperelliptic. Then there is the 2-sheeted covering $C \rightarrow P_C$ ramified over R_C with $|R_C| = 8$. It is seen in this case, as in 3.1, that there is a natural identification between $S^-(C)$ and the set of subsets of R_C consisting of exactly two elements. It is convenient to visualize the elements of $S^-(C)$ as segments joining the points of R_C , these being distributed on a plane in an arbitrary way. Then, if s_1, s_2, s_3, s_4 are four different elements of $S^-(C)$, $s_1 - s_2 = s_3 - s_4$ iff the segments corresponding to them produce one of the following configurations



From II 2.7 it follows that there is a canonical isomorphism between the group $Sp(H_1(C, \mathbb{Z}/2\mathbb{Z}))$ and the group of permutations of the set $S^-(C)$ that preserve the "geometry" defined by these quadruples. Two comments are in order:

a) Although the permutation group Aut (R_c) is clearly a subgroup of the automorphism group of the "geometry", not every such automorphism arises from a permutation of R_c .

b) The automorphisms of the geometry do not preserve the type of the configuration, they may send one quadruple of the first type drawn above into the other. However in a continuous family of *hyperelliptic* curves of genus 3, each of the two configurations will be preserved as the curve is deformed.

3.3.2 C non hyperelliptic. Let $Q_C = \mathbf{P}(H^{1,0}(C))$ be the projective space of hyperplanes in $H^{1,0}(C)$. Then Q_C is a projective plane and the natural map $C \to Q_C$ is an immersion. The degree of C in Q_C is the degree of the canonical bundle, i.e. 4 and C is thus a nonsingular plane quartic. It is again a simple exercise to prove that the odd theta characteristics on C correspond to the set of lines in Q_C that are bitangents to C. Thus, if B_C is the set of bitangents to C in Q_C , there is a natural identification

$$B_C\simeq S^-(C).$$

The theme of the 28 bitangents to a nonsingular plane quartic (28 $= 2^{3-1} (2^3-1)$) is a classic one in geometry, see for instance Weber [6], chapter 12. A triple (s_1, s_2, s_3) of bitangents is called *syzygetic* (resp. *azygetic*) if their six points of contact with C lie (resp. do not lie) in a conic. A triple is syzygetic iff $L_4 = L_1 \otimes L_2 \otimes L_3^{-1}$ is an odd characteristic, where L_1, L_2, L_3 are the line bundles corresponding to s_1, s_2, s_3 . When this happens, the two points of contact of the bitangent s_4 corresponding to L_4 , together with the preceding six, make up the full $8 = 2 \times 4$ common points of the conic with the quartic.

An Aronhold system of bitangents (Weber [6]) is a set of seven bitangents such that any different three of them constitute an azygetic triple. The Aronhold systems are exactly the basis for the "geometry" in $S^-(C)$ defined by the syzygetic triples (in the sense of II 4.3). It follows from II 4.4 that the set of Aronhold systems is a torsor over the symplectic group $Sp(H_1(C, \mathbb{Z}/2\mathbb{Z}))$, in particular that they have the same number of elements.

As any two "geometries" with the same genus are isomorphic (II 1.4), one can also speak of Aronhold systems in the hyperelliptic case. It turns out that they correspond to the following configurations



There are 1,451,520 of them as it is "immediately" checked. Again, it will be observed that the automorphisms of the geometry do not preserve the type of the configuration.