Zeitschrift:	L'Enseignement Mathématique			
Herausgeber:	Commission Internationale de l'Enseignement Mathématique			
Band:	22 (1976)			
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE			
Artikel:	FINITE GEOMETRIES IN THE THEORY OF THETA CHARACTERISTICS			
Autor:	Rivano, Neantro Saavedra			
Kapitel:	§1 Symplectic torsors			
DOI:	https://doi.org/10.5169/seals-48185			

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# II. THE ABSTRACT THEORY OF CHARACTERISTICS

# §1 Symplectic torsors

1.1 Definitions. Recall that, if  $\Gamma$  is a group, a  $\Gamma$ -torsor (or torsor over  $\Gamma$ ) is a non-void set endowed with a simply transitive action of  $\Gamma$  on it. Let (J, e) be a symplectic pair, a symplectic torsor over (J, e) is a pair (S, Q) of a J-torsor S and a mapping  $Q: S \to \mathbb{Z}/2\mathbb{Z}$  having the property

(1.1.1) Q(s) + Q(x+s) + Q(y+s) + Q(x+y+s) = e(x, y)

where  $s \in S$ ,  $x, y \in J$ . It is clearly equivalent to ask this property for a fixed  $s \in S$  or for all  $s \in S$ , and it may be thought of as meaning that Q "is a quadratic form." Indeed, any  $s \in S$  sets an identification  $J \cong S(x \mapsto x+s)$ , and through this identification Q becomes the map  $x \mapsto Q(x+s)$ . The above property means that the map  $q_s: J \to \mathbb{Z}/2\mathbb{Z}$  defined by

(1.1.2)  $q_s(x) = Q(x+s) + Q(s)$ 

is a quadratic form whose associated bilinear form is e. According to 0.4, two possibilities may and do arise for Q: either  $Q^{-1}(0)$  has  $2^{g-1}(2^g+1)$ or  $2^{g-1}(2^g-1)$  elements, where  $g = \dim J/2$  will be called the *genus* of (S, Q). In the first case, (S, Q) will be said to be *even*, *odd* in the second. In what follows, all symplectic torsors will be even unless otherwise stated. This because the symplectic torsors that will appear most often will be even and because of the following simple construction. If (S, Q) is an even (resp. odd) symplectic torsor over (J, e), and  $\overline{Q}$  is defined by  $\overline{Q}(s)$ = Q(s) + 1, then  $(S, \overline{Q})$  is an odd (resp. even) symplectic torsor over (J, e).

For a given (S, Q) the following notation will be used

$$S^+ = Q^{-1}(0) \quad S^- = Q^{-1}(1).$$

The elements of S will be often called *characteristics*, those in  $S^+$  are *positive*, those in  $S^-$  are *negative*.

1.2 Morphisms. Let (S, Q), (S', Q') be symplectic torsors respectively over (J, e), (J', e'). For any map  $f: S \to S'$  we define a map  $\sigma_f: J \times S \to J'$ by the property

 $f(x+s) = \sigma_f(x,s) + f(s);$ 

this can be done because S' is a J'-torsor. Now, the following cocycle-type property for  $\sigma_f$  is immediately checked, where  $x, y \in J, s \in S$ 

$$\sigma_f(x+y+s) = \sigma_f(x, y+s) + \sigma_f(y, s),$$

and from it one infers the equivalence of the following statements:

(i) For any  $s, s' \in S, x \in J$ 

$$\sigma_f(x,s) = \sigma_f(x,s').$$

(ii) For some  $s \in S$ , any  $x, y \in J$ 

$$\sigma_f(x+y,s) = \sigma_f(x,s) + \sigma_f(y,s)$$

(iii) For any  $s \in S$ ,  $x, y \in J$ 

$$\sigma_f(x+y,s) = \sigma_f(x,s) + \sigma_f(y,s).$$

So, when these statements hold, one gets a group homomorphism  $\sigma_f: J \to J'$ and has  $f(x+s) = \sigma_f(x) + f(s)$ .

An *isomorphism* of (S, Q) onto (S', Q') is a bijection  $f: S \to S'$  verifying statements (i) to (iii) above, and also the condition

$$Q' \circ f = Q$$
.

It is clear in this case that  $\sigma_f: J \to J'$  is an isomorphism compatible with e, e'. The group of automorphisms of (S, Q) will be denoted Sp(S, Q), so the mapping  $f \to \sigma_f$  is a group homomorphism  $Sp(S, Q) \to Sp(J, e)$ .

1.3 An example. For any given (J, e) there is a canonical example of an even symplectic torsor, namely  $(Q(J, e), Q_e)$ . The J-torsor Q(J, e) was introduced in 0.2, the map  $Q_e$  in 0.3 where it was also remarked that it has property (1.1.1) and that  $Q_e^{-1}(0)$  has  $2^{g-1}(2^g+1)$  elements.

If (J, e), (J', e') are two symplectic pairs, and if  $\sigma: J \to J'$  is a linear isomorphism compatible with e, e', a map  $Q(\sigma): Q(J, e) \to Q(J', e')$  was defined in 0.4, where it was shown that it is an isomorphism of symplectic torsors. Clearly  $Q(\sigma)$  is canonical in any conceivable way.

Indeed, if one still dares in these days to use the language of category theory, what I just did was to define a functor from the category of symplectic pairs to the category of even symplectic torsors (morphisms = isomorphisms, in both cases). In section 1.4 we will see that this is an equivalence of categories.

1.4 Uniqueness of symplectic torsors. It will be shown here, that for a given symplectic pair (J, e) there is essentially only one symplectic torsor over it. Let (S, Q) be such an object; then there is a map

 $f_s: S \to Q(J, e),$ 

defined by the rule  $s \to q_s$ , where  $q_s$  was defined in (1.1.2). Let us prove that  $f_s$  is an isomorphism of symplectic torsors inducing the identity  $id_J: J \to J$ . The formula

$$q_{x+s}(y) = (x+q_s)(y)$$

is a mere restatement of condition (1.1.1), and the formula

$$Q_e \circ f_s = Q$$

follows from the fact that (S, Q) is even and from the meaning of the Arf invariant recalled in 0.3.

The isomorphisms  $f_s$  are canonical, in the following sense. If (S, Q), (S', Q') are symplectic torsors over (J, e), (J', e'),  $f: S \to S'$  is an isomorphism of symplectic torsors inducing an isomorphism  $\sigma: J \to J'$ , then the following square commutes

Recalling the definitions, one has to check for  $s \in S$ ,  $x \in J$  that

$$Q(\sigma(x) + f(s)) + Q(f(s)) = Q(x+s) + Q(s)$$

which is immediate from the definition of isomorphism in 1.2.

It comes out of this that for any isomorphism  $\sigma: J \to J'$  there exists one and only one isomorphism  $f: S \to S'$  inducing it. In particular, the group homomorphism at the end of 1.2.

$$Sp(S, Q) \rightarrow Sp(J, e)$$

is an isomorphism. A useful application of this is the following: If by some unspecified means one is able to construct two symplectic torsors over a pair (J, e), there is a unique isomorphism between them inducing the identity of J.

1.5 Some notation. a) Let J be a vector space over  $\mathbb{Z}/2\mathbb{Z}$ , S a J-torsor. Let's put

$$E(S) = J \amalg S$$

the disjoint union of J, S; on this set there is a structure of vector space over  $\mathbb{Z}/2\mathbb{Z}$ . In fact there is an exact sequence

$$0 \to J \to E(S) \to \mathbb{Z}/2\mathbb{Z} \to 0$$

where J is sent identically onto itself, and the inverse image of 0 (resp. 1) in E(S) is J (resp. S). The addition law in E(S) reduces to the given one on J when both elements are in J, is the action of J on S when one element is in J and the other in S, and finally s + s' (for  $s, s' \in S$ ) is the unique element  $x \in J$  such that x + s = s' (or equivalently x + s' = s).

b) Given the standard pair  $(J_o, e_o)$ , as in 0.5. I will write  $S_o = Q(J_o, e_o)$ ,  $Q_o = Q_{e_o}$ . Both  $J_o$ ,  $S_o$  identify to  $(\mathbb{Z}/2\mathbb{Z})^{2g}$ , but the following notations will be used in compliance with tradition, where  $u_1, ..., u_{2g}$  is the canonical basis. An element of the form

$$\sum_{i=1}^{g} \left( \varepsilon_{i} \ u_{i} + \varepsilon_{i}^{'} \ u_{i+g} \right)$$

will be written  $\binom{\varepsilon}{\varepsilon'}$  or  $\begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}$  whether it is seen in  $J_o$  or  $S_o$  respectively, where  $\varepsilon$ ,  $\varepsilon'$  are row vectors. In particular, the addition law in  $E(S_o)$  is the following:

$\binom{\boldsymbol{\epsilon}}{\boldsymbol{\epsilon'}}$	$+ \begin{pmatrix} \eta \\ \eta' \end{pmatrix}$	) = (	+ 3 + '3	$\begin{pmatrix} \eta \\ \eta' \end{pmatrix}$
$\binom{\epsilon}{\epsilon'}$	$+\begin{bmatrix}\eta\\\eta'\end{bmatrix}$		ε + ε' +	$\eta \\ \eta' \end{bmatrix}$
$\begin{bmatrix} \varepsilon \\ \varepsilon' \end{bmatrix}$	$+\begin{bmatrix}\eta\\\eta'\end{bmatrix}$		ε + ε' +	$\eta \atop \eta' \end{pmatrix}$

# § 2 FINITE GEOMETRIES ON SETS OF CHARACTERISTICS

2.0 Let's fix for paragraph § 2 a symplectic torsor (S, Q) over a symplectic pair (J, e) of genus g. The letter  $\Sigma$  will stand for either the set  $S^+$  of  $S^-$ , its cardinality is  $2^{g-1} (2^g \pm 1)$  (recall that according to 1.1 we assume