Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 22 (1976)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FINITE GEOMETRIES IN THE THEORY OF THETA CHARACTERISTICS

Autor: Rivano, Neantro Saavedra

Bibliographie

DOI: https://doi.org/10.5169/seals-48185

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

It follows from the last part of prop. 3.2.1 that, in this case, the number of odd characteristics in the fundamental sets is congruent to g mod 4. We will see that this is a general fact.

4.5.1 PROPOSITION. *Let* $O(F)$ *be the number of odd characteristics in a fundamental set* F. *Then* $O(F) \equiv g\ (4)$. *Conversely, for any* $l \equiv g\ (4)$, *and* $l \leqslant 2g + 2$, *there are fundamental sets* F *with* $O(F) = l$.

4.5.2 *Proof.* We may safely restrict ourselves to the case where the symplectic torsor is S_X with its standard basis X, and $F = \{A\} + (X \cup \{X\})$ where $A \subset X$ is of even order $|A| = 2k$ (cf. 4.3). Then, in F there are $2k$ characteristics corresponding to subsets of X with $2k - 1$ elements, $2(g-k) + 1$ characteristics with $2k + 1$ elements, and 1 characteristic with $2(g-k) + 1$ elements, namely the ones obtained adding A to respectively the characteristics of the form $\{s\}$ $(s \in A)$, $\{s\}$ $(s \notin A)$, X. When g is even the second and third types have the same parity; when g is odd the first and third types have the same parity. From these remarks, it is easy to see that the number of elements of the same parity in F and $X \cup \{X\}$ are congruent mod 4, and that with this only restriction, this number can be arbitrary for F by conveniently choosing A. The proposition follows from this and from what was said just before its statement.

4.5.3 In Coble [1], additional material on fundamental sets may be found.

REFERENCES

[1] COBLE, A. An application of finite geometry to the characteristic theory of the odd and even theta functions. *Trans. A.M.S. 7* (1906), pp. 241-276.

[2] FAY, J. Theta functions on Riemann surfaces. *Lecture Notes in Mathematics, No. 352*, Springer Verlag (1973).

[3] IGUSA, J. *Theta Functions.* Springer Verlag (1972).

[4] MUMFORD, D. On the equations defining Abelian varieties I. *Invent. Math. 1* (1966), pp. 287-354.

[5] —— Theta characteristics of an algebraic curve. *Ann. Scient. ENS 4* (1971), pp. 181-192.

[6] WEBER, H. *Lehrbuch der algebra*, Band 2, Braunschweig (1896).

(Reçu le 20 février 1976)

Neantro Saavedra Rivano

Dept. de Matemática
Universidad Simón Bolívar
Apartado 5354
Caracas, Venezuela