

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 22 (1976)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** DIRECTIONAL CLUSTER SET EXAMPLE  
**Autor:** Belna, C. L. / Evans, M. J. / Humke, P. D.  
**DOI:** <https://doi.org/10.5169/seals-48186>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# A DIRECTIONAL CLUSTER SET EXAMPLE

by C. L. BELNA, M. J. EVANS and P. D. HUMKE

## §0. INTRODUCTION

Let  $f$  be a mapping from the open upper half plane  $H$  into the Riemann sphere  $W$ . For each point  $x$  on the real line  $R$ , let  $C(f, x)$  and  $C(f, x, \theta)$  denote respectively the total cluster set of  $f$  at  $x$  and the cluster set of  $f$  at  $x$  in the direction  $\theta$  ( $0 < \theta < \pi$ ); then let  $\Theta(x)$  denote the set of directions  $\theta$  for which  $C(f, x, \theta) = C(f, x)$ . E. F. Collingwood [3, Theorem 2 combined with Theorem 3] established the following result.

**THEOREM C.** *Let  $f: H \rightarrow W$  be continuous. Then the set  $\Theta(x)$  is residual at each point  $x$  of a residual subset  $S$  of  $R$ .*

A. M. Bruckner and Casper Goffman [2, p. 510] raised the question as to whether or not there exists a residual set of directions  $\Theta$  such that  $\Theta \subset \Theta(x)$  for each  $x \in S$ . Here we prove

**THEOREM 1.** *There exists a continuous  $f: H \rightarrow W$  such that  $\bigcap_{x \in Q} \Theta(x)$  is a first category set of directions for each residual subset  $Q$  of  $R$ .*

To construct this function (§3), we use certain sets of J.-P. Kahane [5] as building blocks<sup>1)</sup>. Two important properties of these sets are established in §2, and the necessary technical preliminaries are presented in §1. Finally, in §4 we present an example concerning essential directional cluster sets.

## §1. (1/2)-TRAPEZOIDS AND THEIR FOUR DESCENDANTS

By a (1/2)-trapezoid we mean any closed trapezoid  $T$  having bases  $L$  and  $L'$  which lie respectively on the lines  $y = 0$  and  $y = 1$  and for which  $|L| = 2|L'|$ . (Here and throughout this paper, we use  $|\tau|$  to denote the length of the line segment  $\tau$ .) For each (1/2)-trapezoid  $T$ , we set

$$T(1/2) = \{z \in T: \operatorname{Im}(z) = 1/2\}.$$

<sup>1)</sup> The authors wish to thank Professor John R. Kinney for bringing this paper of J.-P. Kahane to their attention.

For real numbers  $p$  and  $p'$ , let  $\tau_{pp'}$  denote the line segment joining the points  $(p, 0)$  and  $(p', 1)$ . Then the *projection of the segment*  $\tau_{pp'}$  is given by

$$\text{proj}[\tau_{pp'}] = p' - p,$$

and the *projection set of the  $(1/2)$ -trapezoid*  $T$  is given by

$$\text{proj}[T] = \{ \text{proj}[\tau_{pp'}] : (p, 0) \in L \text{ and } (p', 1) \in L' \}.$$

For the remainder of this section, suppose  $T$  is a  $(1/2)$ -trapezoid with bases  $L$  and  $L'$ , and let  $L_1$  and  $L_2$  (resp.,  $L'_1$  and  $L'_2$ ) denote the two line segments that remain when the open middle half of  $L$  (resp.,  $L'$ ) is removed with  $L_1$  (resp.,  $L'_1$ ) lying to the left of  $L_2$  (resp.,  $L'_2$ ). Then the *four descendants* of  $T$  are the  $(1/2)$ -trapezoids  $T_1$ ,  $T_2$ ,  $T_3$ , and  $T_4$  having respective bases  $L_1$  and  $L'_1$ ,  $L_1$  and  $L'_2$ ,  $L_2$  and  $L'_1$ , and  $L_2$  and  $L'_2$ .

Now let  $a$  and  $a'$  denote the respective  $x$ -coordinates of the left endpoints of  $L$  and  $L'$ , and set  $l = |L|$ . Then the following list of facts concerning  $T$  and its descendants can easily be established:

(A) If  $\hat{a} = (a + a')/2$ , then

$$T(1/2) = \{ (x, 1/2) : \hat{a} \leq x \leq \hat{a} + 3l/4 \}$$

and, for each  $k = 1, 2, 3$ , and  $4$ , we have

$$T_k(1/2) = \{ (x, 1/2) : \hat{a} + 3(k-1)l/16 \leq x \leq \hat{a} + 3kl/16 \}.$$

That is, the segments  $T_k(1/2)$  ( $k = 1, 2, 3, 4$ ) partition the segment  $T(1/2)$  into four equal subsegments.

(B) If  $\hat{v} = a' - a$ , then

$$\text{proj}[T] = \{ v : \hat{v} - l \leq v \leq \hat{v} + l/2 \},$$

$$\text{proj}[T_3] = \{ v : \hat{v} - l \leq v \leq \hat{v} - 5l/8 \},$$

$$\text{proj}[T_4] = \{ v : \hat{v} - 5l/8 \leq v \leq \hat{v} - 2l/8 \},$$

$$\text{proj}[T_1] = \{ v : \hat{v} - 2l/8 \leq v \leq \hat{v} + l/8 \},$$

and

$$\text{proj}[T_2] = \{ v : \hat{v} + l/8 \leq v \leq \hat{v} + 4l/8 \}.$$

That is, the intervals  $\text{proj}[T_k]$  ( $k = 1, 2, 3, 4$ ) partition the interval  $\text{proj}[T]$  into four equal subintervals.

§2. KAHANE'S SET  $K_\xi$ .

Let  $\xi$  be a real number, and let  $T_\xi$  be the  $(1/2)$ -trapezoid with bases  $L = \{(x, 0): 0 \leq x \leq 1\}$  and  $L' = \{(x', 1): \xi \leq x' \leq \xi + 1/2\}$ .

Let  $T_{11}, T_{12}, T_{13}$ , and  $T_{14}$  be the four descendants of  $T_\xi$  labeled in such a way that the points

$$\alpha_{1k} = \min \{x: (x, 1/2) \in T_{1k}\}$$

satisfy  $\alpha_{1k} < \alpha_{1(k+1)}$  for  $k = 1, 2$ , and  $3$ . Then let  $T_{2k}$  ( $k = 1, 2, \dots, 4^2$ ) be the  $4^2$  descendants of the trapezoids  $T_{1k}$  ( $k = 1, 2, 3, 4$ ) labeled in such a way that the points

$$\alpha_{2k} = \min \{x: (x, 1/2) \in T_{2k}\}$$

satisfy  $\alpha_{2k} < \alpha_{2(k+1)}$  for  $k = 1, 2, \dots, 4^2 - 1$ . Continuing inductively we arrive at the collection of  $(1/2)$ -trapezoids

$$T_{nk} \quad \text{for } n = 1, 2, \dots \quad \text{and } k = 1, 2, \dots, 4^n$$

where the  $T_{nk}$  ( $k = 1, 2, \dots, 4^n$ ) are the descendants of the  $T_{(n-1)k}$  ( $k = 1, 2, \dots, 4^{n-1}$ ) and are labeled in such a way that the points

$$\alpha_{nk} = \min \{x: (x, 1/2) \in T_{nk}\}$$

satisfy  $\alpha_{nk} < \alpha_{n(k+1)}$  for  $k = 1, 2, \dots, 4^n - 1$ .

If for each  $n$  we set

$$\alpha_{n(4n+1)} = \max \{x: (x, 1/2) \in T_{n4^n}\},$$

then it follows from (A) in §1 that both

$$T_{nk}(1/2) = \{(x, 1/2): \alpha_{nk} \leq x \leq \alpha_{n(k+1)}\}$$

for each  $k = 1, 2, \dots, 4^n$ , and for each  $n$

$$T_\xi(1/2) = \bigcup_{k=1}^{4^n} T_{nk}(1/2).$$

Now, for each  $n$ , let  $K_n = \bigcup_{k=1}^{4^n} T_{nk}$  and define the set

$$K_\xi = \bigcap_{n=1}^{\infty} K_n.$$

The following properties of the set  $K_\xi$  were established by Kahane [5]:

(I)  $K_\xi$  is a compact set of 2-dimensional measure zero.

(II) Let any segment  $\tau$  that extends from one base of  $T_\xi$  to the other be called “admissible”. Then the following two properties hold:

- (i) There are at most two admissible line segments in  $K_\xi$  that pass through any given point  $z \in T_\xi(1/2)$ ; in fact, precisely one such segment passes through each  $z \in T_\xi(1/2)$  that is not one of the  $\alpha_{nk}$ .
- (ii) For each  $\lambda \in [\xi - 1, \xi + 1/2] = \text{proj}[T_\xi]$ , there exists at least one point  $z \in T_\xi(1/2)$  through which there passes an admissible line segment  $\tau$  in  $K_\xi$  with  $\text{proj}[\tau] = \lambda$ ; in fact, for all but countably many such  $\lambda$ , there exists only one such  $z$ .

We now introduce a set-valued function  $\Lambda$  defined on the subsets  $A$  of  $T_\xi(1/2)$  as follows:

$$\Lambda(A) = \{ \text{proj}[\tau] : \tau \subset K_\xi \text{ is an admissible segment with } A \cap \tau \neq \emptyset \}.$$

Then  $K_\xi$  has the two additional properties:

(III)  $\Lambda(A)$  is nowhere dense whenever  $A$  is.

(IV)  $\Lambda(A)$  is of linear (Lebesgue) measure zero whenever  $A$  is.

Proof of (III). Let  $A$  be a nowhere dense subset of  $T_\xi(1/2)$ , and suppose  $\Lambda(A)$  is dense on some subinterval  $I$  of the interval  $\text{proj}[T_\xi]$ . Then, according to (B) in §1, there exist integers  $n$  and  $k$  such that  $\Lambda(A)$  is dense on  $\text{proj}[T_{nk}]$ . Since  $A$  is nowhere dense, there exist integers  $n'$  and  $k'$  such that  $T_{n'k'}(1/2) \subset T_{nk}(1/2)$  and  $A \cap T_{n'k'} = \emptyset$ . By (B) of §1, it follows that

$$\Lambda(A) \cap \text{int}(\text{proj}[T_{n'k'}]) = \emptyset$$

(int  $\equiv$  interior). This contradicts the fact that  $\Lambda(A)$  is dense on  $\text{proj}[T_{nk}]$ , and property (III) is proved.

For the remainder of this paper, we use  $\mu(A)$  and  $\mu^*(A)$  to denote respectively the Lebesgue measure and the Lebesgue outer measure of the linear set  $A$ .

Proof of (IV). Let  $A \subset T_\xi(1/2)$  and suppose  $\mu(A) = 0$ . If we set

$$\hat{A} = A - \{ \alpha_{nk} : n = 1, 2, \dots; k = 1, 2, \dots, 4^n + 1 \},$$

then in view of II (i) it suffices to show that the corresponding set  $\Lambda(\hat{A})$  has linear measure zero.

Let  $\mathcal{I}$  be the collection of all closed intervals on  $T_\xi(1/2)$  of the form  $[\alpha_{nk}, \alpha_{n(k+1)}]$ . Then it is easy to verify that

$$\mu(\hat{A}) = \inf \sum_{j=1}^{\infty} \mu(I_j),$$

where the inf is taken over all sequences  $\{I_j\}$  of intervals in  $\mathcal{I}$  covering  $\hat{A}$ .

Let  $\varepsilon > 0$ . Then there is a sequence  $\{I_j\}$  of intervals in  $\mathcal{I}$  that cover  $\hat{A}$  such that

$$\sum_{j=1}^{\infty} \mu(I_j) < \varepsilon.$$

For each index  $j$ , there exist integers  $n_j$  and  $k_j$  such that

$$I_j = T_{n_j k_j}(1/2);$$

hence, in view of (B) in §1, we have

$$A(\hat{A}) \subset \bigcup_{j=1}^{\infty} \text{proj}[T_{n_j k_j}].$$

Furthermore, combining (A) and (B) of §1, we obtain

$$\mu(\text{proj}[T_{n_j k_j}]) = 2\mu(I_j) \quad (j = 1, 2, \dots).$$

Therefore,

$$\mu^*(A(\hat{A})) \leq \sum_{j=1}^{\infty} \mu(\text{proj}[T_{n_j k_j}]) = 2 \sum_{j=1}^{\infty} \mu(I_j) < 2\varepsilon,$$

and property (IV) is proved.

### §3. PROOF OF THEOREM 1

For each integer  $n$ , let  $\xi_n = 1 + 3n/2$  and set

$$A_n = K_{\xi_n} \cap \{(x, y): \xi_n/2 \leq x \leq \xi_n/2 + 3/4 \text{ and } 1/2 < y \leq 1\}.$$

Then set

$$A_n^* = \{z - (1+i)/2: z \in A_n\} \quad (i = \sqrt{-1}),$$

and define the set

$$A = \bigcup \{A_n^*: n = 0, \pm 1, \pm 2, \dots\}.$$

Let  $\{z_n\}_{n=1}^{\infty}$  be a sequence of points in  $H - A$  whose derived set is  $R$ . Define the function  $f_0$  on  $A \cup \{z_n\}_{n=1}^{\infty}$  by

$$f_0(z) = \begin{cases} 0 & \text{for } z \in \Delta \\ 1 & \text{for } z = z_n (n = 1, 2, \dots) \end{cases}.$$

Then let  $f$  be a continuous extension of  $f_0$  to all of  $H$ . We now show that  $f$  is the desired function.

Let  $Q$  be a residual subset of  $R$ . Then, for each integer  $n$ , the set  $Q_n = Q \cap [3n/4, 3(n+1)/4]$  is a residual subset of the closed interval  $[3n/4, 3(n+1)/4]$ . As a consequence of (III) for  $\xi = \xi_n$ , there exists a residual set of directions  $\Theta_n \subset \{ \theta : 3n/2 \leq \cot \theta \leq 3(n+1)/2 \}$  such that, for each  $\theta \in \Theta_n$ , there exists a segment in  $\Delta$  emanating from a point of  $Q_n$  and having the direction  $\theta$ . Therefore, the set  $\bigcap_{x \in Q} \Theta(x)$  is of the first category on the set  $\{ \theta : 3n/2 \leq \cot \theta \leq 3(n+1)/2 \}$  for each integer  $n$ , and the theorem is proved.

#### §4. AN ESSENTIAL CLUSTER SET EXAMPLE

If  $f$  is a measurable function from  $H$  to  $W$ , then the *essential cluster set*  $C_e(f, x)$  of  $f$  at  $x$  is defined as the set of all values  $w \in W$  for which the upper density of  $f^{-1}(U)$  at  $x$  is positive for every open set  $U$  containing  $w$ ; the *essential cluster set*  $C_e(f, x, \theta)$  of  $f$  at  $x$  in the direction  $\theta$  is the set of all values  $w \in W$  for which the upper density of  $f^{-1}(U)$  along the ray at  $x$  having direction  $\theta$  is positive for every open set  $U$  containing  $w$ . As a supplement to a result of Casper Goffman and W. T. Sledd [4, Theorem 2], the present authors [1] proved the following result concerning the set

$$\Theta^*(x) = \{ \theta : C_e(f, x) \subset C_e(f, x, \theta) \} \quad (x \in R).$$

**THEOREM B.E.H.** *If  $f: H \rightarrow W$  is measurable, then  $\mu(\Theta^*(x)) = \pi$  for almost every and nearly every  $x \in R$ ; furthermore, if  $f$  is continuous, then  $\Theta^*(x)$  is residual for almost every and nearly every  $x \in R$ .*

Again, a natural question to ask is whether or not, for a given function  $f$ , there exists a “large” set of directions  $\Theta^*$  such that  $\Theta^* \subset \Theta^*(x)$  for a “large” set of points  $x \in R$ . As a partial answer, we prove

**THEOREM 2.** *There exists a continuous  $f: H \rightarrow W$  such that the intersection  $\bigcap_{x \in Q} \Theta^*(x)$  is (a) of the first category if  $Q \subset R$  is residual, and (b) of measure zero if  $Q \subset R$  is of full measure.*

*Proof.* Let  $\Delta$  be as in the proof of Theorem 1, and let  $S$  be a closed subset of  $H - \Delta$  that has metric density 1 at each  $x \in R$ . Let  $f$  be a continuous function on  $H$  with  $f(\Delta) = \{0\}$  and  $f(S) = \{1\}$ . Then the proof of (a) is completely analogous to the proof of Theorem 1, and the proof of (b) follows the same line with property (IV) used in place of property (III).

#### REFERENCES

- [1] BELNA, C. L., M. J. EVANS and P. D. HUMKE. Most directional cluster sets have common values. *Fund. Math. To appear.*
- [2] BRUCKNER, A. M. and Casper GOFFMAN. The boundary behavior of real functions in the upper half plane. *Rev. Roumaine Pures Appl.*, XI (1966), pp. 507-518.
- [3] COLLINGWOOD, E. F. Cluster sets and prime ends. *Ann. Acad. Sci. Fenn. Ser. AI*, No. 250/6 (1958), 12 pp.
- [4] GOFFMAN, Casper and W. T. SLEDD. Essential cluster sets. *J. London Math. Soc.* (2), 1 (1969), pp. 295-302.
- [5] KAHANE, J.-P. Trois notes sur les ensembles parfaits linéaires. *Enseignement Math.* 15 (1969), pp. 185-192.

(Reçu le 3 avril 1976)

C. L. Belna, M. J. Evans and P. D. Humke

Department of Mathematics  
Western Illinois University  
Macomb, Illinois 61455  
U.S.A.

vide-leer-empty