

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: DIRECTIONAL CLUSTER SET EXAMPLE
Autor: Belna, C. L. / Evans, M. J. / Humke, P. D.
Kapitel: §0. Introduction
DOI: <https://doi.org/10.5169/seals-48186>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A DIRECTIONAL CLUSTER SET EXAMPLE

by C. L. BELNA, M. J. EVANS and P. D. HUMKE

§0. INTRODUCTION

Let f be a mapping from the open upper half plane H into the Riemann sphere W . For each point x on the real line R , let $C(f, x)$ and $C(f, x, \theta)$ denote respectively the total cluster set of f at x and the cluster set of f at x in the direction θ ($0 < \theta < \pi$); then let $\Theta(x)$ denote the set of directions θ for which $C(f, x, \theta) = C(f, x)$. E. F. Collingwood [3, Theorem 2 combined with Theorem 3] established the following result.

THEOREM C. *Let $f: H \rightarrow W$ be continuous. Then the set $\Theta(x)$ is residual at each point x of a residual subset S of R .*

A. M. Bruckner and Casper Goffman [2, p. 510] raised the question as to whether or not there exists a residual set of directions Θ such that $\Theta \subset \Theta(x)$ for each $x \in S$. Here we prove

THEOREM 1. *There exists a continuous $f: H \rightarrow W$ such that $\bigcap_{x \in Q} \Theta(x)$ is a first category set of directions for each residual subset Q of R .*

To construct this function (§3), we use certain sets of J.-P. Kahane [5] as building blocks¹). Two important properties of these sets are established in §2, and the necessary technical preliminaries are presented in §1. Finally, in §4 we present an example concerning essential directional cluster sets.

§1. (1/2)-TRAPEZOIDS AND THEIR FOUR DESCENDANTS

By a (1/2)-trapezoid we mean any closed trapezoid T having bases L and L' which lie respectively on the lines $y = 0$ and $y = 1$ and for which $|L| = 2|L'|$. (Here and throughout this paper, we use $|\tau|$ to denote the length of the line segment τ .) For each (1/2)-trapezoid T , we set

$$T(1/2) = \{z \in T: \text{Im}(z) = 1/2\}.$$

¹) The authors wish to thank Professor John R. Kinney for bringing this paper of J.-P. Kahane to their attention.