Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: DIRECTIONAL CLUSTER SET EXAMPLE
Autor: Belna, C. L. / Evans, M. J. / Humke, P. D.
Kapitel: §2. Kahane's set K_ξ.
DOI: https://doi.org/10.5169/seals-48186

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 07.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

§2. Kahane's set K_ξ.

Let ξ be a real number, and let T_ξ be the (1/2)-trapezoid with bases

$$L = \{(x, 0): 0 \leqslant x \leqslant 1\} \quad \text{and} \quad L' = \{(x', 1): \xi \leqslant x' \leqslant \xi + 1/2\}.$$

Let T_{11}, T_{12}, T_{13}, and T_{14} be the four descendants of T_ξ labeled in such a way that the points

$$\alpha_{1k} = \min\{x: (x, 1/2) \in T_{1k}\}$$

satisfy $\alpha_{1k} < \alpha_{1(k+1)}$ for $k = 1, 2$, and 3. Then let T_{2k} $(k=1, 2, ..., 4^2)$ be the 4^2 descendants of the trapezoids T_{1k} $(k=1, 2, 3, 4)$ labeled in such a way that the points

$$\alpha_{2k} = \min\{x: (x, 1/2) \in T_{2k}\}$$

satisfy $\alpha_{2k} < \alpha_{2(k+1)}$ for $k = 1, 2, ..., 4^2-1$. Continuing inductively we arrive at the collection of (1/2)-trapezoids

$$T_{nk} \quad \text{for} \quad n = 1, 2, \ldots \quad \text{and} \quad k = 1, 2, \ldots, 4^n$$

where the T_{nk} $(k=1, 2, ..., 4^n)$ are the descendants of the $T_{(n-1)k}$ $(k=1, 2, ..., 4^{n-1})$ and are labeled in such a way that the points

$$\alpha_{nk} = \min\{x: (x, 1/2) \in T_{nk}\}$$

satisfy $\alpha_{nk} < \alpha_{n(k+1)}$ for $k = 1, 2, ..., 4^n-1$.

If for each n we set

$$\alpha_{n(4^n+1)} = \max\{x: (x, 1/2) \in T_{n4^n}\},$$

then it follows from (A) in §1 that both

$$T_{nk}(1/2) = \{(x, 1/2): \alpha_{nk} \leqslant x \leqslant \alpha_{n(k+1)}\}$$

for each $k = 1, 2, ..., 4^n$, and for each n

$$T_\xi(1/2) = \bigcup_{k=1}^{4^n} T_{nk}(1/2).$$

Now, for each n, let $K_n = \bigcup_{k=1}^{4^n} T_{nk}$ and define the set

$$K_\xi = \bigcap_{n=1}^{\infty} K_n.$$

The following properties of the set K_ξ were established by Kahane [5]:

(I) K_ξ *is a compact set of* 2-*dimensional measure zero.*

(II) *Let any segment* τ *that extends from one base of* T_ξ *to the other be called "admissible". Then the following two properties hold:*

(i) *There are at most two admissible line segments in* K_ξ *that pass through any given point* $z \in T_\xi(1/2)$; *in fact, precisely one such segment passes through each* $z \in T_\xi(1/2)$ *that is not one of the* α_{nk}.

(ii) *For each* $\lambda \in [\xi - 1, \xi + 1/2] = \text{proj}\,[T_\xi]$, *there exists at least one point* $z \in T_\xi(1/2)$ *through which there passes an admissible line segment* τ *in* K_ξ *with* $\text{proj}\,[\tau] = \lambda$; *in fact, for all but countably many such* λ, *there exists only one such* z.

We now introduce a set-valued function Λ defined on the subsets A of $T_\xi(1/2)$ as follows:

$$\Lambda(A) = \{\, \text{proj}\,[\tau] : \tau \subset K_\xi \text{ is an admissible segment with } A \cap \tau \neq \phi \,\}.$$

Then K_ξ has the two additional properties:

(III) $\Lambda(A)$ *is nowhere dense whenever* A *is.*

(IV) $\Lambda(A)$ *is of linear (Lebesgue) measure zero whenever* A *is.*

Proof of (III). Let A be a nowhere dense subset of $T_\xi(1/2)$, and suppose $\Lambda(A)$ is dense on some subinterval I of the interval proj $[T_\xi]$. Then, according to (B) in §1, there exist integers n and k such that $\Lambda(A)$ is dense on proj $[T_{nk}]$. Since A is nowhere dense, there exist integers n' and k' such that $T_{n'k'}(1/2) \subset T_{nk}(1/2)$ and $A \cap T_{n'k'} = \phi$. By (B) of §1, it follows that

$$\Lambda(A) \cap \text{int}\,(\text{proj}\,[T_{n'k'}]) = \phi$$

(int $\equiv$ interior). This contradicts the fact that $\Lambda(A)$ is dense on proj $[T_{nk}]$, and property (III) is proved.

For the remainder of this paper, we use $\mu(A)$ and $\mu^*(A)$ to denote respectively the Lebesgue measure and the Lebesgue outer measure of the linear set A.

Proof of (IV). Let $A \subset T_\xi(1/2)$ and suppose $\mu(A) = 0$. If we set

$$\hat{A} = A - \{\, \alpha_{nk} : n = 1, 2, \ldots;\ k = 1, 2, \ldots, 4^n + 1 \,\},$$

then in view of II (i) it suffices to show that the corresponding set $\Lambda(\hat{A})$ has linear measure zero.

Let $\mathscr{I}$ be the collection of all closed intervals on $T_\xi(1/2)$ of the form $[\alpha_{nk}, \alpha_{n(k+1)}]$. Then it is easy to verify that

$$\mu(\hat{A}) = \inf \sum_{j=1}^{\infty} \mu(I_j),$$

where the inf is taken over all sequences $\{I_j\}$ of intervals in $\mathscr{I}$ covering $\hat{A}$.

Let $\varepsilon > 0$. Then there is a sequence $\{I_j\}$ of intervals in $\mathscr{I}$ that cover $\hat{A}$ such that

$$\sum_{j=1}^{\infty} \mu(I_j) < \varepsilon.$$

For each index j, there exist integers n_j and k_j such that

$$I_j = T_{n_j k_j}(1/2);$$

hence, in view of (B) in §1, we have

$$\Lambda(\hat{A}) \subset \bigcup_{j=1}^{\infty} \operatorname{proj}[T_{n_j k_j}].$$

Furthermore, combining (A) and (B) of §1, we obtain

$$\mu(\operatorname{proj}[T_{n_j k_j}]) = 2\mu(I_j) \quad (j = 1, 2, \ldots).$$

Therefore,

$$\mu^*(\Lambda(\hat{A})) \leqslant \sum_{j=1}^{\infty} \mu(\operatorname{proj}[T_{n_j k_j}]) = 2 \sum_{j=1}^{\infty} \mu(I_j) < 2\varepsilon,$$

and property (IV) is proved.

§3. Proof of Theorem 1

For each integer n, let $\xi_n = 1 + 3n/2$ and set

$$\Delta_n = K_{\xi_n} \cap \{(x, y): \xi_n/2 \leqslant x \leqslant \xi_n/2 + 3/4 \quad \text{and} \quad 1/2 < y \leqslant 1\}.$$

Then set

$$\Delta_n^* = \{z - (1+i)/2 : z \in \Delta_n\} \quad (i = \sqrt{-1}),$$

and define the set

$$\Delta = \cup \{\Delta_n^* : n = 0, \pm 1, \pm 2, \ldots\}.$$

Let $\{z_n\}_{n=1}^{\infty}$ be a sequence of points in $H - \Delta$ whose derived set is R. Define the function f_0 on $\Delta \cup \{z_n\}_{n=1}^{\infty}$ by