Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: DIRECTIONAL CLUSTER SET EXAMPLE
Autor: Belna, C. L. / Evans, M. J. / Humke, P. D.
Kapitel: §3. Proof of Theorem 1
DOI: https://doi.org/10.5169/seals-48186

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 21.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Let $\mathscr{I}$ be the collection of all closed intervals on $T_\xi(1/2)$ of the form $[\alpha_{nk}, \alpha_{n(k+1)}]$. Then it is easy to verify that

$$\mu(\hat{A}) = \inf \sum_{j=1}^{\infty} \mu(I_j),$$

where the inf is taken over all sequences $\{I_j\}$ of intervals in $\mathscr{I}$ covering $\hat{A}$.

Let $\varepsilon > 0$. Then there is a sequence $\{I_j\}$ of intervals in $\mathscr{I}$ that cover $\hat{A}$ such that

$$\sum_{j=1}^{\infty} \mu(I_j) < \varepsilon.$$

For each index j, there exist integers n_j and k_j such that

$$I_j = T_{n_j k_j}(1/2);$$

hence, in view of (B) in §1, we have

$$\Lambda(\hat{A}) \subset \bigcup_{j=1}^{\infty} \operatorname{proj}[T_{n_j k_j}].$$

Furthermore, combining (A) and (B) of §1, we obtain

$$\mu(\operatorname{proj}[T_{n_j k_j}]) = 2\mu(I_j) \quad (j = 1, 2, \ldots).$$

Therefore,

$$\mu^*(\Lambda(\hat{A})) \leqslant \sum_{j=1}^{\infty} \mu(\operatorname{proj}[T_{n_j k_j}]) = 2 \sum_{j=1}^{\infty} \mu(I_j) < 2\varepsilon,$$

and property (IV) is proved.

§3. Proof of Theorem 1

For each integer n, let $\xi_n = 1 + 3n/2$ and set

$$\Delta_n = K_{\xi_n} \cap \{(x, y): \xi_n/2 \leqslant x \leqslant \xi_n/2 + 3/4 \quad \text{and} \quad 1/2 < y \leqslant 1\}.$$

Then set

$$\Delta_n^* = \{z - (1+i)/2 : z \in \Delta_n\} \quad (i = \sqrt{-1}),$$

and define the set

$$\Delta = \cup \{\Delta_n^* : n = 0, \pm 1, \pm 2, \ldots\}.$$

Let $\{z_n\}_{n=1}^{\infty}$ be a sequence of points in $H - \Delta$ whose derived set is R. Define the function f_0 on $\Delta \cup \{z_n\}_{n=1}^{\infty}$ by

$$f_0(z) = \begin{cases} 0 & \text{for} \quad z \in \Delta \\ 1 & \text{for} \quad z = z_n \, (n = 1, 2, \ldots) . \end{cases}$$

Then let f be a continuous extension of f_0 to all of H. We now show that f is the desired function.

Let Q be a residual subset of R. Then, for each integer n, the set $Q_n = Q \cap [3n/4, 3(n+1)/4]$ is a residual subset of the closed interval $[3n/4, 3(n+1)/4]$. As a consequence of (III) for $\xi = \xi_n$, there exists a residual set of directions $\Theta_n \subset \{\theta : 3n/2 \leqslant \cot\theta \leqslant 3(n+1)/2\}$ such that, for each $\theta \in \Theta_n$, there exists a segment in Δ emanating from a point of Q_n and having the direction θ. Therefore, the set $\bigcap_{x \in Q} \Theta(x)$ is of the first category on the set $\{\theta : 3n/2 \leqslant \cot\theta \leqslant 3(n+1)/2\}$ for each integer n, and the theorem is proved.

§4. An Essential Cluster Set Example

If f is a measurable function from H to W, then the *essential cluster set* $C_e(f, x)$ *of* f *at* x is defined as the set of all values $w \in W$ for which the upper density of $f^{-1}(U)$ at x is positive for every open set U containing w; the *essential cluster set* $C_e(f, x, \theta)$ *of* f *at* x *in the direction* θ is the set of all values $w \in W$ for which the upper density of $f^{-1}(U)$ along the ray at x having direction θ is positive for every open set U containing w. As a supplement to a result of Casper Goffman and W. T. Sledd [4, Theorem 2], the present authors [1] proved the following result concerning the set

$$\Theta^*(x) = \{\theta : C_e(f, x) \subset C_e(f, x, \theta)\} \quad (x \in R) .$$

Theorem B.E.H. *If* $f : H \to W$ *is measurable, then* $\mu(\Theta^*(x)) = \pi$ *for almost every and nearly every* $x \in R$; *furthermore, if* f *is continuous, then* $\Theta^*(x)$ *is residual for almost every and nearly every* $x \in R$.

Again, a natural question to ask is whether or not, for a given function f, there exists a "large" set of directions Θ^* such that $\Theta^* \subset \Theta^*(x)$ for a "large" set of points $x \in R$. As a partial answer, we prove

Theorem 2. *There exists a continuous* $f : H \to W$ *such that the intersection* $\bigcap_{x \in Q} \Theta^*(x)$ *is* (a) *of the first category if* $Q \subset R$ *is residual, and* (b) *of measure zero if* $Q \subset R$ *is of full measure.*