Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 23 (1977)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS

Autor: Vitushkin, A. G.

Kapitel: §3. Theorem on superpositions of smooth functions

DOI: https://doi.org/10.5169/seals-48931

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 09.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

§ 3. *Theorem on superpositions of smooth functions*

We will denote by $C_s(\mathscr{I}^n)$ the space of n times differentiable functions of n variables defined on the cube $\mathscr{I}^n$ with the norm

$$\|f\| = \sum_{p=1}^{s} \sum_{k_1+k_2+\ldots+k_n=p} \max_{x \in \mathscr{I}^n} \left| \frac{\partial^{k_1+\ldots+k_n} f(x)}{\partial x_1^{k_1} \ldots \partial x_n^{k_n}} \right|$$

THEOREM 2.3.1. *Let the numbers* $s \geqslant 1$, $s' \geqslant 1$ *and natural* n *and* n' *be such that* $\frac{n}{s} > \frac{n'}{s'}$. *Then the set of functions from* $C_s(\mathscr{I}^n)$ *not representable on* $\mathscr{I}^n$ *by superpositions of* S' *times differentiable functions of* n' *variables is a set of second category.*

The space $C_s(\mathscr{I}^n)$ is complete and consequently the set mentioned in the theorem is not empty. The theorem is true for any $s \geqslant 1$, $s' \geqslant 1$ but we will assume for simplicity that s and s' are integers.

LEMMA 2.3.1. *Let* f *and* f' *be* q*-fold superpositions composed of the functions* $\{\varphi^p_{\alpha_1,\ldots,\alpha_p}\}$ *and* $\{\tilde{\varphi}^p_{\alpha_1,\ldots,\alpha_p},\}$ *where all functions composing the superpositions satisfy the condition* Lip 1 *with the constant* L *and for any collection* $p, \alpha_1, \ldots, \alpha_p$

$$\max \left| \varphi^p_{\alpha_1,\ldots,\alpha_p} - \tilde{\varphi}^p_{\alpha_1,\ldots\alpha_p} \right| \leqslant \varepsilon$$

Then

$$\max_{x \in \mathscr{I}^n} \left| f(x) - \tilde{f}(x) \right| \leqslant (L+1)^q \ \varepsilon$$

The lemma can easily be proved by induction in q.

LEMMA 2.3.2. *Let* Ω *be an open subset of* $C_s(\mathscr{I}^n)$ *and* $\Omega^* \subset C(\mathscr{I}^n)$. *If every* $f \in \Omega$ *allows uniform approximations on* $\mathscr{I}^n$ *with any accuracy by functions from* Ω^*, *i.e. the closure of* Ω^* *contains* Ω, *then* $H_\varepsilon(\Omega^*) \geqslant C\left(\frac{1}{\varepsilon}\right)^{n/s}$, *where* $C > 0$ *is independent of* ε.

The lemma is easily reduced to lemma 2.2.1 and lemma 2.2.2.

We denote by Ω_k the set of all functions of $C(\mathscr{I}^n)$ which are k-fold superpositions composed of s' times differentiable functions of n' variables with partial derivatives bounded by the same constant k.

LEMMA 2.3.3. *If* $\frac{n}{s} > \frac{n'}{s'}$ *then for any natural* k *the set* $\Omega_k \cap C_s(\mathscr{I}^n)$ *is nowhere dense in* $C_s(\mathscr{I}^n)$.

By lemma 2.3.1 and the theorem 2.2.1 for any natural k $H_\varepsilon(\Omega_k) \leqslant C\left(\frac{1}{\varepsilon}\right)^{n'/s'}$, where C does not depend on ε. Hence, it follows from the inequality $\frac{n}{s} > \frac{n'}{s'}$ and lemma 2.3.2 that the set $\Omega_k \cap C_s(\mathscr{I}^n)$ is nowhere dense in $C_s(\mathscr{I}^n)$.

Now to prove the theorem we have to notice only that the set of functions from $C_s(\mathscr{I}^n)$ representable by superpositions coincides with $\bigcup_{k=1}^{\infty} (\Omega_k \cap C_s(\mathscr{I}^n))$. By lemma 2.3.3 the sets $\{\Omega_k \cap C_s(\mathscr{I}^n)\}$ are nowhere dense and consequently the set of not representable functions is a set of second category.

CHAPTER 3. — SUPERPOSITIONS OF CONTINUOUS FUNCTIONS

In this chapter we present the proof of the theorem of Kolmogorov given by Kahane [36]. This proof which is based on Baire's theory contains a minimum of concrete constructions and shows that there exists a wide choice of inner functions for Kolmogorov's formula.

§ 1. *Certain improvements of Kolmogorov's theorem*

By the theorem of Kolmogorov any function defined and continuous on the cube $\mathscr{I}^n$ can be represented as

$$f(x_1, \ldots, x_n) = \sum_{q=1}^{2n+1} g_q\Big(\sum_{p=1}^{n} \varphi_{p,q}(x_p)\Big),$$

where $\{\varphi_{p,q}\}$ are specially chosen continuous and monotonic functions which do not depend on f, and where $\{g_q\}$ are continuous functions.

Lorentz [12] has noticed that in the theorem of Kolmogorov the functions $\{g_q\}$ can be chosen independently of q. In fact, by adding constants to the functions $t_q = \sum_{p=1}^{n} \varphi_{p,q}(x_p)$ $(q = 1, \ldots, 2n+1)$ one can make the ranges