

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS

Autor: Vitushkin, A. G.

Kapitel: Chapter 3. — Superpositions of continuous functions

DOI: <https://doi.org/10.5169/seals-48931>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LEMMA 2.3.3. If $\frac{n}{s} > \frac{n'}{s'}$ then for any natural k the set $\Omega_k \cap C_s(\mathcal{I}^n)$ is nowhere dense in $C_s(\mathcal{I}^n)$.

By lemma 2.3.1 and the theorem 2.2.1 for any natural k $H_\varepsilon(\Omega_k) \leq C \left(\frac{1}{\varepsilon}\right)^{n'/s'}$, where C does not depend on ε . Hence, it follows from the inequality $\frac{n}{s} > \frac{n'}{s}$ and lemma 2.3.2 that the set $\Omega_k \cap C_s(\mathcal{I}^n)$ is nowhere dense in $C_s(\mathcal{I}^n)$.

Now to prove the theorem we have to notice only that the set of functions from $C_s(\mathcal{I}^n)$ representable by superpositions coincides with $\bigcup_{k=1}^{\infty} (\Omega_k \cap C_s(\mathcal{I}^n))$. By lemma 2.3.3 the sets $\{\Omega_k \cap C_s(\mathcal{I}^n)\}$ are nowhere dense and consequently the set of not representable functions is a set of second category.

CHAPTER 3. — SUPERPOSITIONS OF CONTINUOUS FUNCTIONS

In this chapter we present the proof of the theorem of Kolmogorov given by Kahane [36]. This proof which is based on Baire's theory contains a minimum of concrete constructions and shows that there exists a wide choice of inner functions for Kolmogorov's formula.

§ 1. *Certain improvements of Kolmogorov's theorem*

By the theorem of Kolmogorov any function defined and continuous on the cube \mathcal{I}^n can be represented as

$$f(x_1, \dots, x_n) = \sum_{q=1}^{2n+1} g_q \left(\sum_{p=1}^n \varphi_{p,q}(x_p) \right),$$

where $\{\varphi_{p,q}\}$ are specially chosen continuous and monotonic functions which do not depend on f , and where $\{g_q\}$ are continuous functions.

Lorentz [12] has noticed that in the theorem of Kolmogorov the functions $\{g_q\}$ can be chosen independently of q . In fact, by adding constants to the functions $t_q = \sum_{p=1}^n \varphi_{p,q}(x_p)$ ($q = 1, \dots, 2n+1$) one can make the ranges

of the functions pairwise disjoint and consequently the functions $\{t_q\}$ can be considered as the restrictions of a single function $\{g_q\}$.

Sprecher [40] has shown that the functions $\{\varphi_{p,q}\}$ can be chosen in the form $\varphi_{p,q}(x_p) = \lambda_p \varphi_q(x_p)$ where $\{\lambda_p\}$ are constants and $\{\varphi_q\}$ -are continuous monotonic functions.

Thus any continuous function can be represented as

$$f(x_1, \dots, x_n) = \sum_{q=1}^{2n+1} g \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right),$$

where the constants $\{\lambda_p\}$ and the continuous monotone functions $\{\varphi_q\}$ do not depend on f , and where g is a continuous function.

Kahane [36] has shown that such a representation is possible with almost every collection of constants $\{\lambda_p\}$ and “quasi every” collection of continuous functions $\{\varphi_q\}$. The precise statement of this theorem will be given below. Here we consider some further results concerning the formula of Kolmogorov.

Doss [38] has shown that for any continuous monotonic functions $\varphi_{p,q}$ ($p=1, 2$; $q=1, 2, 3, 4$) there exists a continuous function $f(x_1, x_2)$ of two variables not representable as a superposition of the form $\sum_{q=1}^4 g_q \left(\sum_{p=1}^2 \varphi_{p,q}(x_p) \right)$, where $\{g_q\}$ are continuous functions.

Bassalygo [39] succeeded in showing that for any continuous functions $\varphi_i(x_1, x_2)$ ($i=1, 2, 3$) there exists a continuous function $f(x_1, x_2)$ that is not equal to any superposition of the form $\sum_{i=1}^3 g_i(\varphi_i(x_1, x_2))$, where $\{g_i\}$ are continuous functions.

Tihomirov showed that Kolmogorov's theorem can be generalized as follows: for any compact K of dimension n there exists a homeomorphic embedding $\Psi(x) = \{\Psi_1(x), \dots, \Psi_{2n+1}(x)\}$, $x \in K$ into $(2n+1)$ -dimensional euclidean space such that any continuous function $f(x)$ on K can be represented in the form $f(x) = \sum_{i=1}^{2n+1} g_i(\Psi_i(x))$, where $\{g_i\}$ are continuous functions of one variable.

In the same paper [36] Kahane has shown that there exist complex numbers λ_p ($p=1, \dots, n$) and complex valued functions φ_q ($q=1, \dots, 2n+1$) possessing the following properties.

1. The function φ_q is a monotonic continuous transformation of the real axis onto the circle $|t| = 1$ ($q=1, \dots, 2n+1$).

2. The function $t_q = \sum_{p=1}^n \lambda_p \varphi_q(x_p)$ maps the cube \mathcal{I}^n into the circle $|t| = 1$.

3. The transformation Ψ given by the equalities $t_q = \sum_{p=1}^n \lambda_p \varphi_q(x_p)$ ($q = 1, \dots, 2n+1$) is one-to-one on \mathcal{I}^n .

4. For any function f continuous on \mathcal{I}^n there exists a function $g(z)$ continuous on the disk $|z| \leq 1$, holomorphic inside that disk, and such that $f = \sum_{q=1}^{2n+1} g \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right)$.

The transformation Ψ gives an embedding of the cube \mathcal{I}^n into the torus $|t| = 1$ ($q = 1, \dots, 2n+1$) such that any function continuous on the cube $\tilde{\mathcal{I}}^n = \Psi(\mathcal{I}^n)$ is represented in the form $f(t_1, \dots, t_{2n+1}) = \sum_{q=1}^{2n+1} g(t_q)$, where g is a function holomorphic in the unit disk. This means in particular that any function continuous on $\tilde{\mathcal{I}}^n$ has an analytic extension to the polydisk $|t_q| \leq 1$ ($q = 1, \dots, 2n+1$).

§ 2. *The theorem of Kahane*

Let M be a complete metric space. We recall that a set is called a set of second category if it is the intersection of a countable family of open sets which are everywhere dense in M . By the theorem of Baire in a complete metric space no set of second category is empty. The massivity of such sets is characterized by the fact that the intersection of a countable family of sets of second category is again a set of second category and consequently is not empty.

We will say that a statement is true for quasi every element of M if it is true for a set of elements of second category.

Let us consider an example. Let Φ be the space with uniform norm consisting of all functions continuous and non-decreasing on the segment \mathcal{I}^1 ($0 \leq t \leq 1$). It can be shown easily that quasi every element of Φ is a strictly increasing function.

In fact, any strictly increasing function belongs to any set defined as $\varphi(r') < \varphi(r'')$, where $r' < r''$ are fixed rational numbers. Any set defined by an inequality of that type is open and everywhere dense in Φ , and the set of all such sets is countable.

Let \mathcal{I}^n be the cube $\{0 \leq x_i \leq 1, i = 1, \dots, n\}$; $C(\mathcal{I}^n)$ -the space of all functions continuous on \mathcal{I}^n with the uniform norm; Φ -the space of functions continuous and non-decreasing on the segment \mathcal{I}^1 (with the uniform norm); $\Phi^k = \Phi \times \dots \times \Phi$ the k -th power of the space Φ .

THEOREM 3.2.1. *Let $\lambda_p (p=1, \dots, n)$ be a collection of rationally independent constants. Then for quasi every collection $\{\varphi_1, \dots, \varphi_{2n+1}\} \in \Phi^{2n+1}$ it is true that any function $f \in C(\mathcal{I}^n)$ can be represented on \mathcal{I}^n in the form*

$$f(x) = \sum_{q=1}^{2n+1} g \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right),$$

where g is a continuous function.

§ 3. The main lemma

We fix a function $f \in C(\mathcal{I}^n)$, positive numbers $\lambda_p (p=1, \dots, n)$ and a positive ε . We will denote by Ω_f the set of all collections $\{\varphi_1, \dots, \varphi_{2n+1}\} \in \Phi^{2n+1}$ for each of which there exists a continuous function h such that $\|h\| \leq \|f\|$ and $\|f(x) - \sum_{q=1}^{2n+1} h \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right)\| < (1-\varepsilon) \|f\|$. The latter inequality is strict and consequently the set Ω_f is open.

The idea of the construction is contained in the following statement.

LEMMA 3.3.1. *If $\|f\| \neq 0$, the numbers $\{\lambda_p\}$ are rationally independent, and $0 < \varepsilon < \frac{1}{2n+2}$, then the corresponding set Ω_f is everywhere dense in Φ^{2n+1} .*

Proof. Let us fix an open set $\Omega \subset \Phi^{2n+1}$ and prove that $\Omega \cap \Omega_f$ is not empty. This will imply that Ω_f is everywhere dense in Φ^{2n+1} .

We choose a number $\delta > 0$ and denote by $\mathcal{I}_q(j)$ the segment defined by the inequality

$$q \cdot \delta + (2n+1)j \cdot \delta \leq t \leq q \cdot \delta + (2n+1)j\delta + 2n\delta \\ (q=1, \dots, 2n+1, j \text{ is an integer}).$$

The value δ will be determined below. Now we notice, firstly, that for any q the segments $\mathcal{I}_q(j) (j=0, \pm 1, \pm 2)$ are pairwise disjoint and every two consecutive segments are separated by an interval of length δ and, secondly, that, every point of the real axis belongs to at least $2n$ of the sets $\sum_j \mathcal{I}_q(j), (q=1, \dots, 2n+1)$.

We denote by $P_q(j_1, \dots, j_n)$ the cube

$$q\delta + (2n+1)j_k\delta \leq x_k \leq q\delta + (2n+1)j_k\delta + 2n\delta \quad (k=1, \dots, n).$$

We emphasise that every point $x \in \mathcal{I}^n$ belongs to at least $n+1$ of the sets

$\sum_{j_1, \dots, j_n} P_q(j_1, \dots, j_n)$ ($q=1, \dots, 2n+1$). We also remark that for any q the cubes $\{P_q(j_1, \dots, j_n)\}$ are pairwise disjoint.

We denote by Ω^* the subset of Φ^{2n+1} consisting of the collections $\varphi_1, \dots, \varphi_{2n+1}$ such that for every q the function φ_q is constant on every one of the segments $\{\mathcal{I}_q(j)\}$. We will assume that δ is so small that $\Omega^* \cap \Omega$ is not empty.

We choose a collection $\{\varphi_1, \dots, \varphi_{2n+1}\} \in \Omega^* \cap \Omega$. We will show that this collection belongs to Ω_f . We put $t_q \equiv \sum_{p=1}^n \lambda_p \varphi_q(x_p)$. Since the numbers $\{\lambda_p\}$ are rationally independent we can change the constants $\{\varphi_q(\mathcal{I}_q(j))\}$ slightly, so that the new values of $t_q(p_q(j_1, \dots, j_n))$ are pairwise different and the collection $\varphi_1, \dots, \varphi_{2n+1}$ remains in $\Omega^* \cap \Omega$.

We denote by $f_q(j_1, \dots, j_n)$ the value of the function f at the center of $P_q(j_1, \dots, j_n)$ and by h the function defined in the following way:

$h(t_q(j_1, \dots, j_n)) = \frac{1}{2n+1} f_q(j_1, \dots, j_n)$ outside the set $\cup_{q, j_1, \dots, j_n} t_q(j_1, \dots, j_n)$
the function h is defined in such a way that it is continuous on the whole real axis and $\|h\| \leq \frac{1}{2n+1} \|f\|$.

Now we estimate the function $|f - \sum_{q=1}^{2n+1} h(t_q)| = \left| \sum_{q=1}^{2n+1} \frac{f}{2n+1} - h(t_q) \right|$.
For any $x \in \mathcal{I}^n$, q, j_1, \dots, j_n

$$\begin{aligned} \left| \frac{f}{2n+1} - h(t_q) \right| &\leq \frac{1}{2n+1} \|f\| + \|h\| \leq \frac{1}{2n+1} \|f\| + \frac{1}{2n+1} \|f\| \\ &= \frac{2}{2n+1} \|f\|. \end{aligned}$$

If $x \in P_q(j_1, \dots, j_n)$, then

$$\begin{aligned} &\left| \frac{f}{2n+1} - h(t_q) \right| \\ &\leq \max_{q, j_1, \dots, j_n} \left| \max_{x \in P_q(j_1, \dots, j_n)} \frac{f(x)}{2n+1} - \min_{x \in P_q(j_1, \dots, j_n)} \frac{f(x)}{2n+1} \right| = \rho. \end{aligned}$$

We recall that every $x \in \mathcal{I}^n$ belongs to at least $n+1$ of the cubes $\{P_q(j_1, \dots, j_n)\}$. Hence

$$\left| f - \sum_{q=1}^{2n+1} h(t_q) \right| \leq (n+1)\rho + n \frac{2}{2n+1} \|f\|.$$

But $\lim_{\delta \rightarrow 0} \rho = 0$, consequently for sufficiently small δ and $\varepsilon < \frac{1}{2n+2}$

$$\left| f - \sum_{q=1}^{2n+1} h(t_q) \right| < (1-\varepsilon) \|f\|.$$

The lemma is proved.

§ 4. *The proof of the theorem*

We denote by F a countable set, everywhere dense in $C(\mathcal{I}^n)$. We choose ε satisfying the condition of lemma 3.3.1 and consider Ω_{f_k} ($f_k \in F$) corresponding to this ε and the collection λ_p mentioned in the theorem. The sets $\{\Omega_{f_k}\}$ are open and by lemma 3.3.1 they are everywhere dense in Φ^{2n+1} . Consequently, according to the definition, almost every element of Φ^{2n+1} belongs to $\Phi^* = \bigcap_{f_k \in F} \Omega_{f_k}$.

We fix a collection $\{\varphi_1, \dots, \varphi_{2n+1}\} \in \Phi^*$ and a function $f \in C(\mathcal{I}^n)$ and show that the desired representation of f takes place. If $f \equiv 0$ then as the function g we can take $g \equiv 0$. We will assume below that $f \not\equiv 0$. According to the definition of Ω_{f_k} there exists for any $f_k \in F$ a function h_k such that

$\left| f_k - \sum_{q=1}^{2n+1} h_k \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right) \right| \leq (1-\varepsilon) \|f_k\|$. The set F is everywhere dense in $C(\mathcal{I}^n)$. Consequently for any $f \in C(\mathcal{I}^n)$ ($f \not\equiv 0$) there exists $h = \gamma(f)$ such that

$$\left| f - \sum_{q=1}^{2n+1} h \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right) \right| < \left(1 - \frac{\varepsilon}{2} \right) \|f\|.$$

We define the sequence of functions $\chi_0, \chi_1, \chi_2, \dots$ by the recurrent equalities

$$\chi_0 = f, \quad \chi_{k+1} = \chi_k - \sum_{q=1}^{\infty} g_k \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right),$$

where $g_k = \gamma(\chi_k)$. The series $\sum_{k=0}^{\infty} g_k$ converges uniformly and consequently the function $g = \sum_{k=0}^{\infty} g_k$ is continuous and

$$f - \sum_{q=1}^{2n+1} g \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right) = 0.$$

The theorem is proved.