

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF
SUPERPOSITIONS AND RELATED TOPICS
Autor: Vitushkin, A. G.
Kapitel: §1. Certain improvements of Kolmogorov 's theorem
DOI: <https://doi.org/10.5169/seals-48931>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LEMMA 2.3.3. If $\frac{n}{s} > \frac{n'}{s'}$ then for any natural k the set $\Omega_k \cap C_s(\mathcal{J}^n)$ is nowhere dense in $C_s(\mathcal{J}^n)$.

By lemma 2.3.1 and the theorem 2.2.1 for any natural k $H_\varepsilon(\Omega_k) \leq C \left(\frac{1}{\varepsilon}\right)^{n'/s'}$, where C does not depend on ε . Hence, it follows from the inequality $\frac{n}{s} > \frac{n'}{s'}$ and lemma 2.3.2 that the set $\Omega_k \cap C_s(\mathcal{J}^n)$ is nowhere dense in $C_s(\mathcal{J}^n)$.

Now to prove the theorem we have to notice only that the set of functions from $C_s(\mathcal{J}^n)$ representable by superpositions coincides with $\bigcup_{k=1}^{\infty} (\Omega_k \cap C_s(\mathcal{J}^n))$. By lemma 2.3.3 the sets $\{\Omega_k \cap C_s(\mathcal{J}^n)\}$ are nowhere dense and consequently the set of not representable functions is a set of second category.

CHAPTER 3. — SUPERPOSITIONS OF CONTINUOUS FUNCTIONS

In this chapter we present the proof of the theorem of Kolmogorov given by Kahane [36]. This proof which is based on Baire's theory contains a minimum of concrete constructions and shows that there exists a wide choice of inner functions for Kolmogorov's formula.

§ 1. *Certain improvements of Kolmogorov's theorem*

By the theorem of Kolmogorov any function defined and continuous on the cube \mathcal{J}^n can be represented as

$$f(x_1, \dots, x_n) = \sum_{q=1}^{2n+1} g_q \left(\sum_{p=1}^n \varphi_{p,q}(x_p) \right),$$

where $\{\varphi_{p,q}\}$ are specially chosen continuous and monotonic functions which do not depend on f , and where $\{g_q\}$ are continuous functions.

Lorentz [12] has noticed that in the theorem of Kolmogorov the functions $\{g_q\}$ can be chosen independently of q . In fact, by adding constants to the functions $t_q = \sum_{p=1}^n \varphi_{p,q}(x_p)$ ($q = 1, \dots, 2n+1$) one can make the ranges

of the functions pairwise disjoint and consequently the functions $\{t_q\}$ can be considered as the restrictions of a single function $\{g_q\}$.

Sprecher [40] has shown that the functions $\{\varphi_{p,q}\}$ can be chosen in the form $\varphi_{p,q}(x_p) = \lambda_p \varphi_q(x_p)$ where $\{\lambda_p\}$ are constants and $\{\varphi_q\}$ are continuous monotonic functions.

Thus any continuous function can be represented as

$$f(x_1, \dots, x_n) = \sum_{q=1}^{2n+1} g \left(\sum_{p=1}^n \lambda_p \varphi_q(x_p) \right),$$

where the constants $\{\lambda_p\}$ and the continuous monotone functions $\{\varphi_q\}$ do not depend on f , and where g is a continuous function.

Kahane [36] has shown that such a representation is possible with almost every collection of constants $\{\lambda_p\}$ and “quasi every” collection of continuous functions $\{\varphi_q\}$. The precise statement of this theorem will be given below. Here we consider some further results concerning the formula of Kolmogorov.

Doss [38] has shown that for any continuous monotonic functions $\varphi_{p,q}$ ($p=1, 2; q=1, 2, 3, 4$) there exists a continuous function $f(x_1, x_2)$ of two variables not representable as a superposition of the form $\sum_{q=1}^4 g_q \left(\sum_{p=1}^2 \varphi_{p,q}(x_p) \right)$, where $\{g_q\}$ are continuous functions.

Bassalygo [39] succeeded in showing that for any continuous functions $\varphi_i(x_1, x_2)$ ($i=1, 2, 3$) there exists a continuous function $f(x_1, x_2)$ that is not equal to any superposition of the form $\sum_{i=1}^3 g_i(\varphi_i(x_1, x_2))$, where $\{g_i\}$ are continuous functions.

Tihomirov showed that Kolmogorov's theorem can be generalized as follows: for any compact K of dimension n there exists a homeomorphic embedding $\Psi(x) = \{\Psi_1(x), \dots, \Psi_{2n+1}(x)\}$, $x \in K$ into $(2n+1)$ -dimensional euclidean space such that any continuous function $f(x)$ on K can be represented in the form $f(x) = \sum_{i=1}^{2n+1} g_i(\Psi_i(x))$, where $\{g_i\}$ are continuous functions of one variable.

In the same paper [36] Kahane has shown that there exist complex numbers λ_p ($p=1, \dots, n$) and complex valued functions φ_q ($q=1, \dots, 2n+1$) possessing the following properties.

1. The function φ_q is a monotonic continuous transformation of the real axis onto the circle $|t| = 1$ ($q=1, \dots, 2n+1$).

2. The function $t_q = \sum_{p=1}^n \lambda_p \varphi_q(x_p)$ maps the cube \mathcal{J}^n into the circle $|t| = 1$.

3. The transformation Ψ given by the equalities $t_q = \sum_{p=1}^n \lambda_p \varphi_q(x_p)$ ($q=1, \dots, 2n+1$) is one-to-one on \mathcal{J}^n .

4. For any function f continuous on \mathcal{J}^n there exists a function $g(z)$ continuous on the disk $|z| \leq 1$, holomorphic inside that disk, and such that $f = \sum_{q=1}^{2n+1} g\left(\sum_{p=1}^n \lambda_p \varphi_q(x_p)\right)$.

The transformation Ψ gives an embedding of the cube \mathcal{J}^n into the torus $|t| = 1$ ($q=1, \dots, 2n+1$) such that any function continuous on the cube $\tilde{\mathcal{J}}^n = \Psi(\mathcal{J}^n)$ is represented in the form $f(t_1, \dots, t_{2n+1}) = \sum_{q=1}^{2n+1} g(t_q)$, where g is a function holomorphic in the unit disk. This means in particular that any function continuous on $\tilde{\mathcal{J}}^n$ has an analytic extension to the polydisk $|t_q| \leq 1$ ($q=1, \dots, 2n+1$).

§ 2. The theorem of Kahane

Let M be a complete metric space. We recall that a set is called a set of second category if it is the intersection of a countable family of open sets which are everywhere dense in M . By the theorem of Baire in a complete metric space no set of second category is empty. The massivity of such sets is characterized by the fact that the intersection of a countable family of sets of second category is again a set of second category and consequently is not empty.

We will say that a statement is true for quasi every element of M if it is true for a set of elements of second category.

Let us consider an example. Let Φ be the space with uniform norm consisting of all functions continuous and non-decreasing on the segment \mathcal{J}^1 ($0 \leq t \leq 1$). It can be shown easily that quasi every element of Φ is a strictly increasing function.

In fact, any strictly increasing function belongs to any set defined as $\varphi(r') < \varphi(r'')$, where $r' < r''$ are fixed rational numbers. Any set defined by an inequality of that type is open and everywhere dense in Φ , and the set of all such sets is countable.