Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 23 (1977)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS

Autor: Vitushkin, A. G.

Kapitel: §2. The theorem of Kahane

DOI: https://doi.org/10.5169/seals-48931

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

2. The function $t_q = \sum_{p=1}^{n} \lambda_p \varphi_q (x_p)$ maps the cube $\mathscr{I}^n$ into the circle $|t| = 1$.

3. The transformation Ψ given by the equalities $t_q = \sum_{p=1}^{n} \lambda_p \varphi_q (x_p)$ $(q = 1, \ldots, 2n+1)$ is one-to-one on $\mathscr{I}^n$.

4. For any function f continuous on $\mathscr{I}^n$ there exists a function $g(z)$ continuous on the disk $|z| \leqslant 1$, holomorphic inside that disk, and such that $f = \sum_{q=1}^{2n+1} g\left(\sum_{p=1}^{n} \lambda_p \varphi_q (x_p)\right)$.

The transformation Ψ gives an embedding of the cube $\mathscr{I}^n$ into the torus $|t| = 1$ $(q = 1, \ldots, 2n+1)$ such that any function continuous on the cube $\tilde{\mathscr{I}}^n = \Psi(\mathscr{I}^n)$ is represented in the form $f(t_1, \ldots, t_{2n+1}) = \sum_{q=1}^{2n+1} g(t_q)$, where g is a function holomorphic in the unit disk. This means in particular that any function continuous on $\tilde{\mathscr{I}}^n$ has an analytic extension to the polydisk $|t_q| \leqslant 1$ $(q = 1, \ldots, 2n+1)$.

§ 2. *The theorem of Kahane*

Let M be a complete metric space. We recall that a set is called a set of second category if it is the intersection of a countable family of open sets which are everywhere dense in M. By the theorem of Baire in a complete metric space no set of second category is empty. The massivity of such sets is characterized by the fact that the intersection of a countable family of sets of second category is again a set of second category and consequently is not empty.

We will say that a statement is true for quasi every element of M if it is true for a set of elements of second category.

Let us consider an example. Let Φ be the space with uniform norm consisting of all functions continuous and non-decreasing on the segment $\mathscr{I}^1$ $(0 \leqslant t \leqslant 1)$. It can be shown easily that quasi every element of Φ is a strictly increasing function.

In fact, any strictly increasing function belongs to any set defined as $\varphi(r') < \varphi(r'')$, where $r' < r''$ are fixed rational numbers. Any set defined by an inequality of that type is open and everywhere dense in Φ, and the set of all such sets is countable.

Let $\mathscr{I}^n$ be the cube $\{ 0 \leqslant x_i \leqslant 1,\ i = 1, \ldots, n \}$; $C(\mathscr{I}^n)$-the space of all functions continuous on $\mathscr{I}^n$ with the uniform norm; Φ-the space of functions continuous and non-decreasing on the segment $\mathscr{I}^1$ (with the uniform norm); $\Phi^k = \Phi \times \ldots \times \Phi$ the k-th power of the space Φ.

THEOREM 3.2.1. *Let λ_p $(p=1, \ldots, n)$ be a collection of rationally independent constants. Then for quasi every collection $\{ \varphi_1, \ldots, \varphi_{2n+1} \} \in \Phi^{2n+1}$ it is true that any function $f \in C(\mathscr{I}^n)$ can be represented on $\mathscr{I}^n$ in the form*

$$f(x) = \sum_{q=1}^{2n+1} g\Big(\sum_{p=1}^{n} \lambda_p \varphi_q(x_p) \Big),$$

where g is a continuous function.

§ 3. *The main lemma*

We fix a function $f \in C(\mathscr{I}^n)$, positive numbers λ_p $(p=1, \ldots, n)$ and a positive ε. We will denote by Ω_f the set of all collections $\{ \varphi_1, \ldots, \varphi_{2n+1} \} \in \Phi^{2n+1}$ for each of which there exists a continuous function h such that $\| h \| \leqslant \| f \|$ and $\| f(x) - \sum_{q=1}^{2n+1} h\big(\sum_{p=1}^{n} \lambda_p \varphi_q(x_p) \big) \| < (1-\varepsilon) \| f \|$. The latter inequality is strict and consequently the set Ω_f is open.

The idea of the construction is contained in the following statement.

LEMMA 3.3.1. *If $\| f \| \neq 0$, the numbers $\{ \lambda_p \}$ are rationally independent, and $0 < \varepsilon < \dfrac{1}{2n+2}$, than the corresponding set Ω_f is everywhere dense in Φ^{2n+1}.*

Proof. Let us fix an open set $\Omega \subset \Phi^{2n+1}$ and prove that $\Omega \cap \Omega_f$ is not empty. This will imply that Ω_f is everywhere dense in Φ^{2n+1}.

We choose a number $\delta > 0$ and denote by $\mathscr{I}_q(j)$ the segment defined by the inequality

$$q \cdot \delta + (2n+1) j \cdot \delta \leqslant t \leqslant q \cdot \delta + (2n+1) j\delta + 2n\delta$$
$$(q=1, \ldots, 2n+1,\ j \text{ is an integer}).$$

The value δ will be determined below. Now we notice, firstly, that for any q the segments $\mathscr{I}_q(j)$ $(j=0, \pm 1, \pm 2)$ are pairwise desjoint and every two consecutive segments are separated by an interval of length δ and, secondly, that, every point of the real axis belongs to at least $2n$ of the sets $\sum_j \mathscr{I}_q(j)$, $(q=1, \ldots, 2n+1)$.