

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS
Autor: Vitushkin, A. G.
Kapitel: §1. Notation
DOI: <https://doi.org/10.5169/seals-48931>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

CHAPTER 4. — LINEAR SUPERPOSITIONS

In this chapter we prove that there exist analytic functions which are not representable by means of linear superpositions of smooth functions of one variable.

§ 1. *Notation*

Throughout we assume that all the functions are defined and continuous for all values of the arguments. If we say that a function is continuously differentiable, we mean by this that its first partial derivatives are defined and continuous for all values of the arguments; $z = (x, y)$ is the point of the plane with coordinates x and y ; $\text{grad } [q(z)]$ is the gradient of the function $q(z)$, that is, the vector-function with coordinates $\frac{\partial q}{\partial x}$ and $\frac{\partial q}{\partial y}$;

$$D \left(\begin{array}{c} q_1, q_2 \\ x, y \end{array} \right) = \begin{vmatrix} \frac{\partial q_1}{\partial x} & \frac{\partial q_1}{\partial y} \\ \frac{\partial q_2}{\partial x} & \frac{\partial q_2}{\partial y} \end{vmatrix}$$

is the Jacobian of the pair of functions q_1 and q_2 .

$q(D)$ is the image of the set D under the mapping effected by the function $q(x, y)$; $q^{-1}(\delta)$ is the complete inverse image of the interval δ on the axis of values of the function $q(x, y)$.

$e(q, t)$ is the set of level t of the function $q = q(x, y)$.

$\tau(e, z)$ is the unit tangent vector to the curve e at the point $z \in e$.

$\gamma(\tau_1, \tau_2)$ is the absolute value of the acute angle between the vectors τ_1 and τ_2 .

$h_1(e)$ is the length of the set e .

$d_1(e)$ is the one-dimensional diameter of the set e .

$O(\gamma)$ is a quantity bounded by a constant depending only on γ .

$\rho(A_1, A_2)$ is the distance between the sets A_1 and A_2 in the sense of deviation, more precisely

$$\rho(A_1, A_2) = \max \left\{ \sup_{z_1 \in A_1} \inf_{z_2 \in A_2} \rho(z_1, z_2), \sup_{z_2 \in A_2} \inf_{z_1 \in A_1} \rho(z_1, z_2) \right\},$$

where $\rho(z_1, z_2)$ is the distance between the points z_1 and z_2 .