

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	23 (1977)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS
Autor:	Vitushkin, A. G.
Kapitel:	§6. The set of linear superpositions in the space of continuous functions is nowhere dense
DOI:	https://doi.org/10.5169/seals-48931

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

is satisfied, where the constant C does not depend on the functions $\{f_i^k(t)\}$. Since F is a bounded set, there exists a subsequence of suffixes n_1, n_2, \dots such that for any $i = 0, 1, \dots, n$; $k = 1, 2, \dots, m_i$; $j = 1, 2, \dots, s_i$ the numerical sequence $f_{i,n_v}^k \rightarrow C_{k,i,j}$ as $v \rightarrow \infty$. From this and the previous inequality it follows that $\{f_{i,n_v}^k(t)\} \in F$ ($v=1, 2, \dots$) is a Cauchy sequence, because it is known that the sequence $f_n(x, y) \in T(F)$ is Cauchy sequence. Consequently there exists an element $\{f_i^k(t)\} \in B$ such that $\|\{f_i^k(t) - f_{i,n_v}^k(t)\}\|_B \rightarrow 0$. Since F is a closed set, $\{f_i^k(t)\} \in F$. The operator $T: B \rightarrow C(G)$ is bounded. Therefore $T(\{f_i^k(t)\}) = f(x, y)$. Consequently $f(x, y) \in T(F)$. This proves the lemma.

The following lemma from the theory of linear operators [28] turns out to be useful.

LEMMA 4.5.3. *Let B_1 and B_2 be Banach spaces. If a linear operator $T: B_1 \rightarrow B_2$ maps bounded closed sets of B_1 onto closed sets of B_2 , then its domain of values is closed.*

Proof of Theorem 4.5.1. The set of superpositions of the form $\sum_{m=1}^N p_m(x, y) f_m(g_m(x, y))$ coincides on G with the set of superpositions of the form $\sum_{i=0}^n \sum_{k=1}^{m_i} p_i^k(x, y) f_i^k(q_i^1(x, y))$. By Lemma 4.5.2 and 4.5.3 the set of the latter superpositions is closed in the space $C(G)$. This proves the theorem.

§ 6. *The set of linear superpositions in the space of continuous functions is nowhere dense*

THEOREM 4.6.1. *For any continuous functions $p_m(x, y)$ and continuously differentiable functions $q_m(x, y)$ ($m=1, 2, \dots, N$) and any region D of the plane of the variables x, y the set of superpositions of the form*

$$\sum_{m=1}^N p_m(x, y) f_m(q_m(x, y)),$$

where $\{f_m(t)\}$ are arbitrary continuous functions, is nowhere dense in the space of all functions continuous in D with uniform convergence.

By Lemma 4.2.2 we can find a subregion $G^* \subset D$, determine a constant $\gamma^* > 0$, and renumber the functions $\{q_m(x, y)\}$, with two indices so that

the functions $\tilde{q}_i^k(x, y)$ ($i = 0, 1, 2, \dots, \tilde{n}$; $k = 1, 2, \dots, \tilde{m}_i$; $\sum_{i=0}^{\tilde{n}} \tilde{m}_i = N$) obtained after the renumbering satisfy conditions (1), (2), (3) of Lemma 4.2.2. We now fix the point $(x_0, y_0) \in G^*$ and the number v so that the line $(y - y_0) + v(x - x_0) = 0$ does not touch at any of the level curves of the functions $\tilde{q}_i^k(x, y)$ ($i = 1, 2, \dots, \tilde{n}$) that pass through (x_0, y_0) . Let $G^{**} \subset G^*$ be a disc with centre at (x_0, y_0) and radius small enough so that the $\{ \tilde{q}_i^k(x, y) \}$ and $q_{N+1}(x, y) = y + vx$ satisfy condition (3) of Lemma 4.2.2 with some constant $\gamma^{**} > 0$. We put $p_{N+1}(x, y) = 1$. By Lemma 4.4.3 we can find a set $G \subset G^{**}$, determine a constant $\lambda > 0$, and again renumber the functions $p_m(x, y)$ and $q_m(x, y)$ ($m = 1, 2, \dots, N+1$) with two indices so that the functions $p_i^k(x, y)$ and

$$q_i^k(x, y) \quad (i = 0, 1, 2, \dots, n+1; k = 1, 2, \dots, m_i; \sum_{i=0}^{n+1} m_i \leq N+1)$$

that is, some functions may be omitted in the renumbering) obtained after the renumbering satisfy conditions (1)-(3) of Lemma 4.2.2, conditions (4')-(6') of § 5, and the condition

$$7 \quad m_{n+1} = 1, \quad p_{N+1}^1 = p_{N+1}(x, y) = 1, \quad q_{N+1}^1 = q_{N+1}(x, y) = y + vx.$$

Let L be the linear space consisting of all system of functions $\{ f_i^k(t) \}$ defined and continuous on the sets $\{ q_i^1(G) \}$ and satisfying the condition

$$\sum_{i=0}^{n+1} \sum_{k=1}^{m_i} p_i^k(x, y) f_i^k(q_i^1(x, y)) \equiv 0 \quad \text{in } G.$$

LEMMA 4.6.1. *L is a finite-dimensional linear space.*

Proof. By Lemma 4.5.1, in the sets $\{ q_i^1(G) \}$ we can select a subset consisting of a finite number of points $\{ t_{i,j} \}$ such that, if $\{ f_i^k(t) \} \in L$ and $f_i^k(t_{i,j}) = 0$ for all k, i, j then $f_i^k(t) \equiv 0$ on $q_i^1(G)$ for all i, k . Thus, the set of functions $\{ f_i^k(t) \}$ is completely determined by a finite set of parameters $\{ f_i^k(t_{i,j}) \}$. Consequently the dimension of the space L is finite. This proves the lemma.

LEMMA 4.6.2. *There exists a natural number μ such that in D the polynomial $(y + vx)^\mu = Q(x, y)$ is not equal to any superposition of the form*

$$\sum_{m=1}^N p_m(x, y) f_m(q_m(x, y)), \quad \text{where } \{ f_m(t) \} \text{ are arbitrary continuous functions.}$$

Proof. We denote by Φ the space of functions of the form $f(y+vx) = f_{n+1}^1(q_{n+1}^1(x, y))$ that are representable on G by superpositions of the form $[\sum_{m=1}^N p_m(x, y)f_m(q_m(x, y))]$. Or, what comes to the same thing

(see properties (4') and (7)), of the form $[\sum_{i=0}^n \sum_{k=1}^{m_i} p_i^k(x, y)f_i^k(q_i^k(x, y))]$.

Thus, functions of Φ satisfy the relation $\sum_{i=0}^{n+1} \sum_{k=1}^{m_i} p_i^k(x, y)f_i^k(q_i^k(x, y)) \equiv 0$

in G . Consequently the linear space Φ is naturally embedded in L . Since L is finite-dimensional (Lemma 4.6.1), Φ is also finite-dimensional. Let l be the dimension of Φ . Since the polynomials $(y+vx), (y+vx)^2, \dots, (y+vx)^{l+1}$ are linearly independent, at least one of them $Q(x, y) = (y+vx)^\mu$ is not equal to any superposition of the form under discussion on G or, consequently, in D . This proves the lemma.

Proof of Theorem 4.6.1. By Lemma 4.6.2 the set of superpositions of the form given in Theorem 4.6.1 does not exhaust all continuous functions on G . Consequently, by Theorem 4.5.1, the set of these superpositions is a closed linear subspace of $C(G)$. Hence we conclude that the set of superpositions under discussion is nowhere dense in $C(G)$, nor consequently in $C(D)$. This proves the theorem.

COROLLARY 4.6.1. *For any continuous functions $p_m(x_1, x_2, \dots, x_n)$ and continuously differentiable functions $q_m(x_1, x_2, \dots, x_n)$ ($m=1, 2, \dots, N$) and any region D of the space of the variables (x_1, x_2, \dots, x_n) the set of superpositions of the form*

$$\sum_{m=1}^N p_m(x_1, x_2, \dots, x_n) f_m(q_m(x_1, x_2, \dots, x_n), x_2, x_3, \dots, x_{n-1}),$$

where $\{f_m(t, x_2, x_3, \dots, x_{n-1})\}$ are arbitrary continuous functions of $(n-1)$ variables, is nowhere dense in the space of all functions continuous in D with uniform convergence.