

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS
Autor: Vitushkin, A. G.
Kapitel: §2. (,)-entropy of the set of linear superpositions
DOI: <https://doi.org/10.5169/seals-48931>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

COROLLARY 5.1.3. For any continuous functions p_i and continuously differentiable functions $q_{1,i}, q_{2,i}, \dots, q_{k,i}$, $k < n$ ($i = 1, 2, \dots, N$) and every region G_n there exists a continuous function that is not equal in G_n to any superposition of the form (V).

§ 2. (ε, δ) -entropy of the set of linear superpositions

We denote by $S(\delta, z)$ the disc of radius δ with centre at z . Let $p(z) = p(x, y)$ and $q(z) = q(x, y)$ be functions defined in a closed region G of the x, y -plane and having the properties:

- a) $p(x, y), \frac{\partial q(x, y)}{\partial x}, \frac{\partial q(x, y)}{\partial y}$ are continuous in G and have modulus of continuity $\omega(\delta)$,
- b) the inequalities $0 < \gamma \leq |\operatorname{grad}[q(r)]| \leq \frac{1}{\gamma}$ and $|p(z)| \leq \frac{1}{\gamma}$, where γ is some constant, are satisfied everywhere in G .

LEMMA 5.2.1. Let $S(\delta, z) \subset G$ and let $\mu_q(t)$ be the function equal to $2 \sqrt{\delta^2 - (t - q(z))^2} |\operatorname{grad}[q(z)]|^{-2}$ on

$$q(z) - \delta |\operatorname{grad}[q(z)]| \leq t \leq q(z) + \delta |\operatorname{grad}[q(z)]|$$

and equal to zero elsewhere. Then

$$\int_{-\infty}^{\infty} |\mu_q(t) - h_1(e(q, t) \cap S(\delta, z))| dt \leq c_1(\gamma) \omega(\delta) \delta^2,$$

where $c_1(\gamma)$ is a constant depending only on γ .

Proof. Let $[a, b] \subset e(q, t) \cap S(\delta, z)$ be the segment of the level curve $e(q, t)$, endpoints a and b , lying on the boundary of $S(\delta, z)$; $[z, a]$ and $[z, b]$ the vectors with origin at z and endpoints at a and b , respectively;

$$\alpha_1 = \gamma([\overrightarrow{z, a}], \operatorname{grad}[q(z)]), \quad \alpha_2 = \gamma([\overrightarrow{z, b}], \operatorname{grad}[q(z)]).$$

We have

$$\begin{aligned} |t - q(z)| &= |q(a) - q(z)| = \left| \int_{s \in [z, a]} \frac{\partial q}{\partial s} ds \right| \\ &= \delta \cos \alpha_1 |\operatorname{grad}[q(z)]| (1 + O(1) \omega(\delta)) \end{aligned}$$

Hence

$$\delta \sin \alpha_1 = \sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2}$$

and similarly

$$\delta \sin \alpha_2 = \sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2}$$

By b) the size of the angle swept out by the tangent vector to the level curve $e(q, t)$ on moving along $[a, b]$ does not exceed $C_2(\gamma) \omega(\delta)$. Therefore

$$h_1([a, b]) = \delta (\sin \alpha_1 + \sin \alpha_2) (1 + 0(\gamma) \omega(\delta)) \\ = 2 \sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2} + 0(\gamma) \delta \omega(\delta).$$

If $\alpha_1 \geq C_3(\gamma) \omega(\delta)$ (C_3 is a sufficiently large constant), then $[a, b] = e(q, t) \cap S(\delta, z)$. Consequently, for

$$|t - q(z)| \leq \theta = \delta \cos [C_3 \omega(\delta)] |\operatorname{grad} [q(z)]| \times (1 + 0(1) \omega(\delta))$$

we have $h_1(e(q, t) \cap S(\delta, z)) = h_1([a, b])$. Since for every t (by b))

$$h_1(e(q, t) \cap S(\delta, z)) \leq C_4(\gamma) \delta (1 + \omega(\delta)),$$

we have

$$\int_{-\infty}^{\infty} |h_1(e(q, t) \cap S(\delta, z)) - \mu_q(t)| dt = \\ = \int_{q(z) - \theta}^{q(z) + \theta} |h_1(e(q, t) \cap S(\delta, z)) - \mu_q(t)| dt + 0(\gamma) \delta^2 \omega(\delta).$$

We now estimate

$$\int_{q(z) - \theta}^{q(z) + \theta} |h_1(e(q, t) \cap S(\delta, z)) - \mu_q(t)| dt = \\ = \int_{q(z) - \theta}^{q(z) + \theta} |h_1([a, b]) - \mu_q(t)| dt \leq \\ \leq 2 \int_{q(z) - \theta}^{q(z) + \theta} (\sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2} \\ - \sqrt{\delta^2 - (t - q(z))^2} |\operatorname{grad} [q(z)]|^{-2}) dt + 0(\gamma) \delta^2 \omega(\delta) \\ = 0(\gamma) \delta^2 \omega(\delta) \int_{-1}^1 \frac{d\tau}{\sqrt{1 - \tau^2}} + 0(\gamma) \delta^2 \omega(\delta) = 0(\gamma) \delta^2 \omega(\delta).$$

Here we have the mean value theorem. This proves the lemma.

LEMMA 5.2.2. Let $p(z), q(z)$ satisfy conditions a) and b); $S(\delta, z) \subset G$; let $f(t)$ be an arbitrary continuous function, uniformly bounded in modulus by the constant m . Then

$$\begin{aligned} & \int \int_{(u, v) \in S(\delta, z)} p(u, v) f(q(u, v)) dudv \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-1} \int_{-\infty}^{\infty} f(t) \mu_q(t) dt + \lambda(z) m \delta^2 \omega(\delta), \end{aligned}$$

where $|\lambda(z)| \leq C_5(\gamma)$.

Proof. Using a) and b) and Lemma 5.2.1 we have

$$\begin{aligned} & \int \int_{S(\delta, z)} p(u, v) f(q(u, v)) dudv \\ &= p(z) \int \int_{(u, v) \in S(\delta, z)} f(q(u, v)) dudv + O(1) m \delta^2 \omega(\delta) \\ &= p(z) \int_{-\infty}^{\infty} \left\{ f(t) \int_{s \in e(q, t) \cap S(\delta, z)} \left| \operatorname{grad} [q(s)] \right|^{-2} ds \right\} dt + O(1) m \delta^2 \omega(\delta) \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-1} \int_{-\infty}^{\infty} \left\{ f(t) \int_{s \in e(q, t) \cap S(\delta, z)} ds \right\} dt + O(\gamma) m \delta^2 \omega(\delta) \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-2} \int_{-\infty}^{\infty} f(t) h_1(e(q, t) \cap S(\delta, z)) dt + O(\gamma) m \delta^2 \omega(\delta) \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-1} \int_{-\infty}^{\infty} f(t) \mu_q(t) dt + O(\gamma) m \delta^2 \omega(\delta). \end{aligned}$$

This proves the lemma.

LEMMA 5.2.3. Suppose that a number $\alpha > 0$ and functions $p(z), q(z), f(t)$ satisfying the conditions of Lemma 5.2.2. are given. If for every integer k such that

$$\min_{z \in G} q(z) \leq t_k = k \delta \frac{\alpha}{m} \leq \max_{z \in G} q(z)$$

and any integer l such that

$$\min_{z \in G} \left| \operatorname{grad} [q(z)] \right| \leq t_l' = l \frac{\alpha}{m} \leq \max_{z \in G} \left| \operatorname{grad} [q(z)] \right|,$$

the inequality

$$\left| \int_{t_k - t_l' \delta}^{t_k + t_l' \delta} f(t) \sqrt{\delta^2 - \left(\frac{t - t_k}{t_l'} \right)^2} dt \right| \leq \alpha \delta^2$$

is satisfied, then for every disc $S(\delta, z) \subset G$

$$\left| \int_{(u,v) \in S(\delta,z)} p(u,v) f(q(u,v)) du dv \right| \leq c_6(\gamma) (\alpha \delta^2 + m \delta^2 \omega(\delta)).$$

Proof. Suppose that a disc $S(\delta, z) \subset G$ is given. By the condition of the lemma there are integers k and l such that $|q(z) - t_k| \leq \delta \alpha / m$ and $\|\operatorname{grad}[q(z)]\| - t_l' \leq \alpha / m$. From Lemma 5.2.2 we obtain

$$\begin{aligned} \left| \int_{(u,v) \in S(\delta,z)} p(u,v) f(q(u,v)) du dv \right| &\leq \frac{|p(z)|}{\|\operatorname{grad}[q(z)]\|} \left| \int_{-\infty}^{\infty} f(t) \mu_q(t) dt \right| \\ &+ c_5(\gamma) m \delta^2 \omega(\delta) \leq \frac{2}{\gamma^2} \left| \int_{t_k - \delta | \operatorname{grad}[q(z)] |}^{t_k + \delta | \operatorname{grad}[q(z)] |} f(t) \sqrt{\delta^2 - \frac{(t - q(z))^2}{\|\operatorname{grad}[q(z)]\|^2}} dt \right| \\ &- \int_{t_k - \delta | \operatorname{grad}[q(z)] |}^{t_k + \delta | \operatorname{grad}[q(z)] |} f(t) \sqrt{\delta^2 - \left(\frac{t - t_k}{t_l'} \right)^2} dt \Big| + \frac{2}{\gamma^2} \alpha \delta^2 + c_5(\gamma) m \delta^2 \omega(\delta) \leq \end{aligned}$$

(by the mean value theorem)

$$\begin{aligned} &\leq \frac{2}{\gamma^2} \alpha \delta^2 + c_5(\gamma) m \delta^2 \omega(\delta) + \frac{2}{\gamma^2} \left(\int_{-1}^1 \frac{\delta m d\tau}{\sqrt{1 - \tau^2}} \right) \delta \frac{\alpha}{m} \\ &+ \frac{2}{\gamma^2} \left(\int_{-1}^1 \frac{\delta^2 m d\tau}{\sqrt{1 - \tau^2}} \right) \frac{\alpha}{m} \leq c_6(\gamma) (\alpha \delta^2 + m \delta^2 \omega(\delta)). \end{aligned}$$

This proves the lemma.

We denote by $F_m = F_m(D; p_1, p_2, \dots, p_N; q_1, q_2, \dots, q_N)$ the set of superpositions of the form

$$f(x, y) = \sum_{i=1}^N p_i(x, y) f_i(q_i(x, y)), \text{ where } \{p_i(x, y)\}$$

and $\{q_i(x, y)\}$ are fixed functions, defined in the closed region D of the x, y plane and satisfying conditions a) and b) with a constant γ not depending on i and $\{f_i(t)\}$ are arbitrary continuous functions, defined on $\{[a_i, b_i]\} = \{[\min_{z \in D} q_i(z); \max_{z \in D} q_i(z)]\}$ and uniformly bounded in modulus by the constant m .

THEOREM 5.2.1. *There exist constants A and B such that if $\varepsilon > Am\omega(\delta)$ then for the (ε, δ) -entropy of the set of functions F_m , $H_{\varepsilon, \delta}(F_m) \leq \frac{B}{\delta} \left(\frac{m}{\varepsilon} \right)^2$, where A and B depend only on γ, N and D .*

Proof. We put

$$R(f(z), \delta) = \max_{S(\delta, z) \subset D} \left| \frac{1}{\pi \delta^2} \iint_{(u, v) \in S(\delta, z)} f(u, v) dudv \right|.$$

We denote by $\mathcal{H}_{\varepsilon, \delta}(F_m)$ the ε -entropy of the space F_m , taking as the distance between the functions $f_1(z), f_2(z) \in F_m$ the number $R(f_1(z) - f_2(z), \delta)$. The inequality $H_{2\varepsilon, \delta}(F_m) \leq \mathcal{H}_{\varepsilon, \delta}(F_m)$ holds owing to the fact that if two functions $f_1(z)$ and $f_2(z)$ are (ε, δ) -distinguishable, then they are ε -distinguishable also in the sense of the metric $R(f_1(z) - f_2(z), \delta)$. We now estimate the value of $\mathcal{H}_{\varepsilon, \delta}(F_m)$. Let k and l be integers such that

$$\min_{z \in D} q_i(z) \leq t_k = k\delta \frac{\alpha}{m} \leq \max_{z \in D} q_i(z)$$

and

$$\min_{z \in D} |\operatorname{grad} [q_i(z)]| \leq t_l' = l \frac{\alpha}{m} \leq \max_{z \in D} |\operatorname{grad} [q_i(z)]|.$$

To compute the function

$$f_\delta(z) = \frac{1}{\pi \delta^2} \iint_{(u, v) \in S(\delta, z)} f(u, v) dudv,$$

where $f(x, y) \in F_m$, $S(\delta, z) \subset D$ to within ε , it is sufficient by Lemma 5.2.3 to give the values of

$$v_i(t_k, t_l') = \frac{1}{\pi \delta^2} \int_{t_k - t_l' \delta}^{t_k + t_l' \delta} f_i(t) \sqrt{\delta^2 - \left(\frac{t - t_k}{t_l'} \right)^2} dt$$

to within $\alpha = \pi \varepsilon / (2 NC_B(\gamma))$ and to assume that δ is small enough so that

$$\varepsilon > \frac{2NC_B(\gamma)m\omega(\delta)}{\pi} = A(\gamma, N)m\omega(\delta).$$

Since $|v_i(t_k, t_l')| \leq C_1 m$, to write the numbers $v_i(t_k, t_l')$ (i, k, l fixed) $\log_2(C_1 m/\alpha)$ binary digits are sufficient. Since

$$|v_i(t_{k+1}, t_l') - v_i(t_k, t_l')| \leq c_8 \frac{1}{\delta^2} \left(\int_{-1}^1 \frac{\delta m d\tau}{\sqrt{1-\tau^2}} \right) \delta \frac{\alpha}{m} = c_9(\gamma) \alpha$$

(here we again use the mean value theorem), to store the numbers $v_i(t_{k+1}, t_l') - v_i(t_k, t_l')$ to within α , $\log_2 C_9$ binary digits are sufficient. Therefore to write the numbers $v_i(t_k, t_l')$ (i, l fixed; k any admissible number)

$C_{10}(\gamma) \left[\log_2 \frac{m}{\alpha} + (b_i - a_i) \frac{m}{\delta \alpha} \right] = \mathcal{H}_{i,l}$ binary digits are sufficient. Consequently the total number of digits sufficient to store all the numbers $v_i(t_k, t_l')$ to within α , that is, to store the functions $f_\delta(z)$ to within ε , is

$$\mathcal{H} = \sum_{i,l} \mathcal{H}_{i,l} \leq N c_{10}(\gamma) \left[\log_2 \frac{m}{\alpha} + (b_i - a_i) \frac{m}{\delta \alpha} \right] \frac{1}{\gamma} \frac{m}{\alpha} \leq \frac{B(\gamma, N, D)}{\delta} \left(\frac{m}{\varepsilon} \right)^2.$$

This proves the theorem.

§ 3. Functional “dimension” of the space of linear superpositions

Suppose that continuous functions $p_i(x, y)$ and continuously differentiable functions $q_i(x, y)$ ($i = 1, 2, \dots, N$) are fixed. Let G be a closed region of the x, y plane. We denote by $F = F(G, \{p_i\}, \{q_i\})$ the set of superpositions of the form $f(x, y) = \sum_{i=1}^N p_i(x, y) f_i(q_i(x, y))$, where $(x, y) \in G$ and $\{f_i(t)\}$ are arbitrary continuous functions of one variable. We are interested in the functional dimension of the set F .

THEOREM 5.3.1. *In every region D of the x, y plane there exists a closed subregion $G \subset D$ such that*

$$r(F(G, \{p_i\}, \{q_i\})) \leq 1.$$

Proof. By Theorem 4.5.1, in D there exists a closed subregion $G^* \subset D$ such that the set of superpositions $F(G^*, \{p_i\}, \{q_i\})$ is closed (in the uniform metric) in $C(G^*)$, and the functions $\{q_i(x, y)\}$ satisfy the condition: for any i , either $\text{grad}[q_i(x, y)] \neq 0$ on G^* or $q_i(x, y) \equiv \text{const}$ on G^* . We show that $r(F(G^*, \{p_i\}, \{q_i\})) \leq 1$. By Banach's open mapping theorem, there exists a constant K such that for any superposition $\sum_{i=1}^N p_i(x, y) f_i(q_i(x, y)) = f(x, y) \in F(G^*, \{p_i\}, \{q_i\})$ there are con-