

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS
Autor: Vitushkin, A. G.
Kapitel: §3. Functional "dimension" of the space of linear superpositions
DOI: <https://doi.org/10.5169/seals-48931>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$|v_i(t_{k+1}, t'_l) - v_i(t_k, t'_l)| \leq c_8 \frac{1}{\delta^2} \left(\int_{-1}^1 \frac{\delta m d\tau}{\sqrt{1-\tau^2}} \right) \delta \frac{\alpha}{m} = c_9(\gamma) \alpha$$

(here we again use the mean value theorem), to store the numbers $v_i(t_{k+1}, t'_l) - v_i(t_k, t'_l)$ to within α , $\log_2 C_9$ binary digits are sufficient. Therefore to write the numbers $v_i(t_k, t'_l)$ (i, l fixed; k any admissible number)

$C_{10}(\gamma) \left[\log_2 \frac{m}{\alpha} + (b_i - a_i) \frac{m}{\delta \alpha} \right] = \mathcal{H}_{i,l}$ binary digits are sufficient. Consequently the total number of digits sufficient to store all the numbers $v_i(t_k, t'_l)$ to within α , that is, to store the functions $f_\delta(z)$ to within ε , is

$$\mathcal{H} = \sum_{i,l} \mathcal{H}_{i,l} \leq N c_{10}(\gamma) \left[\log_2 \frac{m}{\alpha} + (b_i - a_i) \frac{m}{\delta \alpha} \right] \frac{1}{\gamma} \frac{m}{\alpha} \leq \frac{B(\gamma, N, D)}{\delta} \left(\frac{m}{\varepsilon} \right)^2.$$

This proves the theorem.

§ 3. Functional "dimension" of the space of linear superpositions

Suppose that continuous functions $p_i(x, y)$ and continuously differentiable functions $q_i(x, y)$ ($i=1, 2, \dots, N$) are fixed. Let G be a closed region of the x, y plane. We denote by $F = F(G, \{p_i\}, \{q_i\})$ the set of superpositions of the form $f(x, y) = \sum_{i=1}^N p_i(x, y) f_i(q_i(x, y))$, where $(x, y) \in G$ and $\{f_i(t)\}$ are arbitrary continuous functions of one variable. We are interested in the functional dimension of the set F .

THEOREM 5.3.1. *In every region D of the x, y plane there exists a closed subregion $G \subset D$ such that*

$$r(F(G, \{p_i\}, \{q_i\})) \leq 1.$$

Proof. By Theorem 4.5.1, in D there exists a closed subregion $G^* \subset D$ such that the set of superpositions $F(G^*, \{p_i\}, \{q_i\})$ is closed (in the uniform metric) in $C(G^*)$, and the functions $\{q_i(x, y)\}$ satisfy the condition: for any i , either $\text{grad}[q_i(x, y)] \neq 0$ on G^* or $q_i(x, y) \equiv \text{const}$ on G^* . We show that $r(F(G^*, \{p_i\}, \{q_i\})) \leq 1$. By Banach's open mapping theorem, there exists a constant K such that for any superposition

$\sum_{i=1}^N p_i(x, y) f_i(q_i(x, y)) = f(x, y) \in F(G^*, \{p_i\}, \{q_i\})$ there are con-

tinuous functions $\{f_i^*(t)\}$, defined on the sets $\{q_i(G^*)\}$ and satisfying the conditions

$$8) \quad f(x, y) = \sum_{i=1}^N p_i(x, y) f_i^*(q_i(x, y)) \text{ for all } (x, y) \in G^* ;$$

$$9) \quad \max_i \max_{t \in q_i(G^*)} |f_i^*(t)| \geq K \max_{(x, y) \in G^*} |f(x, y)| .$$

Denote by $F_{\lambda\varepsilon} = F_{\lambda\varepsilon}(G^*, \{p_i\}, \{q_i\})$ the set of superpositions $f(x, y) \in F(G^*, \{p_i\}, \{q_i\})$ such that $\max_{(x, y) \in G^*} |f(x, y)| \leq \lambda\varepsilon$. By Theorem 5.2.1

and (8), (9), there exist constants A and B such that if $\omega(\delta) \leq (\lambda AK)^{-1}$ then $H_{\varepsilon, \delta}(F_{\lambda\varepsilon}) \leq B(\lambda K)^2/\delta$. Hence the functional dimension

$$r(F_i(G^*, \{p_i\}, \{q_i\})) \leq \lim_{\lambda \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \frac{\log_2 \log_2 \frac{B(\lambda K)^2}{\delta}}{\log_2 \delta} = 1$$

This proves the theorem.

From Theorem 5.3.1 and the properties of functional dimension (§ 1) we have the following result, which is a stronger form of Theorem 4.6.1.

COROLLARY 5.3.1. *For any continuous functions $\{p_i(x, y)\}$ and continuously differentiable functions $\{q_i(x, y)\}$ and every region D the set of linear superpositions $F(D, \{p_i\}, \{q_i\})$ is nowhere dense in any space of functions that has in every region $G \subset D$ functional "dimension" greater than 1.*

Remark 5.3.1. All the results about linear superpositions of the form $\sum_{i=1}^N p_i(x, y) f_i(q_i(x, y))$ remain valid if we assume that $\{f_i(t)\}$ are arbitrary bounded measurable functions.

§ 4. Variation of superpositions of smooth functions

Let G_n be a closed region of the space of the variables x_1, x_2, \dots, x_n ($n \geq 2$). A function $F(x) = F(x_1, x_2, \dots, x_n)$ is called a superposition of order s generated by the functions of k ($k > 1$) variables

$$f_{\beta_1, \beta_2, \dots, \beta_\alpha}(t_1, t_2, \dots, t_k) \quad (\alpha = 0, 1, 2, \dots, s; \beta_i = 1, 2, \dots, k)$$

if it is defined in G by relations