Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 23 (1977)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON REPRESENTATION OF FUNCTIONS BY MEANS OF SUPERPOSITIONS AND RELATED TOPICS

Autor: Vitushkin, A. G.

Kapitel: §4. Variation of superpositions of smooth functions

DOI: https://doi.org/10.5169/seals-48931

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 20.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

tinuous functions $\{f_i^*(t)\}$, defined on the sets $\{q_i(G^*)\}$ and satisfying the conditions

8) $$f(x, y) = \sum_{i=1}^{N} p_i(x, y) f_i^*(q_i(x, y)) \text{ for all } (x, y) \in G^*;$$

9) $$\max_i \max_{t \in q_i(G^*)} |f_i^*(t)| \geqslant K \max_{(x,y) \in G^*} |f(x, y)|.$$

Denote by $F_{\lambda\varepsilon} = F_{\lambda\varepsilon}(G^*, \{p_i\}, \{q_i\})$ the set of superpositions $f(x,y) \in F(G^*, \{p_i\}, \{q_i\})$ such that $\max_{(x,y) \in G^*} |f(x, y)| \leqslant \lambda\varepsilon$. By Theorem 5.2.1 and (8), (9), there exist constants A and B such that if $\omega(\delta) \leqslant (\lambda AK)^{-1}$ then $H_{\varepsilon,\delta}(F_{\lambda\varepsilon}) \leqslant B(\lambda K)^2/\delta$. Hence the functional dimension

$$r(F_i(G^*, \{p_i\}, \{q_i\})) \leqslant \lim_{\lambda \to \infty} \lim_{\delta \to 0} \lim_{\varepsilon \to 0} \frac{\log_2 \log_2 \frac{B(\lambda K)^2}{\delta}}{\log_2 \delta} = 1$$

This proves the theorem.

From Theorem 5.3.1 and the properties of functional dimension (§ 1) we have the following result, which is a stronger form of Theorem 4.6.1.

COROLLARY 5.3.1. *For any continuous functions* $\{p_i(x, y)\}$ *and continuously differentiable functions* $\{q_i(x, y)\}$ *and every region* D *the set of linear superpositions* $F(D, \{p_i\}, \{q_i\})$ *is nowhere dense in any space of functions that has in every region* $G \subset D$ *functional "dimension" greater than* 1.

Remark 5.3.1. All the results about linear superpositions of the form $\sum_{i=1}^{N} p_i(x, y) f_i(q_i(x, y))$ remain valid if we assume that $\{f_i(t)\}$ are arbitrary bounded measurable functions.

§ 4. *Variation of superpositions of smooth functions*

Let G_n be a closed region of the space of the variables $x_1, x_2, \ldots, x_n$ $(n \geqslant 2)$. A function $F(x) = F(x_1, x_2, \ldots, x_n)$ is called a superposition of order s generated by the functions of k $(k > 1)$ variables

$$f_{\beta_1, \beta_2 \ldots, \beta_\alpha}(t_1, t_2, \ldots, t_k) \; (\alpha = 0, 1, 2, \ldots, s; \; \beta_i = 1, 2, \ldots, k)$$

if it is defined in G by relations

$$\left\{\begin{array}{l} F = f(q_1, q_2, \ldots, q_k), \\ \cdots\cdots\cdots\cdots\cdots\cdots\cdots\cdots \\ q_{\beta_1,\beta_2,\ldots,\beta_\alpha} = f_{\beta_1,\ldots,\beta_\alpha}(q_{\beta_1,\ldots,\beta_\alpha,1} q_{\beta_1,\ldots,\beta_\alpha,2}, \ldots, q_{\beta_1,\ldots,\beta_\alpha,k}), \\ \cdots\cdots\cdots\cdots\cdots\cdots\cdots\cdots \\ q_{\beta_1,\beta_2,\ldots,\beta_{s+1}} = x_{\gamma(\beta_1,\beta_2,\ldots,\beta_{s+1})}, \end{array}\right. \qquad \text{(VI)}$$

where $\gamma(\beta_1, \beta_2, \ldots, \beta_{s+1})$ is a function of the indices $\beta_1, \beta_2, \ldots, \beta_{s+1}$ and takes one of the values $1, 2, \ldots, n$. As before, we assume that the functions $\{\varphi_{\beta_1,\beta_2,\ldots,\beta_\alpha}(t_1, t_2, \ldots, t_k)\}$ are defined for all values of the arguments.

A superposition of any order, generated by functions of one variable, is again a function of one variable. Therefore in this case ($k=1$) we consider superpositions of functions of one variable and the operation of addition, that is, superpositions definable in the following way.

A function $F(x) = F(x_1, x_2, \ldots, x_n)$ $(n>1)$ is called a superposition of order s of the functions $f_{\beta_1,\ldots,\beta_\alpha}(t)$ $(\alpha=0, 1, 2, \ldots, s;\ \beta_i=1, 2)$ if the following relations are satisfied:

$$\left.\begin{array}{l} F = f(q_1+q_2), \\ \cdots\cdots\cdots\cdots\cdots\cdots\cdots\cdots \\ q_{\beta_1,\beta_2,\ldots,\beta_\alpha} = f_{\beta_1,\beta_2,\ldots,\beta_\alpha}(q_{\beta_1,\beta_2,\ldots,\beta_\alpha,1} + q_{\beta_1,\beta_2,\ldots,\beta_\alpha,2}) \\ \cdots\cdots\cdots\cdots\cdots\cdots\cdots\cdots \\ q_{\beta_1,\beta_2,\ldots,\beta_{s+1}} = x_{\gamma(\beta_1,\beta_2,\ldots,\beta_{s+1})}, \end{array}\right\} \qquad \text{(VII)}$$

where $\gamma(\beta_1, \beta_2, \ldots, \beta_{s+1})$ takes one of the values $1, 2, \ldots, n$.

Note that we can represent as superpositions of the form (VII), for example, all rational functions of $x_1, x_2, \ldots, x_n$ since we can write any arithmetic operation by such superpositions, for example, $u \cdot v = e^{\ln u + \ln v} = f(f_1(u) + f_2(v))$.

Let $F(x_1, x_2, \ldots, x_n)$ be a superposition of order s of the continuously differentiable functions $\{f_{\beta_1,\beta_2,\ldots,\beta_\alpha}(t_1, t_2, \ldots, t_k)\}$ and $\tilde{F}(x_1, x_2, \ldots, x_n)$ the superposition of the same form of the continuously differentiable functions $\{\tilde{f}_{\beta_1,\beta_2,\ldots,\beta_\alpha}(t_1, t_2, \ldots, t_k)\}$. We put

$$\varphi_{\beta_1,\beta_2,\ldots,\beta_\alpha} = \tilde{f}_{\beta_1,\ldots,\beta_\alpha} - f_{\beta_1,\ldots,\beta_\alpha} \quad (\alpha=0, 1, 2, \ldots, s;\ \beta_i=1, 2, \ldots, k)$$

$$\mu = \max_{\alpha, \beta_1, \ldots, \beta_\alpha} \sum_{i=1}^{k} \sup_t \left| \frac{\partial f_{\beta_1,\ldots,\beta_\alpha}(t_1, \ldots, t_k)}{\partial t_i} \right|,$$

$$\varepsilon = \max_{\alpha, \beta_1, \ldots, \beta_\alpha} \sup_t \left| \varphi_{\beta_1,\ldots,\beta_\alpha}(t_1, t_2, \ldots, t_k) \right|$$

LEMMA 5.4.1. *The inequality*

$$\sup_{x \in G} \left| \tilde{F}(x_1, x_2, \ldots, x_n) - F(x_1, x_2, \ldots, x_n) \right| \leqslant A(\mu, s)\, \varepsilon .$$

holds, where the constant $A(\mu, s)$ *depends only on* μ *and* s.

Proof. We proceed by induction on s. For definiteness suppose that $k < 1$. Having verified the statement of the lemma for $s = 1$ and having made an appropriate inductive assumption for superpositions of order $s - 1$, we have

$$\sup_{x \in G} \left| \tilde{F}(x_1, x_2, \ldots, x_n) - F(x_1, x_2, \ldots, x_n) \right|$$

$$\leqslant \left| f(\tilde{q}_1, \ldots, \tilde{q}_k) - f(q_1, \ldots, q_k) \right| + \left| \varphi(\tilde{q}_1, \tilde{q}_2, \ldots, \tilde{q}_k) \right|$$

$$\leqslant \mu \max_{\beta_1} \sup_{x \in G} \left| \tilde{q}_{\beta_1} - q_{\beta_1} \right| + \varepsilon \leqslant \mu \cdot A(\mu, s-1)\, \varepsilon + \varepsilon = A(\mu, s)\, \varepsilon .$$

(the last by the indictive assumption). This proves the lemma.

Further, let $\omega(\delta)$ be the common modulus of continuity of all the functions $\left\{ \dfrac{\partial f_{\beta_1, \ldots, \beta_\alpha}(t_1, \ldots, t_k)}{\partial t_i} \right\}$ and, in addition, put

$$\varepsilon' = \max_{\alpha, \beta_1, \ldots, \beta_\alpha} \sum_{i=1}^{k} \sup_{t} \left| \frac{\partial \varphi_{\beta_1, \ldots, \beta_\alpha}(t_1, \ldots, t_k)}{\partial t_i} \right|$$

LEMMA 5.4.2. *We have (for case* $k > 1$*)*

$$\tilde{F}(x_1, \ldots, x_n) - F(x_1, \ldots, x_n) = \sum_{\alpha, \beta_1, \ldots, \beta_\alpha} p_{\beta_1, \ldots, \beta_\alpha}(x_1, x_2, \ldots, x_n)$$

$$\times\, \varphi_{\beta_1, \ldots \beta_\alpha}\left(q_{\beta_1, \ldots, \beta_\alpha, 1}(x_1, \ldots, x_n), \ldots, q_{\beta_1, \ldots, \beta_\alpha, k}(x_1, \ldots, x_n)\right)$$

$$+ R(x_1, x_2, \ldots, x_n) ,$$

where

$$\left| R(x_1, x_2, \ldots, x_n) \right| \leqslant B(\mu, s, k) \left[\varepsilon' + \omega\left(A(\mu, s)\, \varepsilon\right) \right] \varepsilon ,$$

$$p_{\beta_1, \ldots, \beta_\alpha}(x_1, x_2, \ldots, x_n) = \prod_{i=0}^{\alpha-1} \frac{\partial f_{\beta_1, \ldots, \beta_i}}{\partial q_{\beta_1, \ldots, \beta_{i+1}}}$$

(for $\alpha = 0$ $p(x_1, x_2, \ldots, x_n) \equiv 1$),

$B(\mu, s, k)$ *is a constant depending only on* μ, s, k. *For* $k = 1$ *the corresponding equation is slightly different (see Chapter I, (III)) :*

$$\tilde{F}(x_1, \ldots, x_n) - F(x_1, \ldots, x_n)$$
$$= \sum_{\alpha, \beta_1, \ldots, \beta_\alpha} p_{\beta_1, \ldots, \beta_\alpha}(x_1, x_2, \ldots, x_n)\, \varphi_{\beta_1, \ldots, \beta_\alpha}(q_{\beta_1, \ldots, \beta_\alpha, 1}(x_1, \ldots, x_n)$$
$$+ q_{\beta_1, \ldots, \beta_\alpha, 2}(x_1, \ldots, x_n)) + R(x_1, \ldots, x_n)\,.$$

Proof. As in the preceding lemma we proceed by induction on s. Again for definiteness we limit ourselves to the case $k > 1$. For $s = 1$ the assertion of the lemma is easily verified. We assume that it is true for superpositions of order $s - 1$. By Lemma 5.4.1, for superpositions of order s we have

$$\tilde{F}(x_1, \ldots, x_n) - F(x_1, \ldots, x_n) = f(\tilde{q}_1, \tilde{q}_2, \ldots, \tilde{q}_k) - f(q_1, q_2, \ldots, q_k)$$
$$+ \varphi(\tilde{q}_1, \tilde{q}_2, \ldots, \tilde{q}_k) = \varphi(q_1, q_2, \ldots, q_k) + \sum_{\beta_1 = 1}^{k} \frac{\partial f}{\partial q_{\beta_1}} (\tilde{q}_{\beta_1} - q_{\beta_1})$$
$$+ A(\mu, s)\, \varepsilon' \cdot \varepsilon + k \cdot A(\mu, s)\, \omega\big(A(\mu, s)\, \varepsilon\big)\, \varepsilon\,.$$

Since $\tilde{q}_{\beta_1}$ and q_{β_1} $(\beta_1 = 1, 2, \ldots, k)$ are superpositions of order $s - 1$, by the inductive hypothesis we have

$$\tilde{q}_{\beta_1} - q_{\beta_1} = \sum_{\substack{\alpha > 0 \\ \beta_2, \beta_3, \ldots, \beta_\alpha}} \hat{p}_{\beta_1, \ldots, \beta_\alpha}(x_1, x_2, \ldots, x_n)$$
$$\times\, \varphi_{\beta_1, \ldots \beta_\alpha,}\big(q_{\beta_1, \ldots, \beta_\alpha, 1}(x_1, x_2, \ldots, x_n), \ldots, q_{\beta_1, \ldots, \beta_\alpha, k}(x_1, x_2, \ldots, x_n)\big)$$
$$+ \acute{R}(x_1, x_2, \ldots, x_n)\,,$$

where

$$\big|\hat{R}(x_1, x_2, \ldots, x_n)\big| \leqslant B(\mu, s-1, k)\big[\varepsilon' + \omega\big(A(\mu, s-1)\, \varepsilon\big)\big]\, \varepsilon\,,$$
$$\hat{p}_{\beta_1, \ldots, \beta_\alpha}(x_1, \ldots, x_n) = \prod_{i=1}^{\alpha - 1} \frac{\partial f_{\beta_1, \beta_2, \ldots, \beta_i}}{\partial q_{\beta_1, \ldots, \beta_{i+1}}}$$

(for $\alpha = 1$, $\hat{p}_{\beta_1}(x_1, \ldots, x_n) \equiv 1$).

When we now substative the expressions for the differences $\tilde{q}_{\beta_1} - q_{\beta_1}$ in the formula for $\tilde{F} - F$ above, we obtain the required representation of the difference of two superpositions $\tilde{F} - F$. This proves the lemma.