

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: QUADRATIC FORMS IN AN ADELIC SETTING
Autor: Verner, Lawrence
Kapitel: 3. Formulation of Siegel's Theorem
DOI: <https://doi.org/10.5169/seals-48916>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

of a compact set K . The right-hand side of the above formula reduces to the volume of K , while the left-hand side gives the mean value of

$$\text{card} (L - \{0\} \cap K),$$

as L varies over all \mathbf{Z} -lattices in \mathbf{R}^n with volume 1.

We turn now to the adelic mean value formula. Let G be a linear algebraic group defined over \mathbf{Q} , and let X be an algebraic homogeneous space for G , defined over \mathbf{Q} . For $\xi \in X$, let $G_\xi = \{g \in G: g\xi = \xi\}$. We assume that

- a) X has at least one rational point
- b) for any $\xi \in X_{\mathbf{C}}$, both $G_{\mathbf{C}}$ and $(G_\xi)_{\mathbf{C}}$ have finite fundamental groups
- c) for any extension field K of \mathbf{Q} , G_K acts transitively on X_K .

We then have the following result.

THEOREM (Ono [2]). There are canonical measures on the adèle spaces G_A and X_A such that, given any continuous function Φ on X_A with compact support,

$$(A) \quad \frac{\int_{G_A/G_{\mathbf{Q}}} \sum_{x \in X_{\mathbf{Q}}} \Phi(gx) dg}{\tau(G_\xi)} = \int_{X_A} \Phi(x) dx,$$

where ξ is any element of $X_{\mathbf{Q}}$, and $\tau(G_\xi) =$ the invariant measure of $(G_\xi)_A / (G_\xi)_{\mathbf{Q}}$. The analogy to the previous mean value theorem is clear in the cases when $\tau(G) = \tau(G_\xi)$.

3. FORMULATION OF SIEGEL'S THEOREM

Let S and T be square matrices with integral entries of size m and n , respectively. We assume that both are positive definite. For any matrix x , denote $S[x] = {}^t x S x$ (when defined). Let $A(S, T) =$ the number of integral $m \times n$ matrices x such that $S[x] = T$. For each positive integer q , let $A_q(S, T) =$ the number of integral $m \times n$ matrices x , mod q , such that $S[x] \equiv T \pmod{q}$.

A positive definite integral matrix S' is said to be in the same class as S if $S' = S[U]$, for some $U \in SL(m, \mathbf{Z})$. S' is in the same genus as S if for each q , there exists $U \in SL(m, \mathbf{Z})$ such that $S' \equiv S[U] \pmod{q}$. Let S_1, \dots, S_h be the representatives of the classes in genus (S) . Let $E(S_i) =$ the finite group consisting of all $U \in SL(m, \mathbf{Z})$ such that $S_i[U] = S_i$, and put

$e_i = 1 / \# E(S_i)$, where $\#$ denotes cardinality. We now define the “number of representations of T by the genus of S ” as

$$A(\text{genus}(S), T) = \frac{e_1 A(S_1, T) + \dots + e_h A(S_h, T)}{e_1 + \dots + e_h}.$$

Now S is a real symmetric matrix, and so we may view it as a point in \mathbf{R}^{n_1} , where $n_1 = n(n+1)/2$. Similarly, T is a point in \mathbf{R}^{m_1} . Let dt be the usual measure in \mathbf{R}^{m_1} , and let dx be the usual measure in the real vector space of $m \times n$ matrices. Given $\varepsilon > 0$, let B_ε denote the ε -neighborhood of T in \mathbf{R}^{m_1} , and let C_ε denote the set of $x \in M_{m \times n}(\mathbf{R})$ satisfying $S[x] \in B_\varepsilon$. Then B_ε and C_ε are open sets with compact closure, and the following limit is known to exist:

$$A_\infty(S, T) = \lim_{\varepsilon \rightarrow 0} \int_{C_\varepsilon} dx / \int_{B_\varepsilon} dt.$$

THEOREM (Siegel [4]). For $m - n \geq 3$,

$$(S) \quad A(\text{genus}(S), T) = A_\infty(S, T) \lim_q \frac{A_q(S, T)}{q^{mn - (n+1)n/2}}.$$

4. DERIVATION OF SIEGEL'S THEOREM

Let $G = \{g \in SL(m) : S[g] = S\}$, and let $X = \{x \in M_{m \times n} : S[x] = T\}$. If $m \geq 4$, both $G_{\mathbf{C}}$ and $G_{\mathbf{R}}$ have fundamental groups of order 2. Condition (c) of § 2 is the classical Witt theorem for (G, X) . We assume that $X_{\mathbf{Q}}$ is nonempty.

We will show that (A) implies (S). This reduces Siegel's theorem to the computation of the Tamagawa number $\tau(G)$.

Let $\Phi_\infty =$ the constant function 1 on $X_{\mathbf{R}}$, and let $\Phi_p =$ the characteristic function of $X_{\mathbf{Z}_p}$ in $X_{\mathbf{Q}_p}$. Then $\Phi = \Phi_\infty \cdot \prod \Phi_p$ is the characteristic function of $X_{S_\infty} = X_{\mathbf{R}} \cdot \prod X_{\mathbf{Z}_p}$ in $X_{\mathbf{A}}$. Because of the positive definiteness of S , Φ has compact support.

Consider the right-hand side of formula (S). Siegel has shown that there exists an algebraic gauge form dx on X such that $A_\infty(S, T) = \int_{X_{\mathbf{R}}} dx_\infty$, and

$$\lim_q \frac{A_q(S, T)}{q^{mn - (n+1)n/2}} = \prod_p \int_{X_{\mathbf{Z}_p}} dx_p,$$

where dx and dx_p are the positive measures induced on $X_{\mathbf{R}}$ and $X_{\mathbf{Q}_p}$ by dx .