Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 23 (1977)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE EVALUATION OF GAUSSIAN SUMS FOR NON-PRIMITIVE DIRICHLET CHARACTERS

Autor: Joris, Henri

Kapitel: 1. Introduction

DOI: https://doi.org/10.5169/seals-48917

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 24.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

ON THE EVALUATION OF GAUSSIAN SUMS FOR NON-PRIMITIVE DIRICHLET CHARACTERS

by Henri JORIS

1. INTRODUCTION

Let χ be a Dirichlet character mod m. We denote the conductor of χ by f and the corresponding primitive character by ψ. For a natural number α, we have the Gaussian sum

$$\mathcal{G}(\alpha, \chi) = \sum_{k \bmod m} \chi(k) \exp\left(2\pi i \frac{k\alpha}{m}\right).$$

We will write $\tau(\chi)$ for $\mathcal{G}(1, \chi)$. It is well known that

$$\mathcal{G}(\alpha, \chi) = \bar{\chi}(\alpha)\tau(\chi), \tag{1}$$

if χ is primitive, i.e. if $\chi = \psi$. For non primitive characters, (1) does not hold; according to H. Hasse [1], one has the following result:

THEOREM A. Let χ, m, ψ, f be as above. For $\alpha \in \mathbf{N}$ we put $\alpha_0 = \alpha/(\alpha, m)$, $m_0 = m/(\alpha, m)$. Then we have:

$$\left.\begin{aligned} \mathcal{G}(\alpha, \chi) &= 0 \qquad \text{if } f \nmid m_0, \\ \mathcal{G}(\alpha, \chi) &= \frac{\varphi(m)}{\varphi(m_0)} \mu\left(\frac{m_0}{f}\right) \psi\left(\frac{m_0}{f}\right) \bar{\psi}(\alpha_0)\tau(\psi) \\ &\qquad \text{if } f \mid m_0. \end{aligned}\right\} \tag{2}$$

Here, as throughout this note, φ and μ stand for the Euler totient and the Moebius function.

In [1], theorem A is proved in an elementary way, using several steps of reduction. See also [2].

In the present note we give another evaluation of $\mathcal{G}(\alpha, \chi)$, using the functional equation for Dirichlet L-series.

THEOREM B. Let m, χ, f, ψ be as above. We put

$$q = \prod_{\substack{p \mid m \\ p \nmid f}} p\,, \quad R = \frac{m}{fq}$$

where p denotes rational primes. Let the multiplicative function g be defined by

$$g(n) = \mu((n,q))\,\varphi((n,q))\,\bar{\psi}(n)\,.$$

Then we have:

$$\left.\begin{aligned} \mathcal{G}(\alpha,\chi) &= 0 \text{ if } R \nmid \alpha \\ \mathcal{G}(Rn,\chi) &= \mu(q)\,\psi(q)\,\tau(\psi)\,R\,g(n), \quad n = 1, 2, \ldots \end{aligned}\right\} \tag{3}$$

2. PROOF OF THEOREM B

For a Dirichlet character χ mod m let the function $L(s,\chi)$ be given by

$$L(s,\chi) = \sum_{1}^{\infty} \chi(n)\,n^{-s}\,, \quad Re\, s > 1\,.$$

The series defines an analytic function for Re $s > 1$, which can be extended to a meromorphic function on the whole complex plane, with at most one simple pole at $s = 1$. If χ is primitive, then $L(s,\chi)$ satisfies the equation

$$L(1-s,\chi) = m^{s-1}(2\pi)^{-s}\,\Gamma(s)\left(e^{-\frac{\pi i s}{2}} + \chi(-1)\,e^{\frac{\pi i s}{2}}\right)\tau(\chi)\,L(s,\bar{\chi})\,. \tag{4}$$

Because of (1), this can also be written, for Re $s > 1$, as

$$L(1-s,\chi) = m^{s-1}(2\pi)^{-s}\,\Gamma(s)\left(e^{-\frac{\pi i s}{2}} + \chi(-1)\,e^{\frac{\pi i s}{2}}\right)\sum_{1}^{\infty}\mathcal{G}(n,\chi)\,n^{-s}\,. \tag{5}$$

Whereas (4) holds only for primitive characters, (5) turns out to be valid in the general case. In fact, a much more general formula is proved in [3], th. 6.1; if we put there $x = \alpha = 0$, $\alpha_n = \chi(n)$ and observe that $\mathcal{G}(-n,\chi) = \chi(-1)\,\mathcal{G}(n,\chi)$, we get (5) immediately. But also most of the classical (= non-adelic) proofs of (4) will give (5) after very small changes. The only use of the primitivity of χ in these proofs is that they replace $\mathcal{G}(n,\chi)$ by $\bar{\chi}(n)\,\tau(\chi)$ at some stage (See for instance [7]).

Now let χ, m, ψ, f be as in the theorem. We have, by the Euler-product,

$$\begin{aligned} L(1-s,\chi) &= L(1-s,\psi)\prod_{p \mid q}\left(1 - \frac{\psi(p)}{p^{1-s}}\right) \\ &= L(1-s,\psi)\,\mu(q)\,q^{s-1}\,\psi(q)\prod_{p \mid q}(1 - p\bar{\psi}(p)\,p^{-s}) \end{aligned} \tag{6}$$