

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	23 (1977)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON THE EVALUATION OF GAUSSIAN SUMS FOR NON-PRIMITIVE DIRICHLET CHARACTERS
Autor:	Joris, Henri
Kapitel:	2. Proof of theorem B
DOI:	https://doi.org/10.5169/seals-48917

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THEOREM B. Let m, χ, f, ψ be as above. We put

$$q = \prod_{\substack{p|m \\ p \nmid f}} p, \quad R = \frac{m}{fq}$$

where p denotes rational primes. Let the multiplicative function g be defined by

$$g(n) = \mu((n, q)) \varphi((n, q)) \bar{\psi}(n).$$

Then we have:

$$\left. \begin{aligned} \mathcal{G}(\alpha, \chi) &= 0 \text{ if } R \nmid \alpha \\ \mathcal{G}(Rn, \chi) &= \mu(q) \psi(q) \tau(\psi) R g(n), \quad n = 1, 2, \dots \end{aligned} \right\} \quad (3)$$

2. PROOF OF THEOREM B

For a Dirichlet character $\chi \pmod{m}$ let the function $L(s, \chi)$ be given by

$$L(s, \chi) = \sum_1^{\infty} \chi(n) n^{-s}, \quad \operatorname{Re} s > 1.$$

The series defines an analytic function for $\operatorname{Re} s > 1$, which can be extended to a meromorphic function on the whole complex plane, with at most one simple pole at $s = 1$. If χ is primitive, then $L(s, \chi)$ satisfies the equation

$$L(1-s, \chi) = m^{s-1} (2\pi)^{-s} \Gamma(s) \left(e^{-\frac{\pi i s}{2}} + \chi(-1) e^{\frac{\pi i s}{2}} \right) \tau(\chi) L(s, \bar{\chi}). \quad (4)$$

Because of (1), this can also be written, for $\operatorname{Re} s > 1$, as

$$L(1-s, \chi) = m^{s-1} (2\pi)^{-s} \Gamma(s) \left(e^{-\frac{\pi i s}{2}} + \chi(-1) e^{\frac{\pi i s}{2}} \right) \sum_1^{\infty} \mathcal{G}(n, \chi) n^{-s}. \quad (5)$$

Whereas (4) holds only for primitive characters, (5) turns out to be valid in the general case. In fact, a much more general formula is proved in [3], th. 6.1; if we put there $x = \alpha = 0$, $\alpha_n = \chi(n)$ and observe that $\mathcal{G}(-n, \chi) = \chi(-1) \mathcal{G}(n, \chi)$, we get (5) immediately. But also most of the classical (= non-adelic) proofs of (4) will give (5) after very small changes. The only use of the primitivity of χ in these proofs is that they replace $\mathcal{G}(n, \chi)$ by $\bar{\chi}(n) \tau(\chi)$ at some stage (See for instance [7]).

Now let χ, m, ψ, f be as in the theorem. We have, by the Euler-product,

$$\begin{aligned} L(1-s, \chi) &= L(1-s, \psi) \prod_{p|q} \left(1 - \frac{\psi(p)}{p^{1-s}} \right) \\ &= L(1-s, \psi) \mu(q) q^{s-1} \psi(q) \prod_{p|q} (1 - p \bar{\psi}(p) p^{-s}) \end{aligned} \quad (6)$$

Formula (4) is valid if we write f for m and ψ for χ . Eliminating $L(1-s, \chi)$ and $L(1-s, \psi)$ out of the equations (4), (5), (6), and taking into account that $\chi(-1) = \psi(-1)$, we obtain after an easy calculation, for $\operatorname{Re} s > 1$:

$$\sum_1^\infty \mathcal{G}(n, \chi) n^{-s} = R \tau(\psi) \mu(q) \psi(q) R^{-s} L(s, \bar{\psi}) \prod_{p|q} \left(1 - \frac{p\bar{\psi}(p)}{p^s}\right) \quad (7)$$

Using Euler's product, we have for $\operatorname{Re} s > 1$

$$\begin{aligned} L(s, \bar{\psi}) \prod_{p|q} (1 - p\bar{\psi}(p) p^{-s}) \\ = \prod_{p \nmid q} \sum_{k=0}^\infty \bar{\psi}(p^k) p^{-ks} \cdot \prod_{p|q} (1 + (1-p) \sum_{k=1}^\infty \bar{\psi}(p^k) p^{-ks}) \\ = \prod_p \sum_{k=0}^\infty \bar{\psi}(p^k) \mu((p^k, q)) \varphi((p^k, q)) p^{-ks} \\ = \prod_p \sum_{k=0}^\infty g(p^k) p^{-ks} = \sum_1^\infty g(n) n^{-s}. \end{aligned}$$

If we put that into (7) we get

$$\sum_1^\infty \mathcal{G}(n, \chi) n^{-s} = R \tau(\psi) \mu(q) \psi(q) \sum_1^\infty g(n) (Rn)^{-s},$$

and we obtain (3) by comparing the coefficients.

3. In this section we show that one can prove theorem A using theorem B and conversely. Let $K(\alpha)$ resp. $H(\alpha)$ be the right hand side of (2) resp. (3). We want to prove $K(\alpha) = H(\alpha)$. First we show that $K(\alpha) \neq 0$ iff $H(\alpha) \neq 0$.

Now $H(\alpha) \neq 0$ iff $R|\alpha$ and $(\alpha R^{-1}, f) = 1$, and this is equivalent to

$$(\alpha f q, f m) = m \quad (8)$$

On the other hand $K(\alpha) \neq 0$ iff the four conditions hold

$$(\alpha, m)f | m \quad (9)$$

$$\frac{m}{(\alpha, m)f} \text{ is squarefree} \quad (10)$$

$$\left(\frac{m}{(\alpha, m)f}, f \right) = 1 \quad (11)$$

$$\left(\frac{\alpha}{(\alpha, m)}, f \right) = 1 \quad (12)$$

At several places in the following we will use that q is squarefree, $(f, q) = 1$ and the prime divisors of m are precisely the prime divisors of fq .

Let us assume (8). Then $f(\alpha, m) \mid f(\alpha q, m) = m$. Also $\frac{m}{f(\alpha, m)} = \frac{(\alpha q, m)}{(\alpha, m)} \mid q$. This proves (9) and (10). From $f = m/(\alpha q, m)$ we have $(\alpha/(\alpha, m), f) = (\alpha/(\alpha, m), m/(\alpha q, m)) \mid (\alpha/(\alpha, m), m/(\alpha, m)) = 1$.

This proves (12). Finally

$$\left(\frac{m}{(\alpha, m)f}, f \right) = \left(\frac{f(\alpha q, m)}{(\alpha, m)f}, f \right) \mid (q, f) = 1,$$

proving (11).

Conversely assume (9)-(12). From (10) and (11) we infer that $\frac{m}{f(\alpha, m)} \mid q$.

This implies

$$m \mid m \left(\frac{\alpha}{(\alpha, m)}, f \right) = \left(\frac{m\alpha f}{f(\alpha, m)}, mf \right) \mid (q\alpha f, mf). \quad (13)$$

Also

$$f(m, q\alpha) = f(\alpha, m) \left(\frac{m}{(\alpha, m)}, q \frac{\alpha}{(\alpha, m)} \right) = f(\alpha, m) \left(\frac{m}{(\alpha, m)}, q \right).$$

In the last term, the numbers f and $(m/(\alpha, m), q)$ both divide $m/(\alpha, m)$, because of (9), and they are coprime, hence their product divides $m/(\alpha, m)$. This gives $f(m, q\alpha) \mid m$. Together with (13) this implies (8).

It remains to prove that $H(\alpha) = K(\alpha)$ for $\alpha = Rn$, $(n, f) = 1$. We have $m_0 = m/(\alpha, m) = fq/(n, fq) = fq/(n, q)$. Hence

$$\begin{aligned} R &= \frac{m}{fq} = \frac{\varphi(m)}{\varphi(fq)} = \frac{\varphi(m)}{\varphi(f)\varphi(q)} = \frac{\varphi(m)}{\varphi(f)\varphi((n, q))\varphi(\frac{q}{(n, q)})} \\ &= \frac{\varphi(m)}{\varphi((n, q))\varphi(m_0)} \end{aligned}$$

hence

$$R\varphi((n, q)) = \varphi(m)/\varphi(m_0) \quad (14)$$

Also $\mu(q) = \mu((n, q))\mu(q/(n, q))$, so

$$\mu(q)\mu((n, q)) = \mu(q/(n, q)) = \mu(m_0/f). \quad (15)$$

Finally $\alpha_0 = n/(n, q)$, so $\alpha_0 q = nm_0/f$,

hence

$$\psi(\alpha_0)\psi(q) = \psi(m_0/f)\psi(n)$$

or

$$\bar{\psi}(n)\psi(q) = \bar{\psi}(\alpha_0)\psi\left(\frac{m_0}{f}\right) \quad (16)$$

Multiplying (14), (15) and (16) we find $K(\alpha) = H(\alpha)$.

4. SPECIAL CASES

(a) Theorem *B* implies that $\tau(\chi) \neq 0$ if and only if $R = 1$, that is, if and only if m/f is squarefree and has no common divisor with f . We have then

$$\tau(\chi) = \mu\left(\frac{m}{f}\right)\psi\left(\frac{m}{f}\right)\tau(\psi) \quad (17)$$

and

$$\mathcal{G}(\alpha, \chi) = g(\alpha)\tau(\chi) \quad (18)$$

On the other hand, if m/f is not square free or has a common divisor with f , then the right hand side of (17) is zero. So, (17) holds for any character χ . For another proof of this see [4], p. 148.

(b) If $\chi = \chi_0 =$ principal character mod m , then $f = 1$, $\psi \equiv 1$, $\tau(\psi) = 1$, $q = \tilde{m}$ = squarefree kernel of m , $R = m/\tilde{m}$, and $\mathcal{G}(\alpha, \chi_0) = C_m(\alpha)$ = RAMANUJANS SUM.

Theorem *B* gives the well-known formula:

$$C_m(\alpha) = 0 \quad \text{if} \quad \frac{m}{\tilde{m}} \not\mid \alpha$$

$$C_m\left(\frac{m}{\tilde{m}} n\right) = \frac{m}{\tilde{m}} \mu(\tilde{m}) \mu((n, \tilde{m})) \varphi((n, \tilde{m})).$$

From (17) we get for all m

$$C_m(1) = \mu(m).$$

5. Remarks: (a) It is clear that $\mathcal{G}(\alpha, \chi)$ cannot vanish identically. So by 4. (a), formula (1) can only hold if $R = 1$, and if $g(\alpha) = \bar{\chi}(\alpha)$ for all α . But this is only possible if $q = 1$, i.e. if $m = f$. This shows that (1) characterises primitive characters, a fact proved by T.M. Apostol [5].