Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	23 (1977)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	THE GELFAND-NAIMARK THEOREMS FOR C*-ALGEBRAS
Autor:	Doran, Robert S. / Wichmann, Josef
Kapitel:	1. Introduction
DOI:	https://doi.org/10.5169/seals-48924

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## **Download PDF:** 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

## THE GELFAND-NAIMARK THEOREMS FOR C\*-ALGEBRAS

by Robert S. DORAN and Josef WICHMANN

## 1. INTRODUCTION

Many of the Banach spaces which attract attention are at the same time algebras under some multiplication. In spite of this fact their study from this richer point of view was taken up only after the publication in 1932 of Banach's book [6]. One of the early fundamental results in the general theory of Banach algebras was a generalization of the classical theorem of Frobenius that any finite dimensional division algebra over the complex field is isomorphic to the field of complex numbers. S. Mazur [35] announced in 1938 that every complex normed division algebra is isomorphic to the field of complex numbers. S. Mazur [35] announced in 1938 that every complex normed division algebra is isomorphic to the field of complex numbers. Since the first published proof was given by I. M. Gelfand [22] this result is often called the Mazur-Gelfand theorem [43], [55]. As an immediate consequence one obtains the following beautiful characterization of the complex field among normed algebras: any normed algebra satisfying the norm condition  $||xy|| = ||x|| \cdot ||y||$  for all elements x and y is isometrically isomorphic to the field of complex numbers.

Many important Banach algebras carry a natural involution. In the case of an algebra of functions the involution is the operation of taking the complex-conjugate and in the case of an algebra of operators on a Hilbert space it is the operation of taking the adjoint operator. Motivated by these observations the Soviet mathematicians Israel M. Gelfand and Mark A. Naimark [23] proved, under some additional assumptions, the following two theorems:

THEOREM I. Let A be a commutative Banach algebra with involution satisfying  $||x^*x|| = ||x^*|| \cdot ||x||$  for all x in A. Then A is isometrically \*-isomorphic to  $C_0(X)$ , the algebra of all continuous complex-valued functions which vanish at infinity on some locally compact Hausdorff space X.

THEOREM II. Let A be a Banach algebra with involution satisfying  $||x^*x|| = ||x^*|| \cdot ||x||$  for all x in A. Then A is isometrically \*-iso-

The purpose of this paper is to present a thorough discussion of these two representation theorems. We shall trace, as carefully as we have been able, the interesting and rather tangled history which led to their present form. Then proofs of the theorems will be given. Finally, we shall survey some recent developments inspired by the theorems.

# 2. DEFINITIONS AND MOTIVATION

A \*-algebra is a complex associative linear algebra A with a mapping  $x \to x^*$  of A into itself such that for all  $x, y \in A$  and complex  $\lambda$ : (a)  $x^{**} = x$ ; (b)  $(\lambda x)^* = \overline{\lambda} x^*$ ; (c)  $(x+y)^* = x^* + y^*$ ; and (d)  $(xy)^* = y^* x^*$ . The map  $x \to x^*$  is called an *involution*; because of (a) it is clearly bijective. A subalgebra B of A is called a \*-subalgebra if  $x \in B$  implies  $x^* \in B$ .

An algebra which is also a Banach space satisfying  $||xy|| \le ||x|| \cdot ||y||$ for all x and y is called a *Banach algebra*. A Banach algebra which is also a \*-algebra is called a *Banach* \*-*algebra*. The involution in a Banach \*-algebra is said to be *continuous* if there is a constant M such that  $||x^*|| \le M ||x||$ for all x; the involution is *isometric* if  $||x^*|| = ||x||$  for all x.

A norm on a \*-algebra is said to satisfy the B\*-condition if  $||x^*x||$ =  $||x^*|| \cdot ||x||$  for all x; a B\*-algebra is a Banach \*-algebra whose norm satisfies the B\*-condition. A B\*-algebra with isometric involution clearly satisfies the condition  $||x^*x|| = ||x||^2$ . On the other hand, if A is a Banach \*-algebra satisfying  $||x||^2 \leq ||x^*x||$  (in particular if equality holds), then A is easily seen to be a B\*-algebra with isometric involution.

The Banach space C(X) of continuous complex-valued functions on a compact Hausdorff space is a commutative B\*-algebra under point-wise multiplication (fg)(t) = f(t)g(t), involution  $f^*(t) = \overline{f(t)}$ , and supnorm. Similarly, the algebra  $C_0(X)$  of continuous complex-valued functions which vanish at infinity on a locally compact Hausdorff space is a commutative B\*-algebra.

Examples of noncommutative B\*-algebras are provided by the algebra B(H) of bounded linear operators on a Hilbert space H. Multiplication in B(H) is operator composition, the involution  $T \to T^*$  is the usual adjoint operation, and the norm is the operator norm  $||T|| = \sup \{ ||T\xi|| : ||\xi|| \le 1, \xi \in H \}$ . A norm-closed \*-subalgebra of B(H) is called a C\*-algebra; clearly, every C\*-algebra is a B\*-algebra.