

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: REPRESENTATION OF COMPLETELY CONVEX FUNCTIONS BY THE EXTREME-POINT METHOD
Autor: Berg, Christian
Kapitel: 2. Determination of the extreme rays of W
DOI: <https://doi.org/10.5169/seals-48925>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. DETERMINATION OF THE EXTREME RAYS OF W

Inspired by [6] we consider the Green's function

$$G(x, t) = \begin{cases} (1-x)t & \text{for } 0 \leq t < x \leq 1, \\ (1-t)x & \text{for } 0 \leq x \leq t \leq 1. \end{cases}$$

If φ is a continuous function on $[0, 1]$ the unique solution $f \in C([0, 1]) \cap C^2([0, 1])$ to the equations

$$(2.1) \quad f'' = -\varphi \text{ in }]0, 1[, \quad f(0) = f(1) = 0$$

is

$$(2.2) \quad f(x) = \int_0^1 G(x, t) \varphi(t) dt.$$

The successive iterates of G are defined for $x, t \in [0, 1]$ by the equations

$$G_1(x, t) = G(x, t)$$

$$G_n(x, t) = \int_0^1 G(x, y) G_{n-1}(y, t) dy, \quad n \geq 2.$$

It is clear that $G_n(x, t) \geq 0$ for $x, t \in [0, 1]$.

We define recursively a sequence of polynomials $(A_n)_{n \geq 0}$ ¹⁾ by the requirement

$$(2.3) \quad A_0(x) = x, \quad A_n'' = -A_{n-1} \quad \text{and} \quad A_n(0) = A_n(1) = 0$$

for $n \geq 1$.

The polynomial A_n is of degree $(2n + 1)$, and we clearly have

$$(2.4) \quad A_n(x) = \int_0^1 G(x, t) A_{n-1}(t) dt = \int_0^1 G_n(x, t) t dt \quad \text{for}$$

$n \geq 1, x \in [0, 1].$

It follows that $A_n \geq 0$ on $[0, 1]$ for all n , and since $(-1)^k A_n^{(2k)} = A_{n-k}$ for $k \leq n$ we see that $A_n \in W$.

We recall that a ray \mathbf{R}_+x of a cone C is called *extreme*, if an equation $x = f + g$ with $f, g \in C$ is possible only if $f, g \in \mathbf{R}_+x$, cf. [3].

¹⁾ Our terminology is different from that of [6]; $(-1)^n A_n$ is equal to the n 'th Lidstone polynomial of [4] and [6].

PROPOSITION 2.1. *The polynomials Λ_n , $n \geq 0$, lie on extreme rays of W .*

Proof. If $\Lambda_0 = f + g$ with $f, g \in W$ we have $0 = f'' + g''$, but since f'' and g'' are both ≤ 0 , we conclude that f and g are affine. Furthermore, since $f(0) = g(0) = 0$, we conclude that f and g are proportional to Λ_0 .

Suppose now that Λ_{n-1} , $n \geq 1$, lies on an extreme ray of W , and assume that $\Lambda_n = f + g$ where $f, g \in W$. Then $\Lambda_{n-1} = -f'' + (-g'')$, and the induction hypothesis implies that $-f''$ and $-g''$ are proportional to Λ_{n-1} . Therefore we have $f = \lambda \Lambda_n(x) + ax + b$ for certain numbers $\lambda \geq 0$, a, b . Since $0 \leq f \leq \Lambda_n$, we have $f(0) = f(1) = 0$ which implies that $a = b = 0$. This proves that f (and similarly g) are proportional to Λ_n which then lies on an extreme ray of W .

Since $f \mapsto f^*$ is an affine isomorphism of W the polynomials Λ_n^* also lie on extreme rays of W . The following result is a special case of [6], Theorem 1.1.

PROPOSITION 2.2. *Every function $f \in W$ can for $n \geq 1$ be written as*

$$f(x) = \sum_{k=0}^{n-1} ((-1)^k f^{(2k)}(0) \Lambda_k^*(x) + (-1)^k f^{(2k)}(1) \Lambda_k(x)) + R_n(x),$$

where

$$R_n(x) = \int_0^1 G_n(x, t) (-1)^n f^{(2n)}(t) dt \in W.$$

Proof. For $n = 1$ the formula is equivalent with

$$(2.5) \quad f(x) - f(0)(1-x) - f(1)x = R_1(x) = - \int_0^1 G(x, t) f''(t) dt,$$

which follows directly from (2.2), and it is clear that $R_1 \in W$.

Suppose now the formula holds for some $n \geq 1$. Applying (2.5) to $(-1)^n f^{(2n)} \in W$ we get

$$\begin{aligned} (-1)^n f^{(2n)}(x) &= (-1)^n f^{(2n)}(0) \Lambda_0^*(x) + (-1)^n f^{(2n)}(1) \Lambda_0(x) \\ &\quad + \int_0^1 G(x, t) (-1)^{n+1} f^{(2n+2)}(t) dt, \end{aligned}$$

which substituted in the expression for R_n yields the formula for $n+1$ because of (2.4).

To see that $R_n \in W$ we notice that

$$(-1)^k R_n^{(2k)}(x) = \begin{cases} \int_0^1 G_{n-k}(x, t) (-1)^n f^{(2n)}(t) dt & \text{for } 0 \leq k \leq n-1 \\ (-1)^k f^{(2k)}(x) & \text{for } k \geq n. \end{cases}$$

The following lemma is easy to establish, but instead of giving the proof here we refer to [6].

LEMMA 2.3. *There exists a constant $M > 0$ such that*

$$0 \leq \int_0^1 G_n(x, t) dt \leq \frac{M}{\pi^{2n}} \quad \text{for } 0 \leq x \leq 1, n \geq 1.$$

PROPOSITION 2.4. *The only functions $f \in W$ satisfying $f^{(2k)}(0) = f^{(2k)}(1) = 0$ for all $k \geq 0$ are $f(x) = a \sin(\pi x)$ with $a \geq 0$.*

Proof. Suppose $f \in W$ satisfies $f^{(2k)}(0) = f^{(2k)}(1) = 0$ for all $k \geq 0$. Defining $a = \sup \{ \alpha \geq 0 \mid f - \alpha \sin(\pi x) \in W \}$, $g = f - a \sin(\pi x)$ belongs to W because W is closed in \mathbf{R}^I . Furthermore

$$g^{(2k)}(0) = g^{(2k)}(1) = 0 \text{ for all } k \geq 0.$$

Let $\varepsilon > 0$ be given. Since $\varphi = g - \varepsilon \sin(\pi x) \notin W$, there exist $k \geq 0$ and $x_0 \in]0, 1[$ such that $(-1)^k \varphi^{(2k)}(x_0) < 0$, hence

$$(-1)^k g^{(2k)}(x_0) < \varepsilon \pi^{2k} \sin(\pi x_0).$$

By Lemma 1.2 (iii) applied to $(-1)^k g^{(2k)}$ we get

$$(-1)^k g^{(2k)}(t) \leq \varepsilon \pi^{2k+1} \quad \text{for } 0 < t < 1,$$

and therefore by Proposition 2.2 and Lemma 2.3 for $0 < x < 1$

$$\begin{aligned} g(x) &= \int_0^1 G_k(x, t) (-1)^k g^{(2k)}(t) dt \leq \varepsilon \pi^{2k+1} \int_0^1 G_k(x, t) dt \\ &\leq \varepsilon M \pi. \end{aligned}$$

This proves that g is identically zero.

PROPOSITION 2.5. *The extreme rays of W are precisely the rays generated by A_n and $A_{n'}^*$, where $n \geq 0$, and $\sin(\pi x)$.*

Proof. We first show that $\sin(\pi x)$ lies on an extreme ray. If $\sin(\pi x) = f + g$ where $f, g \in W$, we have $f(0) = f(1) = g(0) = g(1) = 0$. Differentiating $2k$ times we similarly get $f^{(2k)}(0) = f^{(2k)}(1) = g^{(2k)}(0) = g^{(2k)}(1) = 0$, and it follows by Proposition 2.4 that f and g are proportional to $\sin(\pi x)$.

We finally have to show that an arbitrary extreme ray is generated by one of the above functions.

Assume that $f \in W$ generates an extreme ray. If $f^{(2k)}(0) = f^{(2k)}(1) = 0$ for all $k \geq 0$ we already know by Proposition 2.4 that f is proportional to $\sin(\pi x)$. Otherwise let n be the smallest number ≥ 0 for which $f^{(2n)}(0) \neq 0$ or $f^{(2n)}(1) \neq 0$. By Proposition 2.2 we then have

$$f(x) = (-1)^n f^{(2n)}(0) A_n^*(x) + (-1)^n f^{(2n)}(1) A_n(x) + R_{n+1}(x),$$

but since f lies on an extreme ray all three terms on the right-hand side lie on this ray.

If $f^{(2n)}(0) \neq 0$ this shows that $(-1)^n f^{(2n)}(1) A_n$ and R_{n+1} are proportional to A_n^* . Therefore $f^{(2n)}(1) = 0$ and $R_{n+1}^{(2n+2)} = f^{(2n+2)}$ is proportional to $(A_n^*)^{(2n+2)} = 0$, so that $f^{(2n+2)} = 0$ and hence $R_{n+1} = 0$ (cf. Proposition 2.2).

If $f^{(2n)}(1) \neq 0$ we similarly get $f^{(2n)}(0) = 0$ and $R_{n+1} = 0$. This shows that f lies on the ray generated by either A_n^* or A_n .

3. DETERMINATION OF A BASE FOR W

There are several ways of determining a base for W . We choose the following set

$$B = \left\{ f \in W \mid \int_0^1 f(x) \sin(\pi x) dx = 1 \right\}.$$

By Lemma 1.2 (ii) we get for $f \in B$ and $x_0 \in]0, 1[$ that

$$1 \geq \frac{1}{\pi} f(x_0) \int_0^1 \sin^2(\pi x) dx = \frac{1}{2\pi} f(x_0),$$

so the functions in B are uniformly bounded by 2π .

It is therefore clear that B is a compact convex base for W .

The extreme points of B are exactly the intersections between B and the extreme rays of W . We see that $2 \sin(\pi x) \in B$.

We claim that the following formulas hold, cf. [4]:

$$(3.1) \quad A_n^*(x) = \frac{2}{\pi^{2n+1}} \sum_{k=1}^{\infty} \frac{\sin(k\pi x)}{k^{2n+1}}, \quad n \geq 0, \quad x \in]0, 1[,$$

$$(3.2) \quad A_n(x) = \frac{2}{\pi^{2n+1}} \sum_{k=1}^{\infty} (-1)^{k+1} \frac{\sin(k\pi x)}{k^{2n+1}}, \quad n \geq 0, \quad x \in]0, 1[.$$