

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME CONTRIBUTIONS OF BENO ECKMANN TO THE DEVELOPMENT OF TOPOLOGY AND RELATED FIELDS
Autor: Hilton, Peter
Kapitel: 5. Simple homotopy type
DOI: <https://doi.org/10.5169/seals-48926>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. SIMPLE HOMOTOPY TYPE

Our final example of Eckmann's work brings us very much into the modern era. It will be understood why I have eschewed the temptation to deal with his very extensive contributions between 1958 and 1973 in detail. However, in 1970, he published a paper with Serge Maumary [70], dedicated to Georges de Rham, to which it will surely repay us to give some attention.

In 1950 Henry Whitehead recast in algebraic terms the theory of simple homotopy types which he had introduced many years earlier. In this theory one considers the collection of finite simplicial complexes¹⁾ homotopically equivalent to a given complex X and introduces the finer classification of *simple* equivalence into this collection. Whitehead showed that a single invariant sufficed to classify homotopy equivalences modulo simple equivalences; this invariant is now known as *Whitehead torsion* and is an element of an abelian group constructed functorially out of $\pi_1 X$. The importance of this theory has come to be recognized more clearly in recent years with the rise of differential topology and algebraic K -theory; indeed, the Whitehead group $\text{Wh } \pi$, to which the Whitehead torsion belongs ($\pi = \pi_1 X$) is a quotient of $K_1 \pi$. Chapman has proved the topological invariance of Whitehead torsion using the techniques of infinite-dimensional topology.

In [70] Eckmann and Maumary give a purely geometric description of the Whitehead group $\text{Wh } X$. They start with the classical description of a simple equivalence $s: Y \rightarrow Y'$ between finite cell complexes, as a sequence of elementary expansions and contractions. For a given finite cell complex X , they then introduce an equivalence relation into the family of maps emanating from X with target a finite cell complex. Thus $f: X \rightarrow Y$ and $f': X \rightarrow Y'$ are equivalent if there exists a simple equivalence $s: Y \rightarrow Y'$ such that $sf \simeq f'$. Let $A(X)$ be the set of equivalence classes thus defined. One may introduce a binary operation into $A(X)$ by means of the *homotopy push-out*—here we adopt a description equivalent to but not identical with that of [70]. Thus let $g: X \rightarrow Y$, $h: X \rightarrow Z$ be maps which we may assume cellular. Replace one or both of g , h by cofibrations; this may be done by the mapping cylinder construction. Thus, assume that g , h are cofibrations and construct the topological push-out of g , h —this will be the double mapping cylinder if g , h are embeddings in mapping cylinders:

¹⁾ There is no problem in replacing simplicial complexes by (finite) CW -complexes.

$$(5.1) \quad \begin{array}{ccc} & g & \\ X & \longrightarrow & Y \\ h \downarrow & & \downarrow u \\ Z & \longrightarrow & M \\ & v & \end{array}$$

Then if $\langle g \rangle$ stands for the class of g , etc., we define, from (5.1),

$$(5.2) \quad \langle g \rangle + \langle h \rangle = \langle ug \rangle$$

Next one makes A a functor to sets, also by means of the homotopy push-out: if $f: X \rightarrow X'$ we take a cofibration g in the class $\langle g \rangle$ and define

$$(5.3) \quad f_* \langle g \rangle = \langle \bar{g} \rangle$$

from the push-out

$$(5.4) \quad \begin{array}{ccc} & g & \\ X & \longrightarrow & Y \\ f \downarrow & & \downarrow \bar{f} \\ X' & \longrightarrow & Y' \\ & \bar{g} & \end{array}$$

The fact that f_* is a homomorphism with respect to the addition (5.2) is proved by purely categorical arguments; so too is the fact that $A(X)$ is an abelian monoid under the addition (5.2); of course the zero of $A(X)$ is $\langle 1_X \rangle$.

Now let $E(X)$ be the subset of $A(X)$ consisting of classes $\langle g \rangle$ such that g is a homotopy equivalence. It is plain that $E(X)$ is a submonoid; however, it is actually an abelian group. First, E is a functor, that is, $f_* EX \subseteq EX'$ if $f: X \rightarrow X'$. Second, let $g: X \rightarrow Y$ be a homotopy equivalence with homotopy inverse $g': Y \rightarrow X$ and set $\langle h \rangle = g'_* \langle g' \rangle$, $h: X \rightarrow Z$, so that $\langle h \rangle \in EX$. Then from (5.1)-(5.4) it follows that

$$\langle g \rangle + \langle h \rangle = \langle g' g \rangle = \langle 1_X \rangle = 0.$$

We have thus defined a functor $E: P \rightarrow Ab$, where P is the category of finite cell-complexes and Ab is the category of abelian groups. Finally, Eckmann and Maumary prove that EX depends only on the 2-skeleton of X ; a general argument then enables them to deduce that E factors through the fundamental group functor.

The gain in conceptual simplicity achieved by this geometric viewpoint is substantial; of course, the hard calculations remain to be done to compute the Whitehead group. One may compare the achievement of this paper with that of [34; 1953], in which Eckmann and Schopf produce a very significant simplification and clarification of the concept of injective hull of a module and a very easy, natural proof of its existence (first proved by Reinhold Baer); or with a very recent paper [81; 1976], in which Eckmann gave a remarkably simple proof of the Dyer-Vasquez theorem that the complement of a higher-dimensional knot $S^{n-2} \subseteq S^n$, $n \geq 4$, is never aspherical unless the knot group is infinite cyclic (thus, if $n \geq 5$, unless the knot is unknotted).

The story goes on. I have on my desk the latest manuscript, a joint paper by Eckmann and Bieri, completed in the spring of 1977, entitled “Relative Homology and Poincaré duality for group pairs”. As I have said, Beno Eckmann remains active and effective—but more is true. The Eckmann touch remains as sure as ever!

PUBLICATIONS OF B. ECKMANN

- [1] Zur Homotopietheorie gefaserter Räume. *Diss. Comm. Math. Helv.* 14 (1942), p. 141.
- [2] Über die Homotopiegruppen von Gruppenräumen. *Comm. Math. Helv.* 14 (1942), p. 234.
- [3] Systeme von Richtungsfeldern in Sphären und stetige Lösungen komplexer linearer Gleichungen. *Comm. Math. Helv.* 15 (1942), p. 1.
- [4] Vektorfelder in Sphären. *Vortrag Schweiz. Math. Ges., Verh. Schweiz. Naturf. Ges.* (1941), p. 85.
- [5] Über stetige Lösungen linearer Gleichungssysteme. *Vortrag Schweiz. Math. Ges., Verh. Schweiz. Naturf. Ges.* (1942), p. 78.
- [6] Über Zusammenhänge zwischen algebraischen und topologischen Problemen. *Vortrag Math. Ver. Bern, Verh. Naturf. Ges. Bern* (1942), p. 14.
- [7] L'idée de dimension, Leçon inaugurale Lausanne 5.2.43. *Revue de Théologie et Philosophie* 127 (1943), p. 1.
- [8] Stetige Lösungen linearer Gleichungssysteme. *Comm. Math. Helv.* 15 (1943), p. 318.
- [9] Gruppentheoretischer Beweis des Satzes von Hurwitz-Radon über die Komposition quadratischer Formen. *Comm. Math. Helv.* 15 (1943), p. 358.
- [10] Über monothetische Gruppen. *Verh. Schweiz. Naturf. Ges.* (1943), p. 63.
- [11] Topologie und Algebra, Antrittsvorlesung ETH 22.5.43. *Viertelj'schrift Naturf. Ges. Zürich*, 31 März 1944 (89), p. 25.
- [12] Über monothetische Gruppen. *Comm. Math. Helv.* 16 (1944), p. 249.
- [13] Harmonische Funktionen und Randwertaufgaben in einem Komplex. *Comm. Math. Helv.* 17 (1945), p. 240.
- [14] Lois de Kirchhoff et fonctions discrètes harmoniques. *Bull. Soc. Vaud. Sc. Nat.* 63, 264 (1945), p. 67.
- [15] Der Cohomologie-Ring einer beliebigen Gruppe. *Comm. Math. Helv.* 18 (1945/46), p. 232.