

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ALTERNATIVE HOMOTOPY THEORIES
Autor: James, I. M.
Kapitel: 4. Ex-HOMOTOPY THEORY
DOI: <https://doi.org/10.5169/seals-48928>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$(t_1 a'_1, \dots, t_n a'_n) \quad (a'_i \in A'_i, t_i \in I)$$

where $t_1^2 + \dots + t_n^2 = 1$. The radius vector through (t_1, \dots, t_n) meets the boundary of the n -cube I^n in a point (x_1, \dots, x_n) , say, where at least one coordinate is equal to 1. Thus a pointed G -homotopy equivalence

$$l: \Sigma (A'_1 * \dots * A'_n) \rightarrow \Sigma A'_1 \wedge \dots \wedge \Sigma A'_n$$

is given by

$$l((t_1 a'_1, \dots, t_n a'_n), s) = ((s x_1, a'_1), \dots, (s x_n, a'_n)).$$

Clearly l is equivariant with respect to the action of the symmetric group on the suspension of the multiple join and on the multiple smash product.

In particular, take $G = O(m)$ and $A'_i = S^{m-1}$, for all i . Let u be a permutation of the multiple join and v the corresponding permutation of the multiple smash product. We distinguish cases according as to whether the degree of the permutation is even or odd. In the even case u is G -homotopic to the identity 1_n on the n -fold join, using elementary rotations as before, and hence v is pointed G -homotopic to the identity 1_n on the n -fold smash product. In the odd case it follows similarly that u is G -homotopic to $1_{n-1} * a$, hence v is pointed G -homotopic to $1_{n-1} \wedge \hat{a}$. Taking $n = 3$, therefore, we see that the automorphisms which appear in (2.2) are trivial, in this example, and so

$$(3.3) \quad 3 [\Sigma * l_m, [\Sigma * l_m, \Sigma * l_m]] = 0$$

in $\pi_G(S^{3m+1}, S^{m+1})$, where l_m denotes the pointed $O(m)$ -homotopy class of the identity on S^m . It is easy to see, incidentally, that the Whitehead square $[\Sigma * l_m, \Sigma * l_m] \in \pi_G(S^{2m+1}, S^{m+1})$ is of infinite order, for all $m \geq 2$.


4. EX-HOMOTOPY THEORY

For our second example of an alternative homotopy theory we take the category of ex-spaces (see [7] for details), which is an enlargement of the category of sectioned bundles mentioned earlier. We recall that, with regard to a given space X , an *ex-space* consists of a space E together with maps

$$X \xrightarrow{\sigma} E \xrightarrow{\rho} X$$

such that $\rho\sigma = 1$. We refer to ρ as the *projection*, to σ as the *section*, and to (ρ, σ) as the *ex-structure*. Let E_i ($i = 1, 2$) be an ex-space with ex-structure

(ρ_i, σ_i) . We describe a map $f : E_1 \rightarrow E_2$ as an *ex-map* if $f\sigma_1 = \sigma_2, \rho_2 f = \rho_1$, as shown in the following diagram.



In particular we refer to $c = \sigma_2 \rho_1$ as the *trivial ex-map*. We also describe a homotopy $h_t : E_1 \rightarrow E_2$ as an *ex-homotopy* if h_t is an ex-map throughout. The set of ex-homotopy classes of ex-maps is denoted by $\pi_X(E_1, E_2)$ and the class of the trivial ex-map by 0.

In particular, suppose that E_i is a sectioned bundle with locally compact fibre. For each point $x \in X$ the fibre $\rho_i^{-1}(x)$ is equipped with basepoint $\sigma_i(x)$. Consider the fibre bundle $M = M_X(E_1, E_2)$ which is formed, in the usual way (see [2]) from the function-spaces of pointed maps $\rho_1^{-1}(x) \rightarrow \rho_2^{-1}(x)$. To each ex-map $f : E_1 \rightarrow E_2$ there corresponds a cross-section $f' : X \rightarrow M$, where $f'(x)$ is given by the restriction of f to the fibre over x , and conversely every such cross-section determines an ex-map. We shall exploit this correspondence in the next section.

Now let P be a principal G -bundle over X , where G is a topological group. For any pointed G -space A the pointed G -bundle $P_{\#}A$ can be regarded as an ex-space, and similarly with pointed G -maps. Thus $P_{\#}$ constitutes a functor from the category of pointed G -spaces to the category of ex-spaces, and determines a function

$$P_{\#} : \pi_G(A_1, A_2) \rightarrow \pi_X(E_1, E_2),$$

where A_i ($i = 1, 2$) is a pointed G -space and $E_i = P_{\#}A_i$. Of course, in general $P_{\#}$ is neither injective nor surjective.

As we have seen in § 1 a functor F in the category of pointed G -spaces defines a functor F in the category of sectioned G -bundles; in many cases such a functor can be extended to the category of ex-spaces. For example, the suspension functor Σ and the loop-space functor Ω can be so extended, also the binary functors product \times , wedge \vee , and smash \wedge . Similarly the notions of Hopf ex-space, etc.; can be introduced, following the standard formal procedure, so that $P_{\#}$ transforms Hopf G -spaces into Hopf ex-spaces, and so forth. Note that ΣE is cogroup-like and ΩE group-like, for any ex-space E .

The Whitehead product theory for ex-spaces has been worked out by Eggar [4]. His definition is such that if A, B, Y are as in §2 and $\alpha \in \pi_G(\Sigma A, Y), \beta \in \pi_G(\Sigma B, Y)$ then

$$(4.1) \quad [P_{\#}\alpha, P_{\#}\beta] = P_{\#}[\alpha, \beta]$$

in $\pi_X(\Sigma(P_{\#}A \wedge P_{\#}B), P_{\#}Y)$. Since we shall only be concerned with elements in the image of $P_{\#}$ we can introduce (4.1) as a piece of notation, without going into the details of Eggar's theory.

5. THE REGISTER THEOREM

In this section we suppose that X is a finite simply-connected CW -complex, although the results obtained can no doubt be generalized. We define the *register* $\text{reg}(X)$ of X to be the number of positive integers r such that, for some abelian group A , the cohomology group $H^r(X; A)$ is non-trivial. If X is a sphere, for example, then $\text{reg}(X) = 1$.

Let $p: M \rightarrow X$ be a fibration with fibre N . If a cross-section $s: X \rightarrow M$ exists then $sp: M \rightarrow M$ is a fibre-preserving map which is constant on the fibre. Conversely if $k: M \rightarrow M$ is a fibre-preserving map which is nulhomotopic on the fibre then M admits a cross-section as shown by Noakes [11]. We use similar arguments to prove

THEOREM (5.1). *Let $k: M \rightarrow M$ be a fibre-preserving map such that $l: N \rightarrow N$ is nulhomotopic, where $l = k|_N$, and let $s, t: X \rightarrow M$ be cross-sections. Then $k^r s$ and $k^r t$ are vertically homotopic, where $r = \text{reg}(X)$.*

The n -section ($n=0, 1, \dots$) of the complex X is denoted by X^n . Since X is connected we have a vertical homotopy of s into t over X^0 . This starts an induction. Suppose that for some $n \geq 1$ and some $q = q(n) \geq 1$ we have a vertical homotopy of $k^q s$ into $k^q t$ over X^{n-1} , so that the separation class

$$d = d(k^q s, k^q t) \in H^n(X; \pi_n(N))$$

is defined. If the cohomology group vanishes then $d = 0$ and $k^q s \simeq k^q t$ over X^n . But in any case the induced endomorphism l_* of $\pi_n(N)$ is trivial, by hypothesis, and so d lies in the kernel of the coefficient endomorphism $l_{\#}$ determined by l_* . Therefore

$$d(k^{q+1} s, k^{q+1} t) = l_{\#} d = 0,$$