**Zeitschrift:** L'Enseignement Mathématique
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique
**Band:** 23 (1977)
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ALTERNATIVE HOMOTOPY THEORIES
**Autor:** James, I. M.
**Kapitel:** 7. The adjoint G-bundle
**DOI:** https://doi.org/10.5169/seals-48928

**Nutzungsbedingungen**
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

**Conditions d'utilisation**
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

**Terms of use**
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 09.01.2026

**ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch**

Given a representation $\phi: SO(m) \to SO(q)$ write

$$J_\phi = J \circ \phi_*: \pi_r SO(m) \to \pi_{r+q}(S^q),$$

where $J$ denotes the usual Hopf-Whitehead homomorphism. For example, if $q > m$ and $\phi$ is the inclusion then

$$J_\phi = (-1)^{m-q} \Sigma_*^{m-q} J, \tag{6.2}$$

by (3.2) of [5] (cf. [8]). If $q = 2m$ and $\phi = 1 \oplus 1$ it is easily seen that

$$J_\phi = 2(-1)^m \Sigma_*^m J. \tag{6.3}$$

Consider the function-space $N = N(S^p, S^q)$ of pointed maps $S^p \to S^q$. We identify $\pi_i(N)$ $(i=0, 1, \ldots)$ with $\pi_{i+p}(S^q)$ in the standard way (see [15]). Let $G$ be a topological group and let

$$\phi: G \to SO(p), \quad \psi: G \to SO(q)$$

be representations of $G$. We regard $S^p$, $S^q$ as pointed $G$-spaces using $\phi$, $\psi$, respectively. Choose a principal $G$-bundle $P$ over $S^n$ with classifying element $\theta \in \pi_{n-1}(G)$, and take $E_1 = P_\#(S^p)$, $E_2 = P_\#(S^q)$. Then the operator $D$ in our exact sequence is given

$$D\alpha = \alpha \circ \Sigma_*^{r+p-q+1} J_\psi \theta - J_\phi \theta \circ \Sigma_*^{n+p-q-1} \alpha, \tag{6.4}$$

where $\alpha \in \pi_{r+p+1}(S^q)$. The case $r = 1$ of this result will be needed in §8 below.

## 7. The adjoint $G$-bundle

Let $X$ be any space and let $P$ be a principal $G$-bundle over $X$. We regard $P$ as a (right) $G$-space in the usual way. By a *principal automorphism* we mean an equivariant fibre-preserving map of $P$ into itself. By the *adjoint $G$-bundle* we mean the sectioned bundle $Q = P_\# G$, where $G$ acts on itself by conjugation. Note that $Q$ is a group ex-space since $G$ is a group $G$-space. We can construct $Q$ from $G \times P$ by identifying

$$(gag^{-1}, b) \sim (a, bg) \qquad (a \in G, b \in P) \tag{7.1}$$

for all $g \in G$. The group ex-structure is given by

$$\{a_1, b\} \cdot \{a_2, b\} = \{a_1 \cdot a_2, b\} \quad (a_1, a_2 \in G),$$

where $\{\ ,\ \}$ denotes the equivalence class of $(\ ,\ )$. Every principal automorphism $f$ of $P$ determines a cross-section $f': X \to Q$ as follows.

Given $x \in X$ choose any $b \in P_x$; then $fb = bg$, for some $g \in G$, and we define $f'x = \{ g, b \}$. This correspondence establishes an isomorphism between the group of principal automorphisms of $P$ and the group of cross-sections of $Q$.

Any element $c$ of the centre of $G$ determines a $G$-map $c_{\#}$ for any $G$-space $A$. Notice that $c_{\#}$ is a principal automorphism in the case of $P$ and that the corresponding cross-section $c'_{\#}$ of $Q$ is given by $c'_{\#} \{ b \} = \{ c, b \}$. When $X$ is a sphere these central cross-sections of $Q$ can be analysed as follows.

Take $X = S^n$ $(n \geqslant 2)$, so that $P$ is a principal $G$-bundle over $S^n$. Let $B^n$ denote the $n$-ball with boundary $S^{n-1}$. Choose a relative homeomorphism $(B^n, S^{n-1}) \to (S^n, x_0)$ and lift this to a map $k: (B^n, S^{n-1}) \to (P, G)$. The homotopy class $\theta \in \pi_{n-1}(G)$ of $l = k|S^{n-1}$ classifies the bundle according to clutching theory.

Let $c \in G$ be central and let $\lambda: I \to G$ be a path such that $\lambda(0) = e$, $\lambda(1) = c$. Consider the map $\Lambda: B^n \times I \to Q$ which is given by

$$\Lambda(y, t) = \{ \lambda(t), k(y) \} \qquad (y \in B^n, t \in I).$$

The boundary of $B^n \times I$ is the sphere

$$B^n \times 0 \cup S^{n-1} \times I \cup B^n \times 1,$$

and $\Lambda$ maps $S^{n-1} \times I$ into $G \subset Q$ by

$$\Lambda(y, t) = (ly) . (\lambda t) . (ly)^{-1},$$

using (7.1). Let us compare this with the map $\Lambda'$ of the boundary which agrees with $\Lambda$ on $B^n \times \dot{I}$ but is given on $S^{n-1} \times I$ by $\Lambda'(y, t) = \lambda t$. Now $\lambda$ can be regarded as a vertical homotopy of $e'_{\#}$ into $c'_{\#}$ over $\{ x_0 \}$ and $\Lambda$ represents the obstruction

$$\delta = \delta(e'_{\#}, c'_{\#}; \lambda) \in \pi_n(G)$$

to extending this vertical homotopy over $S^n$. Since $\Lambda \mid (B^n \times I)^{\cdot}$ is nul-homotopic, however, it follows that $\delta$ is also represented by $d: \tilde{\Sigma} S^{n-1} \to G$, where

$$d(y, t) = (ly) . (\lambda t) . (ly)^{-1}.$$

For example, take $G = SO(m)$, with $m$ even. Take $c = -e$ and

$$\lambda(t) = e \cos \pi t + b \sin \pi t \qquad (0 \leqslant t \leqslant 1),$$

where $b$ denotes the matrix

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \qquad (m/2 \quad \text{summands}).$$

Then $\delta = F\theta$, by definition, where

$$F: \pi_{n-1} SO(m) \to \pi_n SO(m)$$

denotes the Bott suspension, as in [6].

Now let $A_i$ $(i=1, 2)$ be a locally compact pointed $G$-space and write $E_i = P_{\#}A_i$. Recall that $N = N(A_1, A_2)$ denotes the function-space of pointed maps $A_1 \to A_2$. Given a pointed $G$-map $f: A_1 \to A_2$ we can construct an ex-map $P_{\#}f: E_1 \to E_2$ and a pointed $G$-map $\bar{f}: G \to N$, where $\bar{f}(g) = g_{\#} \circ f = f \circ g_{\#}$. I assert

PROPOSITION (7.2). *The ex-maps*

$$P_{\#}f, P_{\#}f \circ P_{\#}c: E_1 \to E_2$$

*are ex-homotopic if and only if*

$$\bar{f}^*\delta \in D\pi_1(N) \subset \pi_n(N),$$

*where $\delta$ is as above.*

Here $D$ is the operator which occurs in the modified exact sequence of the evaluation fibration derived from the function-space bundle, as in §6. The proof of (7.2) is by naturality, as follows.

First observe that $\bar{f}$ extends to a fibre-preserving map $\hat{f}: Q \to M$, where $M = M_X(E_1, E_2)$ denotes the function-space bundle. To see this we note that $f$ determines a pointed $G$-map $F: A_1 \times G \to A_2$, where

$$F(x, g) = f(xg) \qquad (x \in A_1, g \in G).$$

Hence $P_{\#}f: E_1 \times Q \to E_2$ is defined and we take $\hat{f}$ to be the adjoint.

We have $X = S^n$ so that the evaluation fibrations can be modified as in §6. Clearly

$$\Gamma_0(\hat{f}) \circ k \simeq l \circ \Omega^n(\bar{f}) \tag{7.3}$$

as shown below, where $k$ is defined by subtracting the cross-section $e'_{\#}$ and $l$ by subtracting $\hat{f} \circ e'_{\#}$.

$$\begin{array}{ccc} \Omega^n(G) & \xrightarrow{k} & \Gamma_0(Q) \\ {\scriptstyle \Omega^n(\bar{f})}\downarrow & & \downarrow{\scriptstyle \Gamma_0(\hat{f})} \\ \Omega^n(N) & \xrightarrow[l]{} & \Gamma_0(M) \end{array}$$

Hence we obtain a commutative diagram as follows, relating the modified exact sequences for $Q$ and $M$.

$$\begin{array}{ccc} \pi_n(G) & \xrightarrow{u_*} & \pi(\Gamma(Q)) \\ \bar{f}_* \Big\downarrow & & \Big\downarrow (\Gamma(\hat{f}))_* \\ \pi_n(N) & \xrightarrow[v_*]{} & \pi(\Gamma(M)) \end{array}$$

Recall that $\delta$ is the obstruction to extending $\lambda$ to a vertical homotopy of $e'_{\#}$ into $c'_{\#}$. Hence $\bar{f}_*\delta$ is the obstruction to extending $\bar{f} \circ \lambda$ to a vertical homotopy of $\hat{f} \circ e'_{\#}$ into $\hat{f} \circ c'_{\#}$. Hence it follows, as explained in the previous section, that $\hat{f} \circ e'_{\#}$ and $\hat{f} \circ c'_{\#}$ are vertically homotopic if and only if $\delta \in D\pi_1(N)$. Finally we use the correspondence between ex-maps and cross-sections to obtain (7.2) as stated.

## 8. Examples

Let $X$ be a finite simply-connected complex and let $P$ be a principal $SO(m)$-bundle over $X$. Consider the antipodal self-map $a$ of $S^{m-1}$. The unreduced suspension $\hat{a}$ is a pointed $SO(m)$-map of $S^m$ into itself. Hence $P_{\#}\hat{a}$ is an ex-map of $E = P_{\#}S^m$ into itself; let $\sigma \in \pi_X(E, E)$ denote the ex-homotopy class. Since $\hat{a}$ is of degree $(-1)^m$ we can apply (5.3) and obtain that

$$2^r \Sigma_* \sigma = 2^r \qquad (m \text{ even}), \tag{8.1}$$

where $r = \operatorname{reg}(X)$. It follows at once that

$$2^{r+1}[\iota_{\Sigma E}, \iota_{\Sigma E}] = 0 \qquad (m \text{ even}), \tag{8.2}$$

by (2.1) and (3.1), and hence from (3.3) that

$$[\iota_{\Sigma E}, [\iota_{\Sigma E}, \iota_{\Sigma E}]] = 0 \qquad (m \text{ even}). \tag{8.3}$$

Here $\iota_{\Sigma E}$ denotes the ex-homotopy class of the identity on $\Sigma E$. Similar results, but under more restrictive conditions, have been obtained by Eggar [4]. It can also be shown that the quadruple Whitehead products

$$[[\iota_{\Sigma E}, \iota_{\Sigma E}], [\iota_{\Sigma E}, \iota_{\Sigma E}]], [\iota_{\Sigma E}, [\iota_{\Sigma E}, [\iota_{\Sigma E}, \iota_{\Sigma E}]]]$$

are trivial, whether $m$ is even or odd.