

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 23 (1977)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ALTERNATIVE HOMOTOPY THEORIES
Autor: James, I. M.
Kapitel: 8. Examples
DOI: <https://doi.org/10.5169/seals-48928>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Hence we obtain a commutative diagram as follows, relating the modified exact sequences for Q and M .

$$\begin{array}{ccc} \pi_n(G) & \xrightarrow{u_*} & \pi(\Gamma(Q)) \\ \bar{f}_* \downarrow & & \downarrow (\hat{\Gamma(f)})_* \\ \pi_n(N) & \xrightarrow{v_*} & \pi(\Gamma(M)) \end{array}$$

Recall that δ is the obstruction to extending λ to a vertical homotopy of $e'_{\#}$ into $c'_{\#}$. Hence $\bar{f}_*\delta$ is the obstruction to extending $\bar{f} \circ \lambda$ to a vertical homotopy of $\hat{f} \circ e'_{\#}$ into $\hat{f} \circ c'_{\#}$. Hence it follows, as explained in the previous section, that $\hat{f} \circ e'_{\#}$ and $\hat{f} \circ c'_{\#}$ are vertically homotopic if and only if $\delta \in D\pi_1(N)$. Finally we use the correspondence between ex-maps and cross-sections to obtain (7.2) as stated.

8. EXAMPLES

Let X be a finite simply-connected complex and let P be a principal $SO(m)$ -bundle over X . Consider the antipodal self-map a of S^{m-1} . The unreduced suspension \hat{a} is a pointed $SO(m)$ -map of S^m into itself. Hence $P_{\#}\hat{a}$ is an ex-map of $E = P_{\#}S^m$ into itself; let $\sigma \in \pi_X(E, E)$ denote the ex-homotopy class. Since \hat{a} is of degree $(-1)^m$ we can apply (5.3) and obtain that

$$(8.1) \quad 2^r \Sigma_* \sigma = 2^r \quad (m \text{ even}),$$

where $r = \text{reg}(X)$. It follows at once that

$$(8.2) \quad 2^{r+1} [\iota_{\Sigma E}, \iota_{\Sigma E}] = 0 \quad (m \text{ even}),$$

by (2.1) and (3.1), and hence from (3.3) that

$$(8.3) \quad [\iota_{\Sigma E}, [\iota_{\Sigma E}, \iota_{\Sigma E}]] = 0 \quad (m \text{ even}).$$

Here $\iota_{\Sigma E}$ denotes the ex-homotopy class of the identity on ΣE . Similar results, but under more restrictive conditions, have been obtained by Eggar [4]. It can also be shown that the quadruple Whitehead products

$$[[[\iota_{\Sigma E}, \iota_{\Sigma E}], [\iota_{\Sigma E}, \iota_{\Sigma E}]], [\iota_{\Sigma E}, [\iota_{\Sigma E}, [\iota_{\Sigma E}, \iota_{\Sigma E}]]]]$$

are trivial, whether m is even or odd.

In particular, let X be a sphere. For m even (8.1) shows that $2\Sigma_*\sigma = 2$ and (8.2) that $4[\iota_{\Sigma E}, \iota_{\Sigma E}] = 0$. However, more precise results can be obtained by using the methods of §7, as follows. Take $X = S^n$ ($n \geq 2$) and let $\theta \in \pi_{n-1} SO(m)$ be the classifying element of P . We apply (7.2) with $f = 1$ and, using (6.4), obtain

THEOREM (8.4). *Let m be even. Then $\Sigma_*\sigma = 1$ if and only if $\Sigma_*JF\theta$ is contained in the image of*

$$D: \pi_{m+2}(S^{m+1}) \rightarrow \pi_{n+m+1}(S^{m+1}),$$

where

$$D\alpha = \Sigma_*J\theta \circ \Sigma_*^{n-1}\alpha - \alpha \circ \Sigma_*^2J\theta.$$

In the stable range, where $m > n$, the homomorphism D is trivial and $F\theta = \theta \circ \eta$, as in §6 of [6], where η generates the 1-stem. In this range it does not matter whether we deal with ex-maps or over-maps, and so (8.4) agrees with (4.5) of [8].

Now let ι_m denote the pointed $SO(m)$ -homotopy class of the identity on S^m , so that $\iota_{\Sigma E} = P_\# \Sigma_* \iota_m$. Represent the Whitehead square

$$w(\Sigma S^m) = [\Sigma_* \iota_m, \Sigma_* \iota_m]$$

by a pointed $SO(m)$ -map $f: \Sigma(S^m \wedge S^m) \rightarrow \Sigma S^m$. Then $P_\# f$ represents the Whitehead square

$$w(\Sigma E) = P_\# w(\Sigma S^m) = [\iota_{\Sigma E}, \iota_{\Sigma E}].$$

We apply (7.2) again and, using (6.2)-(6.4), obtain

THEOREM (8.5). *Let m be even. Then $2w(\Sigma E) = 0$ if and only if $w_{m+1} \circ \Sigma_*^{m+1} JF\theta$ lies in the image of*

$$D: \pi_{2m+2}(S^{m+1}) \rightarrow \pi_{n+2m+1}(S^{m+1}),$$

where $D\alpha = \alpha \circ \Sigma_*^{m+2}J\theta - 2\Sigma_*^m J\theta \circ \Sigma_*^{n-1}\alpha$.

Here $w_{m+1} \in \pi_{2m+1}(S^{m+1})$ denotes the ordinary Whitehead square of the generator of $\pi_{m+1}(S^{m+1})$. Unless $m = 2$ or 6 we have $w_{m+1} \neq 0$ and (8.5) determines the order of $w(\Sigma E)$. When $m = 2$ or 6 it would be interesting to know when $w(\Sigma E) = 0$, i.e. when ΣE is a Hopf ex-space, but unfortunately our method does not apply.

For some examples where the order of $w(\Sigma E)$ is (precisely) 4, consider the transgression $\Delta: \pi_n(S^m) \rightarrow \pi_{n-1} SO(m)$ in the homotopy exact

sequence of the standard fibration of $SO(m+1)$. Take $\theta = \Delta\phi$, where $\phi \in \Sigma_*\pi_{n-1}(S^{m-1})$. Then $\Sigma_*J\theta = 0$ and so D is trivial. However it follows from (4.1) and (6.3) of [6] that

$$\Sigma_*JF\theta = [\Sigma_*\iota_m, \Sigma_*\phi].$$

This Whitehead product is non-zero if, for example, $m = n$ and $\phi = \iota_m$ with $m \neq 2, 6$. Of course E is trivial as a bundle, in these examples, although not as a sectioned bundle.

REFERENCES

- [1] ARKOWITZ, M. The generalized Whitehead product. *Pacific J. of Math.* 12 (1962), 7-23.
- [2] BECKER, J. C. On the existence of A_k -structure on stable vector bundles. *Topology* 9 (1970), 367-384.
- [3] COHEN, D. E. Products and carrier theory. *Proc. London Math. Soc.* (3) 7 (1957), 219-248.
- [4] EGGAR, M. H. D. Phil. thesis (Oxford 1970).
- [5] JAMES, I. M. Reduced product spaces. *Ann. of Math.* 62 (1955), 170-197.
- [6] ——— On the Bott suspension. *J. of Math. Kyoto Univ.* 9 (1969), 161-188.
- [7] ——— Ex-homotopy theory. *Illinois J. of Math.* 15 (1971), 324-337.
- [8] ——— On sphere-bundles with certain properties. *Quart. J. of Math. Oxford* (2), 22 (1971), 353-370.
- [9] ——— On the maps of one fibre space into another. *Comp. Math.* 23 (1971), 317-328.
- [10] ——— Products between homotopy groups. *Comp. Math.* 23 (1971), 329-345.
- [11] NOAKES, J. L. D. Phil. thesis (Oxford 1974).
- [12] PUPPE, D. Homotopiemengen und ihre induzierten abbildungen I. *Math. Zeitschr.* 69 (1958), 299-344.
- [13] SEGAL, G. B. and JAMES, I. M. Note on equivariant homotopy type (*to appear*).
- [14] SUGAWARA, M. H-spaces and spaces of loops. *Math. J. Okayama Univ.* 5 (1955), 5-11.
- [15] WHITEHEAD, G. W. On products in homotopy groups. *Ann. of Math.* 47 (1946), 460-475.
- [16] ——— On mappings into group-like spaces. *Comm. Math. Helv.* 28 (1954), 320-328.
- [17] WHITEHEAD, J. H. C. On the homotopy type of ANR's. *Bull. Amer. Math. Soc.* 54 (1958), 1125-1132.
- [18] ——— On certain theorems of G. W. Whitehead. *Ann. of Math.* 58 (1953), 418-428.
- [19] WILSON, G. $S0(3)$ -equivariant maps of spheres. *Quart. J. of Math. Oxford* (2), 27 (1976), 263-265.

(Reçu le 6 mai 1977)

I. M. James

Oxford University Mathematical Institute