Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 24 (1978)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SINGULAR INTEGRAL EQUATION CONNECTED WITH QUASICONFORMAL MAPPINGS IN SPACE

Autor: Ahlfors, Lars V.

Kapitel: 3. Invariance properties

DOI: https://doi.org/10.5169/seals-49703

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

is the normalized and symmetrized Jacobian; it carries the quasiconformal data of the mapping.

The Riemannian metric $ds^2 = {}^t dx\,(MF)\,dx$ is conformally flat, a condition expressed by the vanishing of the conformal curvature tensor. For $n = 3$ this tensor is identically zero, but there is instead an integrability condition.

Let $F(x, t)$ be a one-parameter family of homeomorphisms such that $F(x, 0) = x$, $\dot{F}(x, 0) = f(x)$. Under suitable regularity conditions $(DF)^{\cdot}_0 = Df$, $(XF)^{\cdot}_0 = Df - \frac{1}{n}\, tr\, Df \cdot 1_n$, and $(MF)^{\cdot}_0 = Df + {}^t Df - \frac{2}{n}\, tr\, Df \cdot 1_n$. This motivates introducing the differential operator S defined by

$$(Sf)_{ij} = \frac{1}{2}(D_i f_j + D_j f_i) - \frac{1}{n}\,\delta_{ij}\, D_k f_k\,.$$

(The summation convention is in force in this paper). Note that Sf has values in SM_n.

There is a formal adjoint S^* which maps SM_n-valued functions on $\mathbf{R}^n$-valued functions. It is defined by

$$(S^*\varphi)_i = D_j \varphi_{ij}\,,$$

and it satisfies

$$\int_\Omega Sf \cdot \varphi dx = -\int_\Omega f \cdot S^*\varphi\, dx \tag{1}$$

when either f or φ has compact support. ($Sf.\,\varphi$ and $f \cdot S^*\varphi$ are the dot products $Sf_{ij}\,\varphi_{ij}$ and $f_i\,(S^*\varphi)_i$, respectively; dx is the euclidean volume element.)

Equation (1) defines Sf and $S^*\varphi$ as *distributions* even if f and φ are not differentiable. We are always assuming that f is continuous and φ locally integrable.

3. Invariance properties

In (1) we prefer to regard $\varphi\, dx$ as a matrix-valued measure, so that the pairing

$$< Sf, \varphi dx > = \int_\Omega Sf \cdot \varphi\, dx$$

is between a function and a measure. Similarly, $S^*(\varphi dx) = (S^*\varphi)\, dx$ is a vector-valued measure.

Let A be a Möbius transformation. We define the *pull-backs* of vector- and SM_n-valued functions by

$$(A^* f)(x) = (DA)^{-1} f(Ax)$$
$$(A^* \varphi)(x) = (DA)^{-1} \varphi(Ax) DA$$

and for the corresponding measures by

$$A^*(f\,dx) = |\det A|\, {}^t DA f(Ax)\, dx$$
$$A^*(\varphi\, dx) = |\det A|\, (DA)^{-1} \varphi(Ax) DA\,.$$

These definitions are chosen so that the pairings are invariant:

$$< A^* f, A^* g dx > = < f, g dx >$$
$$< A^* v, A^* \varphi\, dx > = < v, \varphi\, dx >\,.$$

There is a basic identity

$$S(A^* f)(x) = (DA)^{-1} Sf(Ax) DA \tag{2}$$

which may be expressed as a commutativity relation $SA^* = A^* S$, applicable to functions, but not to measures. It implies the relation $S^* A^* = A^* S^*$, which is valid for measures in the sense that

$$S^*(A^* \varphi\, dx) = A^*(S^* \varphi\, dx)\,, \tag{3}$$

but not for functions. It should be noted that (2) and (3) are true only because A is conformal.

A function is transformed into a measure by multiplication with a fixed invariant measure $\rho\, dx$. The invariance means that $A^*(\rho\, dx) = \rho\, dx$, or $\rho(Ax)\,|\det DA| = \rho(x)$; we assume also that A leaves Ω invariant. In these circumstances it makes sense to consider the operator $S^* \rho S$ which takes f to $S^*[\rho(Sf)dx]$ and commutes with A^* : $(S^* \rho S) A^* = A^*(S^* \rho S)$.

There are three classical cases in which Ω is invariant under a transitive group $G(\Omega)$ of Möbius transformations:

(i) $\Omega = \mathbf{R}^n$. $G(\Omega)$ is the group of euclidean motions, and $\rho = 1$.

(ii) $\Omega = \mathrm{B}(1) = \{ x : |x| < 1 \}$. $G = G(B)$ is the group of non-euclidean motions, and $\rho = (1 - |x|^2)^{-n}$.

(iii) Ω is the one-point compactification of $\mathbf{R}^n$, identified with S^n in $\mathbf{R}^{n+1}$. The group is formed by the rotations of the sphere, and $\rho = (1 + |x|^2)^{-n}$.