Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 24 (1978)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SINGULAR INTEGRAL EQUATION CONNECTED WITH QUASICONFORMAL MAPPINGS IN SPACE

Autor: Ahlfors, Lars V.

Kapitel: 5. FUNDAMENTAL SOLUTIONS

DOI: https://doi.org/10.5169/seals-49703

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 18.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

with

$$\Delta(x, y) = (1-2Q(y))(1-2Q(x-y^*)) = (1-2Q(y-x^*))(1-2Q(x)) .$$

Observe that $\Delta(x, y) = {}^t\Delta(y, x)$ and $\Delta(x, y)^2 = 1_n$ so that $\Delta(x, y) \in O(n)$. The matrix $\Delta(x, y)$ generalizes the angle arg $(1-\bar{x}y)/(1-\bar{y}x)$.

It is useful to note that $|Ax - Ay|^2 = |A'(x)|\,|A'(y)|\,|x-y|^2$ for any Möbius transformation A, and $[Ax, Ay]^2 = |A'(x)|\,|A'(y)|\,[x, y]^2$ if $A \in G$. There is an important relation between $T_y x$ and $T_x y$ expressed by

$$T_y x = -\Delta(x, y)\, T_x y . \tag{4}$$

We refer to [2, 3, 4, 5] for the elementary proofs of these formulas.

5. Fundamental solutions

A continuous mapping $f : B \to \mathbf{R}^n$ will be called a *deformation*. In this paper we shall assume, mainly for simplicity, that f is continuous on the boundary $S(1)$, and that $x \cdot f(x) = 0$ on $S(1)$; this means that f maps B on itself when regarded as an infinitesimal mapping.

A deformation is *trivial* if $Sf = 0$. There are very few trivial deformations: a complete list is given in [3].

It is customary to say that f is a *quasiconformal* deformation if $\|Sf\| \in L^\infty(B)$; here $\|Sf\|$ is the function whose value at x is the square norm of the matrix $Sf(x)$. More generally, we shall also consider functions with $\|Sf\| \in L^p(B)$; we abbreviate to $Sf \in L^p$, and we denote the L^p-norm of the square norm by $\|Sf\|_p$. The same convention will prevail for all matrix-valued functions.

We shall say that f is *harmonic* if $S^* \rho\, Sf = 0$, $\rho = (1-|x|^2)^{-n}$. Because of the invariance, if f is harmonic and $A \in G$, then $A^* f$ is also harmonic. Harmonicity in this sense is not the same as requiring the components to be harmonic with respect to the Poincaré metric.

There are n linearly independent solutions of the equation $S^* \gamma = 0$ which are homogeneous of degree $1-n$. We denote them by $\gamma_{..,k}$, $k = 1, \ldots, n$, the elements being

$$\gamma_{ij,k}(x) = |x|^{-n}(\delta_{ik}x_j + \delta_{jk}x_i - \delta_{ij}x_k) + (n-2)\,|x|^{-n-2}\, x_i x_j x_k .$$

There is a unique vector-valued function $g_{.k}(x)$ with components $g_{ik}(x)$ such that $g_{.k}(x) = 0$ for $|x| = 1$ and $\rho\, Sg_{.k} = \gamma_{..,k}$ so that

$S^* \rho S g_{.k} = 0$, or more precisely a Dirac distribution concentrated at 0. It is easy to see that $g = g_{ik}$, which we regard as a Green's matrix, will be of the form $g_{ik}(x) = a(|x|)\,\delta_{ik} + b(|x|)\,x_i x_k$; the explicit expressions for $a(r)$ and $b(r)$ are unimportant, except that g is of order $O((1-|x|^2)^{n+1})$ for $|x| \to 1$ and $O(|x|^{-n+2})$ for $x \to 0$ (if $n = 2$ the latter is replaced by $O(\log 1/|x|)$).

If $U \in O(n)$ it is immediate that $g(Ux) = Ug(x)^t U$. If we replace x by $T_x y$ and U by $-\Delta(x, y)$ it follows with the help of (4) that

$$\Delta(y, x)\, g(T_y x) = g(T_x y)\, \Delta(y, x)\,. \tag{5}$$

We now define the Green's matrix with singularity at y by

Definition 1.

$$\begin{aligned} g_{.k}(x, y) &= (1 - |y|^2)(T_y^* g_{.k})(x) = (1 - |y|^2)\, T_y'(x)^{-1}\, g(T_y x) \\ &= [x, y]^2\, \Delta(y, x)\, g(T_y x)\,. \end{aligned} \tag{6}$$

It is clear that $(S^* \rho S)_1\, g(x, y) = 0$ (the subscript indicates that the operator applies to the first variable). In view of (5) we can read off the symmetry property

LEMMA 1. $g(x, y) = {}^t g(y, x)$.

This symmetry plays a prominent role in H. Weyl's classical paper [9] which has been a strong inspiration for this work.

If $A \in G$ it is an easy consequence of (6) that

$$g(Ax, Ay) = A'(x)\, g(x, y)^t A'(y)$$

or, in a more suggestive form,

$$A_1^* A_2^*\, g(x, y) = g(x, y)\,,$$

where A_1^* is A^* applied to the first variable and the first index, and similarly for A_2^*.

Next we define

Definition 2.

$$\gamma_{..,k}(x, y) = \rho(x)\, S_1 g_{.k}(x, y) = (1 - |y|^2)\, \rho(x)\, (S_1 T_y^* g_{.k})(x).$$

It is evident by invariance that $S_1^*\, \gamma_{..,k}(x, y) = 0$. When x and y are transformed by the same $A \in G$ one finds

$$A_1^* A_2^*\, \gamma_{..,.}(x, y)\, dx = \gamma_{..,.}(x, y)\, dx$$

where A_1^* acts on x and the double index, A_2^* on y and the single index. For $A = T_y$ this leads to the explicit formula

$$\gamma_{..,k}(x, y) = \frac{(1 - |y|^2)^{n+1}}{[x, y]^{2n}} \Delta(y, x) \gamma_{..,k}(T_y x) \Delta(x, y) .$$

We note that $\gamma_{..,.}(x, 0) = \gamma_{..,.}(x)$ and $\gamma_{..,.}(0, y) = -(1 - |y|^2)^{n+1} \gamma_{..,.}(y)$.

We shall need to apply S to either variable in $\gamma_{..,.}(x, y)$. For this purpose we introduce

Definition 3. $\Gamma_{ij,hk}(x, y) = [S_2 \gamma_{ij,.}(x, y)]_{hk}$.

Because differentiations with respect to x and y commute it is clear that $S_1^* \Gamma_{..,hk}(x, y) = 0$. Moreover, starting from the relation $g_{ik}(x, y) = g_{ki}(y, x)$ it is not difficult to derive the following symmetry property:

LEMMA 2. $\rho(y) \Gamma_{ij,hk}(x, y) = \rho(x) \Gamma_{hk,ij}(y, x)$.

It follows, in particular, that $S_2^* \rho(y) \Gamma_{ij,..}(x, y) = 0$.

It is also important to know the asymptotic behavior of $\Gamma_{ij,hk}(x, y)$ when $x - y \to 0$. We observe first that

$$\begin{aligned} \rho(y) \Gamma_{ij,hk}(0, y) &= -(1 - |y|^2)^{-n} [S(1 - |y|^2)^{n+1} \gamma_{ij,.}(y)]_{hk} \\ &= -S_{ij,hk}(y) + R_{ij,hk}(y) \end{aligned}$$

where $S_{ij,hk}(y) = [S \gamma_{ij,.}(y)]_{hk}$ is homogeneous of degree $-n$ and $R_{ij,hk}(y)$ is homogeneous of degree $2 - n$. The explicit expression for $\Gamma_{ij,hk}(x, y)$ reads

$$\Gamma_{ij,..}(x, y) = \frac{(1 - |y|^2)^n}{[x, y]^{2n}} \Delta(x, y) \Gamma_{ij,..}(0, T_x y) \Delta(y, x) .$$

Elementary estimates show that

$$(7) \qquad |\Gamma_{ij,hk}(x, y) + S_{ij,hk}(x - y)| \leqq C_n |x - y|^{1-n} [x, y]^{-1}$$

with constant C_n.

6. POTENTIALS

Given an SM_n-valued function v on B we define its *potential* as the vector-valued function $I v$ with components

$$I v(y)_k = \int_B v_{ij}(x) \gamma_{ij,k}(x, y) \, dx .$$