

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SINGULAR INTEGRAL EQUATION CONNECTED WITH
QUASICONFORMAL MAPPINGS IN SPACE
Autor: Ahlfors, Lars V.
Kapitel: 6. POTENTIALS
DOI: <https://doi.org/10.5169/seals-49703>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

where A_1^* acts on x and the double index, A_2^* on y and the single index. For $A = T_y$ this leads to the explicit formula

$$\gamma_{\dots,k}(x, y) = \frac{(1 - |y|^2)^{n+1}}{[x, y]^{2n}} \Delta(y, x) \gamma_{\dots,k}(T_y x) \Delta(x, y).$$

We note that $\gamma_{\dots}(x, 0) = \gamma_{\dots}(x)$ and $\gamma_{\dots}(0, y) = -(1 - |y|^2)^{n+1} \gamma_{\dots}(y)$.

We shall need to apply S to either variable in $\gamma_{\dots}(x, y)$. For this purpose we introduce

Definition 3. $\Gamma_{ij,hk}(x, y) = [S_2 \gamma_{ij,\dots}(x, y)]_{hk}$.

Because differentiations with respect to x and y commute it is clear that $S_1^* \Gamma_{\dots,hk}(x, y) = 0$. Moreover, starting from the relation $g_{ik}(x, y) = g_{ki}(y, x)$ it is not difficult to derive the following symmetry property:

LEMMA 2. $\rho(y) \Gamma_{ij,hk}(x, y) = \rho(x) \Gamma_{hk,ij}(y, x)$.

It follows, in particular, that $S_2^* \rho(y) \Gamma_{ij,\dots}(x, y) = 0$.

It is also important to know the asymptotic behavior of $\Gamma_{ij,hk}(x, y)$ when $x - y \rightarrow 0$. We observe first that

$$\begin{aligned} \rho(y) \Gamma_{ij,hk}(0, y) &= -(1 - |y|^2)^{-n} [S(1 - |y|^2)^{n+1} \gamma_{ij,\dots}(y)]_{hk} \\ &= -S_{ij,hk}(y) + R_{ij,hk}(y) \end{aligned}$$

where $S_{ij,hk}(y) = [S \gamma_{ij,\dots}(y)]_{hk}$ is homogeneous of degree $-n$ and $R_{ij,hk}(y)$ is homogeneous of degree $2 - n$. The explicit expression for $\Gamma_{ij,hk}(x, y)$ reads

$$\Gamma_{ij,\dots}(x, y) = \frac{(1 - |y|^2)^n}{[x, y]^{2n}} \Delta(x, y) \Gamma_{ij,\dots}(0, T_x y) \Delta(y, x).$$

Elementary estimates show that

$$(7) \quad |\Gamma_{ij,hk}(x, y) + S_{ij,hk}(x - y)| \leq C_n |x - y|^{1-n} [x, y]^{-1}$$

with constant C_n .

6. POTENTIALS

Given an SM_n -valued function v on B we define its *potential* as the vector-valued function Iv with components

$$Iv(y)_k = \int_B v_{ij}(x) \gamma_{ij,k}(x, y) dx.$$

The integral converges if $v \in L^p(B)$ for some p with $n < p \leq \infty$. In fact, one proves that

$$|Iv(y)| \leq C_{n,p} \|v\|_p (1 - |y|)^{1-n/p}$$

if $p < \infty$ and

$$|Iv(y)| \leq C_n \|v\|_\infty (1 - |y|)(1 + \log 1/(1 - |y|))$$

if $p = \infty$. In any event $Iv(y)$ vanishes at a fixed rate for $|y| \rightarrow 1$.

The forming of the potential is an invariant operation in the sense that $IA^*v = A^*Iv$ for every $A \in G$. The potential is harmonic outside the support of v , for $(S^* \rho S)_2 \gamma_{ij..}(x, y) = 0$.

The following theorem serves to recover f from Sf and its boundary values:

THEOREM 1. *If $Sf \in L^p(B)$, $p > n$, then*

$$(8) \quad c_n f(y) = -ISf(y) + c_n Hf(y)$$

with

$$Hf(y) = \frac{1}{c_n} \int_{S(1)} \gamma_{ij..}(x, y) x_j f_i d\sigma(x).$$

Moreover, Hf is the unique harmonic function with the same boundary values as f , and if $x \cdot f = 0$ on $S(1)$ it can also be written in the form

$$Hf(y) = \frac{1}{c_n} \int_{S(1)} \frac{(1 - |y|^2)^{n+1}}{|x - y|^{2n}} \Delta(x, y) f(x) d\sigma(x).$$

Remarks. $d\sigma$ refers to the $(n-1)$ -dimensional measure on $S(1)$, and $c_n = 2(n-1)\omega_n/n$ where ω_n is the total measure of $S(1)$. We are assuming that f has a continuous extension to $S(1)$. Actually, this is automatically true if we assume the side condition in the form $x \cdot f(x) \rightarrow 0$ as $|x| \rightarrow 1$, for it can be shown that $Sf \in L^p$ forces f to satisfy a uniform Hölder condition.

The proof is a straight-forward application of Stokes' formula. The passage from the differentiable to the distributional case is elementary. The fact that a harmonic function is uniquely determined by its boundary values can be demonstrated as follows: Suppose that f is harmonic and zero on $S(1)$. It is readily shown that

$$\int_{S(r)} Sf(x)_{ij} \gamma_{ij,k}(x) d\sigma = 0$$

for all r . Therefore $ISf(0) = 0$ and hence $f(0) = 0$ by (8). If this result is applied to $(T_y^{-1})^* f$ it follows that $f(y) = 0$ for arbitrary y , so that f is indeed identically zero.

7. COMPUTATION OF SIv

It is easy to show that $S_{ij,hk}(y) = [S\gamma_{ij,\cdot}(y)]_{hk}$ is a Calderon-Zygmund kernel for any choice of the indices; in other words, it is homogeneous of degree $-n$, and its mean-value over the unit sphere is 0. If $v \in L^p$, $1 < p < \infty$, it follows by the Calderon-Zygmund theory that the principal value

$$\text{pr. v. } \int_B v_{ij}(x) S_{ij,hk}(x-y) dx$$

exists almost everywhere, and that it is the limit in $L^p(B)$ of the corresponding truncated integrals. In view of (7) it follows that the integral

$$(9) \quad \Gamma v(y)_{hk} = \int_B v_{ij}(x) \Gamma_{ij,hk}(x,y) dx$$

will also exist as a principal value almost everywhere. One finds, however, that the remainder in (7) makes it possible to assert merely that the principal value is a limit in $L^{p'}$ for any $p' < p/n$. In these circumstances it is natural to assume that $v \in L^p(B)$ for all $p \geq 1$.

THEOREM 2. *If $v \in L^p(B)$ with $p > n$, then $SIv \in L^{p'}(B)$ for all $1 \leq p' < p/n$, and*

$$(10) \quad SIv = -b_n v + \Gamma v$$

where $b_n = 4 \omega_n / (n+2)$ and Γv is defined by (9).

Proof. Let φ be an SM_n -valued test-function. The definition of SIv as a distribution leads to the following formal computation:

$$\begin{aligned} \int_B SIv(y)_{hk} \varphi(y)_{hk} dy &= - \int_B Iv(y)_k S^* \varphi(y)_k dy \\ &= - \int_B S^* \varphi(y)_k dy \int_B v_{ij}(x) \gamma_{ij,k}(x,y) dx \\ &= - \int_B v_{ij}(x) dx \int_B S^* \varphi(y)_k \gamma_{ij,k}(x,y) dy \\ &= - \int_B v_{ij}(x) dx [b_n \varphi_{ij}(x) - \int_B \varphi(y)_{hk} \Gamma_{ij,hk}(x,y) dy]. \end{aligned}$$