Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 24 (1978)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SINGULAR INTEGRAL EQUATION CONNECTED WITH QUASICONFORMAL MAPPINGS IN SPACE

Autor: Ahlfors, Lars V.

Kapitel: 7. Computation of SIv

DOI: https://doi.org/10.5169/seals-49703

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

for all r. Therefore $ISf(0) = 0$ and hence $f(0) = 0$ by (8). If this result is applied to $(T_y^{-1})^* f$ it follows that $f(y) = 0$ for arbitrary y, so that f is indeed identically zero.

7. Computation of SIv

It is easy to show that $S_{ij,hk}(y) = [S\gamma_{ij,\cdot}(y)]_{hk}$ is a Calderon-Zygmund kernel for any choice of the indices; in other words, it is homogeneous of degree $-n$, and its mean-value over the unit sphere is 0. If $v \in L^p$, $1 < p < \infty$, it follows by the Calderon-Zygmund theory that the principal value

$$\text{pr. v.} \int_B v_{ij}(x)\, S_{ij,hk}(x-y)\, dx$$

exists almost everywhere, and that it is the limit in $L^p(B)$ of the corresponding truncated integrals. In view of (7) it follows that the integral

$$\Gamma v(y)_{hk} = \int_B v_{ij}(x)\, \Gamma_{ij,hk}(x, y)\, dx \tag{9}$$

will also exist as a principal value almost everywhere. One finds, however, that the remainder in (7) makes it possible to assert merely that the principal value is a limit in $L^{p'}$ for any $p' < p/n$. In these circumstances it is natural to assume that $v \in L^p(B)$ for all $p \geqq 1$.

Theorem 2. *If* $v \in L^p(B)$ *with* $p > n$, *then* $SIv \in L^{p'}(B)$ *for all* $1 \leqslant p' < p/n$, *and*

$$SI\, v = -b_n v + \Gamma v \tag{10}$$

where $b_n = 4\,\omega_n/(n+2)$ *and* Γv *is defined by* (9).

Proof. Let φ be an SM_n-valued test-function. The definition of SIv as a distribution leads to the following formal computation:

$$\begin{aligned}
\int_B SI\, v(y)_{hk}\, \varphi(y)_{hk}\, dy &= -\int_B Iv(y)_k S^* \varphi(y)_k\, dy \\
&= -\int_B S^* \varphi(y)_k\, dy \int_B v_{ij}(x)\, \gamma_{ij,k}(x, y)\, dx \\
&= -\int_B v_{ij}(x)\, dx \int_B S^* \varphi(y)_k\, \gamma_{ij,k}(x, y)\, dy \\
= -\int_B v_{ij}(x)\, dx\, &\left[b_n \varphi_{ij}(x) - \int_B \varphi(y)_{hk}\, \Gamma_{ij,hk}(x, y)\, dy\right].
\end{aligned}$$

The justification, by means of the Zygmund-Calderon theory, is routine, and (10) follows.

Taken together, Theorems 1 and 2 lead to a very striking result:

THEOREM 3. *An* SM_n*-valued function* $v \in L^p(B)$, $p > n$, *is of the form* $v = Sf$ *with* $f = 0$ *on* $S(1)$ *if and only if it satisfies the homogeneous integral equation* $\Gamma v = -a_n v$ *with* $a_n = c_n - b_n = 2(n-2)(n+1)\omega_n / n(n+2)$.

Indeed, if v is of this form, Theorem 1 implies $c_n f = -Iv$, hence $c_n v = -SIv$, and consequently $\Gamma v = (b_n - c_n) v$ by Theorem 2. Conversely, if $\Gamma v = -a_n v$ then $SIv = -c_n v$ by (10), and $f = Iv$ vanishes on $S(1)$.

The point of Theorem 3 is that the solvability of $Sf = v$ (with an extra condition on f) has been reduced to an integral equation.

THEOREM 4. *For any* $v \in L^p(B)$, $p > n$, $S^* \rho [\Gamma v + a_n v] = 0$.

Proof. Let f be a vector-valued test-function. Theorem 3 applies to Sf, and we obtain by use of Lemma 2

$$\int_B S^* \rho \Gamma v \cdot f dx = -\int_B \rho(x) \Gamma v(x)_{ij} Sf(x)_{ij} dx$$

$$= -\int_B \rho(x) Sf(x)_{ij} dx \int_B v(y)_{hk} \Gamma_{hk,ij}(y, x) dy$$

$$= -\int_B \rho(y) v(y)_{hk} dy \int_B Sf(x)_{ij} \Gamma_{ij,hk}(x, y) dx$$

$$= -\int_B \rho(y) v(y)_{hk} \Gamma Sf(y)_{hk} dy = a_n \int_B \rho(y) v(y)_{hk} Sf(y)_{hk} dy$$

$$= -a_n \int_B S^* \rho v \cdot f dy$$

and hence $S^* \rho \Gamma v = -a_n S^* v$.

THEOREM 5. *Every* v *which is in all* $L^p(B)$ *has a unique representation in the form* $v = v' + v''$ *where* v' *and* v'' *are in all* $L^p(B)$ *while* v' *is in the image of* SI *and* v'' *is in the kernel of* $S^* \rho$.

As a consequence of Theorems 3 and 4 the representation is given by

$$c_n v = -SIv + (\Gamma v + a_n v).$$

It is unique, for if $SI = \Gamma v + a_n v$, then $S^* \rho SIv = 0$ so that Iv is harmonic and 0 on $S(1)$, hence identically zero.