Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INVARIANTS OF FINITE REFLECTION GROUPS
Autor: Flatto, Leopold
Kapitel: 1. Chevalley's Theorem
DOI: https://doi.org/10.5169/seals-49704

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

(1.10) and Lemma 1.2 yield

$$(1.11) \qquad \delta_m = \frac{1}{|G|} \sum_{\sigma \varepsilon G} (Tr\, \sigma)_m = \frac{1}{|G|} \sum_{\sigma \varepsilon G} \sum_{|a|=m} \omega^a(\sigma) .$$

Multiply both sides of (1.11) by t^m and sum over m from 0 to ∞. We get

$$\sum_{m=0}^{\infty} \delta_m t^m = \frac{1}{|G|} \sum_{m=0}^{\infty} \sum_{\sigma \varepsilon G} \sum_{|a|=m} \omega^a(\sigma) t^m$$

$$= \frac{1}{|G|} \sum_{\sigma \varepsilon G} \{ \sum_{m=0}^{\infty} \omega_1^m(\sigma) t^m \ldots \sum_{m=0}^{\infty} \omega_n^m(\sigma) t^m \}$$

$$= \frac{1}{|G|} \sum_{\sigma \varepsilon G} \frac{1}{(1-\omega_1(\sigma) t) \ldots (1-\omega_n(\sigma) t)}$$

CHAPTER II

INVARIANT THEORETIC CHARACTERIZATION OF FINITE REFLECTION GROUPS

1. Chevalley's Theorem

We showed in chapter I that we can always find a finite number of homogeneous invariants forming a basis for the invariants of G and that this set must contain at least n elements, where $n = \dim V$. We show that this lower bound is attained only for the finite reflection groups. We first define these groups.

Definition 2.1. Let σ be a linear transformation acting on the n-dimensional vector space V. σ is a reflection $\Leftrightarrow$ σ fixes an $n-1$ dimensional hyperplane π and σ is of finite order > 1. π is called the reflecting hyperplane (r.h.) of σ.

Remark. Choose $v \notin \pi$. and let $\sigma v = \zeta v + p$, $p \in \pi$. If $\zeta = 1$, then $\sigma^m v = v + mp$, contradicting that σ is of finite order. Hence $\zeta \neq 1$. Let $v' = v + (\zeta - 1)^{-1} p$ and choose $p_1, \ldots, p_{n-1}$ as a basis for π. Then $\sigma p_i = p_i$, $1 \leqslant i \leqslant n-1$, $\sigma v' = \zeta v'$. ζ is a root of 1 in k which is distinct from 1, as σ is of finite order > 1. Thus σ is a reflection iff relative to some basis, the matrix for σ is diagonal, $n-1$ of the diagonal entries equalling 1 and the remaining one equalling a root of 1 in k distinct from 1.

DEFINITION 2.2. G is a finite reflection group acting on $V \Leftrightarrow G$ is a finite group generated by reflections on V.

As an example of a finite reflection group, let $G = S_n$. It is well known that S_n is generated by transpositions. The transposition of the variables x_i, x_j $(i \neq j)$ fixes the hyperplane $x_i - x_j = 0$, so that it is a reflection.

We have the following result

THEOREM 2.1 (Chevalley [4]). *Let G be a finite reflection group acting on the n-dimensional vector space V. The invariants of G have a basis consisting of n homogeneous elements which are algebraically independent over k.*

Let $k[x]$ denote the ring of polynomials in $x_i, \ldots, x_n$ with coefficients in k. We prove the following.

LEMMA 2.1. Let $I_1, \ldots, I_m$ be invariant polynomials of G, $I_1 \notin (I_2, \ldots, I_m)$ = the ideal in $k[x]$ generated by $I_2, \ldots, I_m$. Suppose that $P_1 I_1 + \ldots + P_m I_m = 0$, the P_i's being polynomials with P_1 homogeneous. Then $P_1 \in \mathscr{I}$, where $\mathscr{I}$ is the ideal in $k[x]$ generated by the homogeneous invariants of positive degree.

Proof of Lemma 2.1. The proof proceeds by induction on $\deg P_1$. Suppose $\deg P_1 = 0$, so that $P_1 = c \in k$. If $c \neq 0$, then $I_1 \in (I_2, \ldots, I_m)$, contrary to assumption. Hence $c = 0 \Rightarrow P_1 \in \mathscr{I}$. Let $\deg P_1 = n > 0$. Let σ be a reflection in G and $L = 0$ the equation of its r.h. (L is a linear homogeneous polynomial). We have $P_1(x) I_1(x) + \ldots + P_m(x) I_m(x) = 0$, $P_1(\sigma x) I_1(x) + \ldots + P_m(\sigma x) I_m(x) = 0$. Hence $[P_1(\sigma x) - P_1(x)] I_1(x) + \ldots + [P_m(\sigma x) - P_m(x)] I_m(x)$. For $L(x) = 0$, $\sigma(x) = x$, so that $P_i(\sigma x) - P_i(x) = 0$ whenever $L(x) = 0$, $1 \leqslant i \leqslant m$. Since $L(x)$ is irreducible it follows that

$$\frac{P_i(\sigma x) - P_i(x)}{L(x)}$$

is a polynomial, $1 \leqslant i \leqslant m$. We have

$$\left[\frac{P_1(\sigma x) - P_1(x)}{L(x)}\right] I_1(x) + \ldots + \left[\frac{P_m(\sigma x) - P_m(x)}{L(x)}\right] I_m(x) = 0 .$$

$$\deg \left[\frac{P_1(\sigma x) - P_1(x)}{L(x)}\right] < \deg P_1(x) ,$$

so that by the induction hypothesis

$$\frac{P_1(\sigma x) - P_1(x)}{L(x)} \equiv 0 \pmod{\mathscr{I}}.$$

Hence $P_1(\sigma x) \equiv P_1(x) \pmod{\mathscr{I}}$. Since the σ's generate G, this congruence holds for $\sigma \in G$. We conclude that

$$P_1(x) \equiv \frac{1}{|G|} \sum_{\sigma \varepsilon G} P_1(\sigma x) \pmod{\mathscr{I}}.$$

The polynomial $\frac{1}{|G|} \sum_{\sigma \varepsilon G} P_1(\sigma x)$ is invariant and homogeneous of degree $n \geqslant 1$. Hence it $\in \mathscr{I}$, so that $P_1 \in \mathscr{I}$.

Proof of Theorem 2.1. We choose $I_1, \ldots, I_r$ to be homogeneous invariants of positive degree forming a minimal basis for $\mathscr{I}$. Hilbert's proof of Theorem 1.1 shows that $I_1, \ldots, I_r$ form a basis for the invariants of G. We show that $I_1, \ldots, I_r$ are algebraically independent, so that $r = n$.

Suppose, to the contrary, that $I_1, \ldots, I_r$ are algebraically dependent. Choose $H(y_1, \ldots, y_r)$ to be a polynomial of minimal positive degree so that $H(I_1(x), \ldots, I_r(x)) = 0$. Let x-degree of any monomial $y_1^{a_1} \ldots y_r^{a_r}$ be $d_1 a_1 + \ldots + d_r a_r$, where $d_i = \deg I_i$. We may assume that all x-degrees of the monomials appearing in H are the same. Let

$$H_i(x) = \frac{\partial H}{\partial y_i}(I_1(x), \ldots, I_r(x)), \quad 1 \leqslant i \leqslant r.$$

The H_i's are invariant homogeneous polynomials, as all monomials in H have equal x-degree. Since $H(y_1, \ldots, y_n)$ is of positive degree, some $\frac{\partial H}{\partial y_i} \neq 0$, It follows that the corresponding $H_i(x) \neq 0$, as H was chosen to be of minimal degree; i.e. not all H_i's $= 0$. We relabel indices so that $H_1, \ldots, H_s$, $1 \leqslant s \leqslant r$, are ideally independent (i.e. none of the H_i's is in the ideal generated by the others) and $H_{s+j} \in (H_1, \ldots, H_s)$. $1 \leqslant j \leqslant r - s$. Thus $H_{s+j} = \sum_{i=1}^{s} V_{ji} H_i$, $1 \leqslant j \leqslant r - s$, where each V_{ji} is a homogeneous polynomial of degree $d_i - d_{s+j}$ (V_{ji} is interpreted to be 0 if this degree is negative). Differentiating the relation $H(I_1(x), \ldots, I_r(x)) = 0$ with respect to x_k, we obtain

$$\text{(2.1)} \qquad \sum_{i=1}^{r} H_i \frac{\partial I_i}{\partial x_k} = \sum_{i=1}^{s} H_i \frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} H_{s+l} \frac{\partial I_{s+l}}{\partial x_k}$$

$$= \sum_{i=1}^{s} H_i \left[\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k} \right] = 0.$$

Since

$$\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k}$$

is homogeneous of degree $d_i - 1$, we conclude from Lemma 2.1 that

$$\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k} = \sum_{j=1}^{r} B_j I_j, \; 1 \leqslant i \leqslant s, \tag{2.2}$$

where the B_j's are homogeneous and each term in (2.2) is homogeneous of degree $d_i - 1$. This forces $B_i = 0$. Multiply both sides of (2.2) by x_k and sum over k. We conclude, by Euler's identity for homogeneous polynomials,

$$d_i I_i + \sum_{l=1}^{r-s} V_{li} d_{s+l} I_{s+l} = \sum_{j=1}^{r} A_j I_j, \tag{2.3}$$

the A_j's being homogeneous with $A_i = 0$.

(2.3) shows that $I_i \in (I_1, \ldots, I_{i-1}, I_{i+1}, \ldots, I_r)$, contradicting the minimality of the basis $I_1, \ldots, I_r$. Hence $I_1, \ldots, I_r$ are algebraically independent and $r = n$.

2. The Theorem of Shephard and Todd

We obtain in this section a converse to Chevalley's Theorem, thereby obtaining an invariant theoretical characterization of finite reflection groups. We first prove several preliminary results.

Lemma 2.2. Let H be a finite group of linear transformations acting on the n-dimensional space V and fixing the $n - 1$ dimensional hyperplane π. The elements of H have a common eigenvector $v \in V - \pi$. Let $\sigma(v) = \zeta(\sigma) v$, $\sigma \in H$. $\zeta(\sigma)$ is an isomorphism from H into the multiplicative group of the roots of unity in k. It follows that H is a cyclic group.

Remark. The above lemma is a consequence of Maschke's Theorem proven in section 2.3. We provide another proof below.

Proof. Let $\sigma_1 \in H$, $\sigma_1 \neq e$ (the identity of H). By the remark following Definition 2.1, there exists $v \in V - \pi$ such that $\sigma_1(v) = \zeta_1 v$, ζ_1 being a root of unity $\neq 1$. For $\sigma \in H$, let $\sigma(v) = \zeta(\sigma) v + p(\sigma)$, $\zeta(\sigma) \in k$ and $p(\sigma) \in \pi$. Let $\sigma^* = \sigma_1^{-1} \sigma^{-1} \sigma_1 \sigma$. Then $\sigma^*(v) = v + (1 - \zeta_1) p(\sigma)$. Since σ^* is of finite order, $(1 - \zeta_1) p(\sigma) = 0 \Rightarrow p(\sigma) = 0$. Hence $\sigma(v) = \zeta(\sigma) v$. $\zeta(\sigma)$ is clearly an isomorphism from H into U, the multiplicative group of