Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INVARIANTS OF FINITE REFLECTION GROUPS
Autor: Flatto, Leopold
Kapitel: 2. The Theorem of Shephard and Todd
DOI: https://doi.org/10.5169/seals-49704

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Since

$$\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k}$$

is homogeneous of degree $d_i - 1$, we conclude from Lemma 2.1 that

$$\frac{\partial I_i}{\partial x_k} + \sum_{l=1}^{r-s} V_{li} \frac{\partial I_{s+l}}{\partial x_k} = \sum_{j=1}^{r} B_j I_j, \; 1 \leqslant i \leqslant s, \tag{2.2}$$

where the B_j's are homogeneous and each term in (2.2) is homogeneous of degree $d_i - 1$. This forces $B_i = 0$. Multiply both sides of (2.2) by x_k and sum over k. We conclude, by Euler's identity for homogeneous polynomials,

$$d_i I_i + \sum_{l=1}^{r-s} V_{li} d_{s+l} I_{s+l} = \sum_{j=1}^{r} A_j I_j, \tag{2.3}$$

the A_j's being homogeneous with $A_i = 0$.

(2.3) shows that $I_i \in (I_1, \ldots, I_{i-1}, I_{i+1}, \ldots, I_r)$, contradicting the minimality of the basis $I_1, \ldots, I_r$. Hence $I_1, \ldots, I_r$ are algebraically independent and $r = n$.

2. The Theorem of Shephard and Todd

We obtain in this section a converse to Chevalley's Theorem, thereby obtaining an invariant theoretical characterization of finite reflection groups. We first prove several preliminary results.

Lemma 2.2. Let H be a finite group of linear transformations acting on the n-dimensional space V and fixing the $n - 1$ dimensional hyperplane π. The elements of H have a common eigenvector $v \in V - \pi$. Let $\sigma(v) = \zeta(\sigma) v$, $\sigma \in H$. $\zeta(\sigma)$ is an isomorphism from H into the multiplicative group of the roots of unity in k. It follows that H is a cyclic group.

Remark. The above lemma is a consequence of Maschke's Theorem proven in section 2.3. We provide another proof below.

Proof. Let $\sigma_1 \in H$, $\sigma_1 \neq e$ (the identity of H). By the remark following Definition 2.1, there exists $v \in V - \pi$ such that $\sigma_1(v) = \zeta_1 v$, ζ_1 being a root of unity $\neq 1$. For $\sigma \in H$, let $\sigma(v) = \zeta(\sigma) v + p(\sigma)$, $\zeta(\sigma) \in k$ and $p(\sigma) \in \pi$. Let $\sigma^* = \sigma_1^{-1} \sigma^{-1} \sigma_1 \sigma$. Then $\sigma^*(v) = v + (1 - \zeta_1) p(\sigma)$. Since σ^* is of finite order, $(1 - \zeta_1) p(\sigma) = 0 \Rightarrow p(\sigma) = 0$. Hence $\sigma(v) = \zeta(\sigma) v$. $\zeta(\sigma)$ is clearly an isomorphism from H into U, the multiplicative group of

the roots of unity in k. U is known to be cyclic ([22], Vol. 1, p. 112). It follows that $\zeta(H)$, a subgroup of U, is cyclic and so H is cyclic.

THEOREM 2.2. *Let G be a finite group acting on the n-dimensional space V. Let $I_1, \ldots, I_n$ be homogeneous polynomials forming a basis for the invariants of G. Let $d_1, \ldots, d_n$ be the respective degrees of $I_1, \ldots, I_n$. Then*

$$\prod_{i=1}^{n} d_i = |G|, \quad \sum_{i=1}^{n} (d_i - 1) = r \tag{2.4}$$

where r = number of reflections in G.

Proof. By Theorem 1.2, $I_1, \ldots, I_n$ are algebraically independent. Let $I(x)$ be a homogeneous invariant of degree m. Then I is a linear combination of the monomials $I_1^{a_1} \ldots I_n^{a_n}$ where $a_1 d_1 + \ldots a_n d_n = m$. Furthermore, these monomials are linearly independent over k, as $I_1, \ldots, I_n$ are algebraically independent over k. It follows that the dimension δ_m of homogeneous invariants of degree m = number of non-negative integer solutions to $a_1 d_1 + \ldots + a_n d_n = m$. Hence

$$\sum_{m=0}^{\infty} \delta_m t^m = \frac{1}{(1-t^{d_1}) \ldots (1-t^{d_n})}. \tag{2.5}$$

(1.9) and (2.5) yield

$$\frac{1}{|G|} \sum_{\sigma \varepsilon G} \frac{1}{(1-\omega_1(\sigma)t) \ldots (1-\omega_n(\sigma)t)} = \frac{1}{(1-t^{d_1}) \ldots (1-t^{d_n})} \tag{2.6}$$

Expand both sides of (2.6) in powers of $(1-t)$. Let $\mathscr{R}$ = set of reflections in G and $\zeta(\sigma)$ = eigenvalue of the reflection σ which $\neq 1$. We have

$$\begin{aligned} &\frac{1}{|G|} \sum_{\sigma \varepsilon G} \frac{1}{(1-\omega_1)(\sigma)t) \ldots (1-\omega_n(\sigma)t)} \\ &= \frac{1}{|G|} \frac{1}{(1-t)^n} + \frac{1}{|G|} \sum_{\sigma \varepsilon \mathscr{R}} \frac{1}{1-\zeta(\sigma)} \frac{1}{(1-t)^{n-1}} + \ldots \end{aligned} \tag{2.7}$$

$$\begin{aligned} &\frac{1}{(1-t^{d_1}) \ldots (1-t^{d_n})} = \prod_{i=1}^{n} \frac{1}{d_i(1-t) - \binom{d_i}{2}(1-t)^2 + \ldots \pm (1-t)^{d_i}} \\ &= \frac{1}{\prod_{i=1}^{n} d_i} \frac{1}{(1-t)^n} + \frac{\frac{1}{2} \sum_{i=1}^{n} (d_i - 1)}{\prod_{i=1}^{n} d_i} \frac{1}{(1-t)^{n-1}} + \ldots \end{aligned} \tag{2.8}$$

Equating coefficients of (2.7), (2.8), we get

$$\prod_{i=1}^{n} d_i = |G|, \quad \sum_{i=1}^{n} (d_i - 1) = 2 \sum_{\sigma \varepsilon \mathscr{R}} \frac{1}{1 - \zeta(\sigma)} . \tag{2.9}$$

We evaluate the sum

$$\sum_{\sigma \varepsilon \mathscr{R}} \frac{1}{1 - \zeta(\sigma)} :$$

Let π be any r.h. Let $H_\pi = \{\sigma \mid \sigma \in G \text{ and } \sigma \text{ fixes } \pi\}$. Thus H_π is the subgroup of G consisting of the identity and those reflections in G with r.h. π. Applying Lemma 2.2 to H_π, we conclude that there exists $v \notin \pi$ such that $\sigma(v) = \zeta(\sigma) v$ for $\sigma \in H_\pi$. Let $H'_\pi = H_\pi - \{e\}$. Since $\zeta(\sigma^{-1}) = (\zeta(\sigma))^{-1}$, we obtain

$$\sum_{\sigma \varepsilon H'_\pi} \frac{1}{1 - \zeta(\sigma)} = \sum_{\sigma \varepsilon H'_\pi} \frac{1}{1 - \zeta(\sigma^{-1})} \tag{2.10}$$

$$= \sum_{\sigma \varepsilon H'_\pi} \left(1 - \frac{1}{1 - \zeta(\sigma)}\right) = |H'_\pi| - \sum_{\sigma \varepsilon H'_\pi} \frac{1}{1 - \zeta(\sigma)} .$$

Hence

$$\sum_{\sigma \varepsilon H'_\pi} \frac{1}{1 - \zeta(\sigma)} = \frac{|H'_\pi|}{2} . \tag{2.11}$$

Summing both sides of (2.11) over all r.h. π, we get

$$\sum_{\sigma \varepsilon \mathscr{R}} \frac{1}{1 - \zeta(\sigma)} = \frac{r}{2} . \tag{2.12}$$

(2.9), (2.12) yield Theorem 2.2.

THEOREM 2.3. *Let $f_1, \ldots, f_n$ be polynomials in the variables $x_1, \ldots, x_n$. $f_1, \ldots, f_n$ are algebraically independent over $k \Leftrightarrow$*

$$\frac{\partial(f_1, \ldots, f_n)}{\partial(x_1, \ldots, x_n)} \neq 0 .$$

Proof. Suppose that $f_1, \ldots, f_n$ are algebraically independent. Then $G(f_1, \ldots, f_n) = 0$ for some polynomial $G = G(y_1, \ldots, y_n)$. Assume that $G(y_1, \ldots, y_n)$ is of minimal positive degree. Differentiating this relation with respect to x_j, we get

$$\sum_{i=1}^{n} \frac{\partial G}{\partial y_i}(f_1, \ldots, f_n) \frac{\partial f_i}{\partial x_j} = 0, \ 1 \leqslant j \leqslant n . \tag{2.13}$$

(2.13) is a system of linear equations (with coefficients in $k(x_1, \ldots, x_n)$) in the unknowns $H_i(x) = \frac{\partial G}{\partial y_i}(f_1, \ldots, f_n)$, $1 \leqslant i \leqslant n$. $\frac{\partial G}{\partial y_i} \neq 0$ for some i, as G is not constant, and $\deg \frac{\partial G}{\partial y_i} < \deg G$. It follows that the corresponding $H_i(x) \neq 0$. Thus the linear system (2.13) has a non-zero solution, so that its determinant

$$\frac{\partial (f_1, \ldots, f_n)}{\partial (x_1, \ldots, x_n)} \neq 0 .$$

Conversely, let $f_1, \ldots, f_n$ be algebraically independent. For each i, $x_i, f_1, \ldots, f_n$ are algebraically dependent. Hence there exists a polynomial $G_i(x_i, y_1, \ldots, y_n)$ of minimal positive degree in x_i such that $G_i(x_i, f_1, \ldots, f_n) = 0$. Differentiating these relations with respect to x_k, we get

$$\sum_{j=1}^{n} \frac{\partial G_i}{\partial y_j}(x_i, f_1, \ldots, f_n) \frac{\partial f_j}{\partial x_k} + \frac{\partial G_i}{\partial x_k}(x_i, f_1, \ldots, f_n)\, \delta_{ik}, \ 1 \leqslant k \leqslant n , \tag{2.14}$$

δ_{ik} denoting the Kronecker symbol. (2.14) may be rewritten in matrix notation as

$$\left(\frac{\partial G_i}{\partial y_j}\right) \cdot \left(\frac{\partial f_i}{\partial x_j}\right) = D \tag{2.15}$$

where the entries of D are

$$- \delta_{ij} \frac{\partial G_i}{\partial x_j} .$$

$\det D \neq 0$, as x_i – degree of $\frac{\partial G_i}{\partial x_i} < x_i$ – degree of G_i, $1 \leqslant i \leqslant n$.

It follows from (2.15) that $\frac{\partial (f_1, \ldots, f_n)}{\partial (x_1, \ldots, x_n)} \neq 0$.

THEOREM 2.4. (Shephard and Todd [19]). *Let G be a finite group acting on the n-dimensional space V. Suppose there exists a basis of n homogeneous polynomials for the invariants of G. Then G is a finite reflection group.*

Proof. Let H be the subgroup of G generated by the reflections in G. By assumption G has n basic homogeneous invariants which, by Theorem 1.2, are algebraically independent. Since H is a finite reflection group, we conclude from Chevalley's Theorem that H has n basic homogeneous invariants $J_1, \ldots, J_n$ which are algebraically independent. Each I_i is invariant under H so that $I_i = I_i(J_1, \ldots, J_n)$, the latter quantity denoting a polynomial in the J_i's. We may assume that $I_i(J_1, \ldots, J_n)$ is a linear combination of monomials $J_1^{a_1} \ldots J_n^{a_n}$ whose x-degree $=$ deg I_i. We have

$$\frac{\partial(I_1, \ldots, I_n)}{\partial(x_1, \ldots, x_n)} = \frac{\partial(I_1, \ldots, I_n)}{\partial(J_1, \ldots, J_n)} \cdot \frac{\partial(J_1, \ldots, J_n)}{\partial(x_1, \ldots, x_n)} \tag{2.16}$$

By Theorem 2.3,

$$\frac{\partial(I_1, \ldots, I_n)}{\partial(x_1, \ldots, x_n)} \neq 0$$

and (2.16) then shows that

$$\frac{\partial(I_1, \ldots, I_n)}{\partial(J_1, \ldots, J_n)} \neq 0 .$$

It follows that there is a rearrangement $k_1, \ldots, k_n$ of $1, \ldots, n$ so that

$$\frac{\partial I_{k_1}}{\partial J_1} \cdots \frac{\partial I_{k_n}}{\partial J_n} \neq 0 .$$

Hence $I_{k_i}(J_1, \ldots, J_n)$ is of positive degree in J_i and deg $I_{k_i} \geqslant$ deg J_i, $1 \leqslant i \leqslant n$. Applying Theorem 2.2 both to G and H, we obtain

$$\prod_{i=1}^{n} \deg J_i = |H|, \quad \prod_{i=1}^{n} \deg I_i = |G| \tag{2.17}$$

$$\sum_{i=1}^{n} (\deg J_i - 1) = \sum_{i=1}^{n} (\deg I_i - 1) = r \tag{2.18}$$

where r = number of reflections in G = number of reflections in H.

Since deg $I_{k_i} \geqslant$ deg J_i, $1 \leqslant i \leqslant n$, we conclude from (2.18) that deg $I_{k_i} =$ deg J_i, $1 \leqslant i \leqslant n$. Hence $\prod_{i=1}^{n} \deg I_i = \prod_{i=1}^{n} \deg J_i$, and we conclude from (2.17) that $|G| = |H|$. Thus $G = H$ and G is a finite reflection group.