

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CARTIER DUALITY AND FORMAL GROUPS OVER Z
Autor: Rotman, Joseph
Kapitel: §2. Groups
DOI: <https://doi.org/10.5169/seals-49705>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

- (i) *The contravariant functor $\mathcal{A} \rightarrow \mathcal{B}$ given by $A \mapsto A^*$ is an antiequivalence of categories taking products to coproducts and final objects to initial objects.*
- (ii) *The restriction of this functor is an equivalence $(C\mathcal{A})^{op} \rightarrow G\mathcal{B}$.*

Several remarks are in order. First, we shall not define “linearly compact”; its role is to guarantee that A and A^{**} are isomorphic vector spaces, and this is false for discrete infinite dimensional spaces. Second, the proof of (ii) is a routine inspection of the various diagrams, once statement (i) has been proved.

There are at least two papers giving a Cartier duality between certain categories of commutative topological k -algebras and of cocommutative k -coalgebras, where k is a commutative ring. (Ditters [2]; Morris and Pareigis [5]). We present a version of Cartier duality between certain commutative Z -algebras (= commutative rings) and cocommutative Z -coalgebras; actually, our proof works if one replaces Z by any principal ideal domain that is neither a field nor a complete discrete valuation ring. Thus, our theorem is weaker than those of Ditters and Morris-Pareigis in that the ground rings k are restricted; it is stronger than their results in that we need not assume the algebras are topological algebras. Indeed, it is easy to see our category of commutative algebras is a proper, full subcategory of the corresponding categories of Ditters and of Morris-Pareigis. We add that our proof is quite easy and all details are given.

§2. GROUPS

All groups are abelian and are written additively.

DEFINITION. A subgroup A' of a group A is *cofinite* if A/A' is f.g. free (f.g. abbreviates “finitely generated”).

Of course, A' cofinite implies $A = A' \oplus A''$, where $A'' \cong A/A'$.

DEFINITION. The *cofinite topology* on a group A is that (linear) topology having a fundamental system of neighborhoods of 0 consisting of all cofinite subgroups of A .

It is clear that A is a topological group in the cofinite topology.

Suppose $A = Z^I$ for some index set I . We may also topologize A with the *product topology*, i.e., equip each factor Z with the discrete topology and consider A in the corresponding product topology. The first lemma shows that the cofinite topology gives a coordinate-free description of the product topology.

LEMMA 1. If $A = Z^I$ and I is countable, then the cofinite topology coincides with the product topology.

Proof. It is easy to see that, in either topology (and for any index sets I and J), every homomorphism $f: Z^I \rightarrow Z^J$ is sequentially continuous (if $x_n \rightarrow x$, then $f(x_n) \rightarrow f(x)$); if we assume I and J countable, then Z^I and Z^J are first countable (even metrizable), and so f is continuous.

Assume A' is cofinite in A , and A has the product topology. For finite n , we see Z^n is discrete (in either topology), whence the natural map $\pi: A \rightarrow A/A' \cong Z^n$ is continuous and $A' = \pi^{-1}(\{0\})$ is open.

Now assume A has the cofinite topology. If $U_i = \prod_{j \in I} X_j$, where $X_j = Z$ if $j \neq i$ and $X_j = \{0\}$ if $j = i$, then U_i is cofinite, hence open. It follows easily that every basic open set in the product topology is open in cofinite topology.

One may prove that Lemma 1 is true for any set I whose cardinal is nonmeasurable [6].

DEFINITION. The *completion* of a group A is $\varprojlim A/A'$, where A' ranges over all cofinite subgroups of A ; we denote $\varprojlim A/A'$ by A^\wedge . There is a canonical map $\lambda: A \rightarrow A^\wedge$; we say A is *complete* if λ is an isomorphism.

COROLLARY 2. If $A = Z^I$, where I is countable, then A is complete.

Proof: It is easy to see that, in the product topology, A is complete in the usual metric. By Lemma 1 and [4, Theorem 13.7], the two notions of completeness coincide.

The following remarkable result of Łos is the reason we need not mention linear compactness. Let us denote $\text{Hom}_Z(A, Z)$ by A^* .

LEMMA 3. (*Łos*)

- (i) Let $A = Z^N = \prod_{n=1}^{\infty} \langle e_n \rangle$. If $G = Z$ or $G = Z^{(I)}$, the direct sum of card I copies of Z , then the map $f \mapsto (f|_{\langle e_n \rangle})$ is an isomorphism $\text{Hom}_Z(A, G) \xrightarrow{\sim} \sum_{n=1}^{\infty} \text{Hom}_Z(\langle e_n \rangle, G)$.
- (ii) If I is countable, then $(Z^I)^* \cong Z^{(I)}$.
- (iii) If I is countable and either $A = Z^I$ or $A = Z^{(I)}$, then A is reflexive in the sense that the natural map $A \rightarrow A^{**}$ is an isomorphism.

Proof: [4; §94]. This Lemma is true if Z is replaced by any principal ideal domain that is neither a field nor a complete discrete valuation ring.

Again the countability assumption is too strong; one only needs the cardinal of I nonmeasurable. Also, part (i) is true for groups G other than Z and $Z^{(I)}$, namely, “slender” groups.

For any index sets I and J , there is a natural imbedding $Z^I \otimes Z^J \rightarrow Z^{I \times J}$ given by $(m_i) \otimes (n_j) \mapsto (m_i \otimes n_j)$.

LEMMA 4. *Assume I and J are countable. Then if $Z^I \otimes Z^J$ and $Z^{I \times J}$ are given the cofinite topology, then $Z^I \otimes Z^J$ is a dense subspace of $Z^{I \times J}$.*

Proof: By “subspace” we mean that the cofinite topology on $Z^I \otimes Z^J$ coincides with the relative topology $Z^I \otimes Z^J$ inherits from the larger space $Z^{I \times J}$. Let us write $A = Z^I \otimes Z^J$ and $G = Z^{I \times J}$. If G' is cofinite in G , then

$$A/G' \cap A \cong (A + G')/G' \subset G/G' ,$$

whence $G' \cap A$ is cofinite in A . Assume that A' is cofinite in A . Now A' is cofinite in A if and only if there are finitely many $f_i \in A^*$ with $A' = \cap \ker f_i$. Moreover, if $f \in A^*$ and $A' = \ker f$, then there exists a cofinite G' in G with $G' \cap A = A'$ if and only if there is $\tilde{f} \in G^*$ extending f . Thus it suffices to show we may extend $f \in (Z^I \otimes Z^J)^*$ to $\tilde{f} \in (Z^{I \times J})^*$. But this follows easily from the adjoint isomorphism and Lemma 3:

$$\begin{aligned} \text{Hom}(Z^I \otimes Z^J, Z) &= \text{Hom}(Z^I, \text{Hom}(Z^J, Z)) \\ &= \text{Hom}(Z^I, Z^{(J)}) \\ &= Z^{(I \times J)} = \text{Hom}(Z^{I \times J}, Z) . \end{aligned}$$

We have shown that $Z^I \otimes Z^J$ is a subspace of $Z^{I \times J}$; it is dense because it contains the dense subgroup $Z^{(I \times J)}$.

We remark that Lemma 4 is false for some subgroups of $Z^{I \times J}$; for example, if $A = Z^{(I \times J)} \oplus \langle x \rangle$, where x has each coordinate 1, then $Z^{(I \times J)}$ is cofinite in A ; the corresponding functional f on A cannot extend to $Z^{I \times J}$, for every $\tilde{f} \in (Z^{I \times J})^*$ that vanishes on $Z^{(I \times J)}$ must be 0 [4; Theorem 94.4].

LEMMA 5. *If I and J are countable, there is a natural isomorphism*

$$(Z^I \otimes Z^J)^\wedge \cong (Z^{(I)} \otimes Z^{(J)})^* .$$

(Recall: $^\wedge$ means completion and $*$ means dual space).

Proof: Since $Z^{(I)} \otimes Z^{(J)} \cong Z^{(I \times J)}$, the right hand side is $Z^{I \times J}$. By Lemma 4, $Z^I \otimes Z^J$ is a dense subspace of $Z^{I \times J}$, so that both have the same completion. This finishes the argument, for $Z^{I \times J}$ is complete, by Corollary 2.

COROLLARY 6. *If I and J are countable, then $(Z^I \otimes Z^J)^\wedge \cong Z^K$, where K is countable.*

Proof: Indeed, we have just seen that we may take $K = I \times J$.

LEMMA 7. *Assume A and B torsion-free. If A' is cofinite in A and B' is cofinite in B , then there is a natural isomorphism*

$$A \otimes B / (A' \otimes B + A \otimes B') \cong A/A' \otimes B/B'.$$

Proof: Define $\theta: A \otimes B \rightarrow A/A' \otimes B/B'$ by $a \otimes b \mapsto \bar{a} \otimes \bar{b}$ (where bar denotes appropriate coset); let $K = \ker \theta$. As A and B are torsion-free, they are \mathbb{Z} -flat, and so there is a commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \longrightarrow & A \otimes B & \xrightarrow{\theta} & A/A' \otimes B/B' \longrightarrow 0 \\ & & \uparrow & & \uparrow = & & \uparrow \\ & & 0 \rightarrow A' \otimes B + A \otimes B' \rightarrow A \otimes B \rightarrow A \otimes B / (A' \otimes B + A \otimes B') \rightarrow 0 & & & & \end{array}$$

The dotted arrow exists and is an epimorphism, by diagram-chasing; it is an isomorphism because both right hand terms are f.g. free of the same rank (to compute the bottom quotient, observe that $A = A' \oplus A''$, $B = B' \oplus B''$, where $A'' \cong A/A'$ and $B'' \cong B/B'$).

LEMMA 8. *Let $A = \mathbb{Z}^I$ and $B = \mathbb{Z}^J$, where I and J are countable. The subgroups of $A \otimes B$ of the form $A' \otimes B + A \otimes B'$, where A' is cofinite in A and B' is cofinite in B , form a fundamental system of neighborhoods at 0 for the cofinite topology of $A \otimes B$.*

Proof: First of all, Lemma 7 shows that each of these special subgroups of $A \otimes B$ is cofinite.

Next, assume C is cofinite in $A \otimes B$, so there is an exact sequence

$$0 \longrightarrow C \longrightarrow A \otimes B \xrightarrow{\theta} F \longrightarrow 0$$

with F f.g. free. Define $A' = \{a \in A: \theta(a \otimes b) = 0 \text{ for all } b \in B\}$ and, similarly, $B' = \{b \in B: \theta(a \otimes b) = 0 \text{ for all } a \in A\}$. Clearly $A' \otimes B + A \otimes B' \subset C$. Now A' is pure in A and B' is pure in B , so that A/A' and B/B' are torsion-free. Also, A' is closed in A (and B' is closed in B)

because θ is continuous (I and J are countable), so that A/A' is complete. By considering maximal independent subsets of A and B and observing that only finitely many elements of A are involved in lifting a (finite) basis of F , we see that A/A' has finite rank (similarly for B/B'). As the only finite rank complete groups are f.g. free, it follows that A' and B' are cofinite.

§3. FORMAL GROUPS

DEFINITION. Let \mathcal{A} denote the category of all commutative rings with 1 whose underlying additive group is of the form \mathbb{Z}^I , where $\text{card } I \leq \aleph_0$.

Note that $\mathbb{Z}[[x_1, \dots, x_n]]$, formal power series over \mathbb{Z} in n variables, is an object of \mathcal{A} . Further, \mathcal{A} has an initial object, namely, \mathbb{Z} .

LEMMA 9. *Every $A \in \text{obj } \mathcal{A}$ is a complete topological ring in the cofinite topology.*

Proof: By Lemma 1 and Corollary 2, we know A is a complete topological group. It remains to show that multiplication $m: A \times A \rightarrow A$ is continuous, and, for this it suffices to prove the corresponding homomorphism $m': A \otimes A \rightarrow A$ is continuous; this is so because every homomorphism is continuous in the cofinite topology.

The next lemma is taken almost verbatim from [1; p. 12].

LEMMA 10. *If $A \in \text{obj } \mathcal{A}$, then A has a fundamental system of neighborhoods of 0 consisting of cofinite ideals.*

Proof: Let A' be a cofinite subgroup of A . Since multiplication is continuous, there is a cofinite subgroup W of A with $W^2 \subset A'$. Since W is cofinite, it has a f.g. free complement $\langle a_1, \dots, a_r \rangle$. For each j , the continuity of $x \mapsto a_j \cdot x$ at 0 implies the existence of a cofinite $W_j \subset W$ with $a_j W_j \subset A'$. If $U = \bigcap_{j=1}^r W_j$, then U is cofinite in A . Moreover, $a_j U \subset A'$ for all j and $WU \subset A'$ (in fact, $W^2 \subset A'$ and $U \subset W$); hence $AU \subset A'$. Since $1 \in A$, we have $U \subset AU$, so that A/AU is f.g. Now if $(AU)_*$ is the pure subgroup of A generated by AU , then $(AU)_*$ is also an ideal, is cofinite, and $(AU)_* \subset A'_* = A'$ (for A' is already pure).

LEMMA 11. *\mathcal{A} has coproducts.*