Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	24 (1978)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ORIGINS OF THE COHOMOLOGY OF GROUPS
Autor:	Mac Lane, Saunders
Kapitel:	9. Spectral Sequences
DOI:	https://doi.org/10.5169/seals-49687

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

pretation for higher n, long sought for, is still missing (and may even not be there!). Eckmann introduced G-finite cohomology groups (1947) and showed their connection with the Hopf-Freudenthal theory of the ends of a group. Eckmann's work, and the paper of Eilenberg-Mac Lane on complexes with operators, again emphasized the connection of cohomology groups of groups with covering spaces. There was a systematic presentation of the subject in the Cartan seminar of 1950/51, entitled "Cohomologie des groupes, suite spectrale, faisceaux". In this seminar Eilenberg first described the cohomology groups axiomatically, and then proved their existence. Subsequent exposés by Cartan emphasized the calculation of the cohomology by free resolutions complete with an abstract version of the comparison theorem. A decisive example of the effective use of such resolutions is the calculation of the cohomology of a cyclic group—carried out here in exposé 3. (I am sensitive to the advantage of using resolutions for this purpose, because in 1948 I had calculated the cohomology of cyclic groups directly from the bar resolution without the general comparison theoremthe direct method worked but was much more cumbersome.) Subsequent exposés made a number of applications-to the Brauer group, the Wedderburn theorem, the theorem of Maschke on complete reducibility of linear representations of a finite group, and P. A. Smith's theorem.

Further applications to pure group theory have been limited. One small but striking one is the homology proof by Gaschutz [1966]:

THEOREM. A finite non-abelian *p*-group has an automorphism of p^{th} power order which is not an inner automorphism.

This conference in Zurich has exhibited more examples of the use of homology in group theory.

9. Spectral Sequences

The results stimulated by group cohomology were not confined just to group theory. For example, the problem of computing the cohomology groups $H^n(G, A)$ for the case when G itself is a group extension (say, cyclic by cyclic) immediately leads to the study of a spectral sequence. Specifically, if

$$1 \to K \to G \to Q \to 1 \tag{1}$$

is a short exact sequence of (multiplicative) groups and A is a left G module there is a spectral sequence E_r^{pq} with

$$E_2^{pq} \cong H^p(Q, H^q(K, M))$$
⁽²⁾

converging to the graded group associated with a filtration of the cohomology $E^{p+q}(G, M)$. In (2), the cohomology $H^{q}(K, M)$ of the subgroup K is suitably interpreted as a Q-module, so that the outside cohomology is defined. The essential portions of such a spectral sequence were discovered by R. Lyndon in his 1946 Harvard thesis, at about the same time that Leray was formulating the general notion of a spectral sequence. Lyndon did use his formulation for computation. Some years later [1953], Hochschild and Serre formulated a spectral sequence like that of (2) in the conventional language, so such a sequence is usually called a Hochschild-Serre spectral sequence. (There are actually several different constructions of such a sequence, and some residual uncertainty as to whether these constructions all yield the same spectral sequence). The essential observation is that computing cohomology or homology in a fiber situation like that of (1) inevitably leads to the spectral sequence technology—whether the fiber situation is group theoretic, as with the exact sequence (1), or a fiber space, as in the case so effectively exploited by Serre in topology.

10. TRANSFER

The operation of *transfer* was well known in group theory, beginning with Burnside's work on monomial representations. If H is a subgroup of index n in G, the transfer from G to H is a homomorphism.

$$t: G / [G, G] \to H / [H, H] \tag{1}$$

between the factor-commutator groups. To define it, choose representatives $x_1, ..., x_n$ of the right cosets of H in G, so that $G = \bigcup Hx_i$ and write $\rho(x)$ for the representative x_i of the coset Hx. Then t is

$$t(g) = \prod_{i=1}^{n} (x_i g) [\rho(x_i g)]^{-1}$$
(2)

This map t is independent of the choice of the set of representatives $x_i, ..., x_n$.

Since the factor commutator group G/[G, G] in (1) is simply the 1dimensional homology group H_1 (G, Z), the transfer can be regarded as a map in homology.

$$t: H_1(G, \mathbb{Z}) \to H_1(H, \mathbb{Z})$$

In 1953 Eckmann extended this map to apply in all dimensions, both in homology and cohomology. Using the standard homogeneous complexes