

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 24 (1978)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ORIGINS OF THE COHOMOLOGY OF GROUPS  
**Autor:** Mac Lane, Saunders  
**Kapitel:** 10. Transfer  
**DOI:** <https://doi.org/10.5169/seals-49687>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$$E_2^{pq} \cong H^p(Q, H^q(K, M)) \quad (2)$$

converging to the graded group associated with a filtration of the cohomology  $E^{p+q}(G, M)$ . In (2), the cohomology  $H^q(K, M)$  of the subgroup  $K$  is suitably interpreted as a  $Q$ -module, so that the outside cohomology is defined. The essential portions of such a spectral sequence were discovered by R. Lyndon in his 1946 Harvard thesis, at about the same time that Leray was formulating the general notion of a spectral sequence. Lyndon did use his formulation for computation. Some years later [1953], Hochschild and Serre formulated a spectral sequence like that of (2) in the conventional language, so such a sequence is usually called a Hochschild-Serre spectral sequence. (There are actually several different constructions of such a sequence, and some residual uncertainty as to whether these constructions all yield the same spectral sequence). The essential observation is that computing cohomology or homology in a fiber situation like that of (1) inevitably leads to the spectral sequence technology—whether the fiber situation is group theoretic, as with the exact sequence (1), or a fiber space, as in the case so effectively exploited by Serre in topology.

## 10. TRANSFER

The operation of *transfer* was well known in group theory, beginning with Burnside's work on monomial representations. If  $H$  is a subgroup of index  $n$  in  $G$ , the transfer from  $G$  to  $H$  is a homomorphism.

$$t : G/[G, G] \rightarrow H/[H, H] \quad (1)$$

between the factor-commutator groups. To define it, choose representatives  $x_1, \dots, x_n$  of the right cosets of  $H$  in  $G$ , so that  $G = \cup Hx_i$  and write  $\rho(x)$  for the representative  $x_i$  of the coset  $Hx$ . Then  $t$  is

$$t(g) = \prod_{i=1}^n (x_i g) [\rho(x_i g)]^{-1} \quad (2)$$

This map  $t$  is independent of the choice of the set of representatives  $x_1, \dots, x_n$ .

Since the factor commutator group  $G/[G, G]$  in (1) is simply the 1-dimensional homology group  $H_1(G, \mathbf{Z})$ , the transfer can be regarded as a map in homology.

$$t : H_1(G, \mathbf{Z}) \rightarrow H_1(H, \mathbf{Z})$$

In 1953 Eckmann extended this map to apply in all dimensions, both in homology and cohomology. Using the standard homogeneous complexes

$B(G)$  and  $B(H)$  for the groups  $G$  and  $H$ , he defined a cochain transformation  $t$  for any  $G$ -module  $A$  and any cochain  $f$  by

$$(tf)(g_0, \dots, g_p) = \sum_{i=1}^n x_i^{-1} f(x_i g_0 (\rho(x_i g_0))^{-1}, \dots, x_i g_n (\rho(x_i g_n))^{-1})$$

This map, up to chain homology, is again independent of the choice of the representatives  $x_j$ , so yields a homomorphism

$$t: H^p(H, A) \rightarrow H^p(G, A).$$

On the other hand, each cochain of  $G$  over  $A$  automatically restricts to a cochain of  $H$  over  $A$ ; this process defines the *restriction* map

$$r: H^p(G, A) \rightarrow H^p(H, A).$$

Eckmann proved that the composite  $tr$  of these maps is the endomorphism given by multiplication by  $n$  in  $H^p(G, A)$ : He made a variety of applications. The notion of transfer was also used by Artin and Tate (see below) in class field theory.

The discovery of the homology of a group had the feature that it exhibited a “non-obvious” construction on groups; in much the same way, the discovery of transfer produced a non-obvious homomorphism between cohomology groups. Thus it is that recently Kahn and Priddy have been able to construct the transfer homeomorphism for the generalized cohomology of an  $n$ -fold covering  $\Pi: E \rightarrow B$ . This transfer applies to the cohomology with coefficients in any strict  $\Omega$ -spectrum; when applied to the Eilenberg-Mac Lane spectrum  $K(\Pi, n)$ , the generalized cohomology is ordinary cohomology and the transfer agrees with the classical one. Using this transfer, they prove a conjecture of Mahowald and Whitehead about a “canonical map” of the  $n$ -fold suspension  $\Sigma^n \mathbf{R}P^{n-1}$  of the real projective  $n - 1$  space into the  $n$  sphere. This map  $\lambda$  is the adjoint of the map

$$\mathbf{R}P^{n-1} \rightarrow O_n \rightarrow \Omega^n S^n.$$

Here the first arrow takes a line through the origin in  $\mathbf{R}^n$  into the reflection in the plane perpendicular to that line; while the second arrow represents each element of  $O^n$  as a map of  $(\mathbf{R}_n \cup \infty, \infty)$  into itself, and hence as an element of the  $n^{\text{th}}$  loop space of  $S^n$ .

The result of Kan and Priddy is that  $\lambda$  is an epimorphism of 2-primary components in stable homotopy.