

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: COINCIDENCE-FIXED-POINT INDEX
Autor: Dold, Albrecht
Kapitel: § 1. The coincidence-fixed-point (c.f.p.) index
DOI: <https://doi.org/10.5169/seals-49689>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§ 1. THE COINCIDENCE-FIXED-POINT (C.F.P.) INDEX

(1.1) Let $p: E \rightarrow B$ denote a euclidean neighborhood retract over B (abbrev. ENR_B), where B , and hence E , is an ENR. Altogether this means that $p: E \rightarrow B$ embeds as a neighborhood retract into the projection $\mathbf{R}^n \times \mathbf{R}^m \rightarrow \mathbf{R}^m$, for some m, n . We refer the reader to [2], §1, for the precise definitions but remark that every smooth submersion and every fibration (with base and total space ENR) qualifies for $p: E \rightarrow B$.

We consider continuous maps $g: D_g \rightarrow E$, $\varphi: D_\varphi \rightarrow B$, where D_g, D_φ are open subsets of E , and $pg = p|_{D_g}$ (i.e., g is fibre-preserving). We let $\text{Fix}(g) = \{x \in D_g \mid gx = x\}$ and $\text{Coinc}(\varphi, p) = \{x \in D_\varphi \mid \varphi x = px\}$, and we assume that $\text{Fix}(g) \cap \text{Coinc}(\varphi, p)$ is compact. Under these circumstances we shall define an integer $J(g, \varphi) \in \mathbf{Z}$ which is akin to the Hopf fixed-point index. It “counts” the points in $\text{Fix}(g) \cap \text{Coinc}(\varphi, p)$ in a weighted and homotopy-invariant fashion. It is the Hopf index of g resp. φ if B is a single point resp. p is the identity map of B .

(1.2) By definition [2], 1.1 of an ENR_B, we have that E is a fibre-preserving neighborhood retract of some $\mathbf{R}^n \times B$. In fact, for the present purpose we can use any product $Y \times B$, i.e. we'll use mappings $E \xrightarrow{i} V \xrightarrow{r} E$ such that $V \subset Y \times B$ is open, $ri = id$, and i, r are maps over B . In formulas,

$$(1.3) \quad ix = (i'x, px), \quad \text{where } i': E \rightarrow Y,$$

$$(1.4) \quad pr(y, b) = b, \quad \text{for } (y, b) \in V,$$

$$(1.5) \quad r(i'x, px) = x, \quad \text{for } x \in E.$$

Consider the following sequence of maps

$$(1.6) \quad D_g \cap D_\varphi \xrightarrow{(g, \varphi)} E \times B \xrightarrow{i' \times id} Y \times B \supset V \xrightarrow{r} E.$$

Its composite $[g, \varphi]$ is defined in $D_V = (i'g, \varphi)^{-1} V$ which is an open subset of $(D_g \cap D_\varphi)$, and hence of E . Thus

$$(1.7) \quad [g, \varphi]: D_V \rightarrow E, \quad [g, \varphi](x) = r(i'gx, \varphi x).$$

If $x \in D_g \cap \text{Coinc}(\varphi, p)$ then

$$(i'g, \varphi)x = (i'gx, px) = (i'gx, pgx) = igx \in V,$$

hence $[g, \varphi]x$ is defined and equals $rigx = gx$. It follows that $\text{Fix}(g) \cap \text{Coinc}(\varphi, p) \subset \text{Fix}[g, \varphi] = \{x \in D_V \mid [g, \varphi]x = x\}$. Conversely, x

$= [g, \varphi] x$ implies $px = p[g, \varphi]x = pr(i'gx, \varphi x) = \varphi x$, hence $x \in \text{Cinc}(\varphi, p)$, and $gx = [g, \varphi]x = x$. Altogether

$$(1.8) \quad \text{Fix}(g) \cap \text{Cinc}(\varphi, p) = \text{Fix}[g, \varphi].$$

In particular, $[g, \varphi]: D_V \rightarrow E$ has a compact fixed-point set, and we can assign to it its Hopf-index $I[g, \varphi] \in \mathbf{Z}$ — for instance as in [1], VII,5.10. Furthermore,

(1.9) PROPOSITION AND DEFINITION. *The Hopf-index $I[g, \varphi] \in \mathbf{Z}$ depends only on (g, φ) , not on the choice of the neighborhood retraction i, r .* We denote this integer by $J(g, \varphi)$, and call it the *c.f.p.-index* of (g, φ) ; thus $J(g, \varphi) = I[g, \varphi]$.

Proof. Because the range B of the maps φ, p is ENR, these two maps are homotopic in a neighborhood of $\text{Cinc}(\varphi, p)$. In fact (cf. [1], IV,8.6), there is an open neighborhood U of $\text{Cinc}(\varphi, p)$ in D_φ , and a deformation $\vartheta_t: U \rightarrow B$, $0 \leq t \leq 1$, such that

$$(1.10) \quad \vartheta_0 = p|U, \quad \vartheta_1 = \varphi|U, \quad \vartheta_t x = px \text{ for } x \in \text{Cinc}(\varphi, p) \text{ and all } t.$$

Consider then two neighborhood retractions

$$\begin{aligned} E &\xrightarrow{i} V \xrightarrow{r} E, \quad V \subset Y \times B; \quad ix = (i'x, px), \\ E &\xrightarrow{j} W \xrightarrow{s} E, \quad W \subset Z \times B; \quad jx = (j'x, px), \end{aligned}$$

as above, and the corresponding maps $[g, \varphi]_1, [g, \varphi]_2$ as defined by 1.6. We have to show $I([g, \varphi]_1) = I([g, \varphi]_2)$. In order to do so we can (cf. [1], VII,5.11) restrict attention to an arbitrary open neighborhood N of $\text{Fix}([g, \varphi]_i) = \text{Fix}(g) \cap \text{Cinc}(\varphi, p)$. And we shall show that $[g, \varphi]_i|N$ are homotopic ($i=1, 2$) without moving the fixed point set, provided N is sufficiently small. The homotopy is given by the formula

$$(1.11) \quad \theta_t x = s(j'r(i'gx, \vartheta_t x), \varphi x).$$

This is defined for (x, t) such that $x \in D_g \cap U$, $v = (i'gx, \vartheta_t x) \in V$, and $w = (j'rv, \varphi x) \in W$; the set of all such (x, t) is an open subset D_θ of $E \times [0, 1]$. If $x \in \text{Fix}(g) \cap \text{Cinc}(\varphi, p)$ then

$$v = (i'gx, \vartheta_t x) = (i'x, px) = ix \in V, \text{ and } rv = x,$$

hence

$$w = (j'rv, \varphi x) = (j'x, px) = jx \in W, \text{ and } \theta_t x = sw = x.$$

Therefore, $(\text{Fix}(g) \cap \text{Coinc}(\varphi, p)) \times [0, 1] \subset D_\theta$, and $(\text{Fix}(g) \cap \text{Coinc}(\varphi, p)) \subset \text{Fix}(\theta_t)$ for all t . It follows that

$$N = \{x \in E \mid (x, t) \in D_\theta \text{ for all } t\}$$

is an open neighborhood of $\text{Fix}(g) \cap \text{Coinc}(\varphi, p)$ in which the deformation θ is defined (by 1.11).

Suppose now $x \in N$ is a fixed point of θ_t , thus $x = s(j'r(i'gx, \vartheta_t x), \varphi x)$. Apply p , using 1.4 for s , and get $px = \varphi x$, hence $\vartheta_t x = px$ by 1.10, hence $r(i'gx, \vartheta_t x) = r(i'gx, px) = r(i'gx, pgx) = rigx = gx$, hence $x = \theta_t x = s(j'gx, px) = s(j'gx, pgx) = sjgx = gx$; altogether, $x \in \text{Coinc}(\varphi, p) \cap \text{Fix}(g)$. It follows that the fixed point set $\text{Fix}(\theta_t) = \text{Fix}(g) \cap \text{Coinc}(\varphi, p)$ for all t . In particular, $\cup_{t \in [0, 1]} \text{Fix}(\theta_t)$ is compact, hence (cf. [1].VII,5.15) all θ_t have the same Hopf-index $I(\theta_t)$. But $r(i'gx, \vartheta_0 x) = r(i'gx, px) = r(i'gx, pgx) = gx$, hence $\theta_0 x = s(j'gx, \varphi x) = [g, \varphi]_2 x$. To calculate θ_1 we first remark that $p[g, \varphi]_1 x = \varphi x$, by 1.7 and 1.4; also $r(i'gx, \vartheta_1 x) = r(i'gx, \varphi x) = [g, \varphi]_1 x$, hence $\theta_1 x = s(j'[g, \varphi]_1 x, p[g, \varphi]_1 x) = sj[g, \varphi]_1 x = [g, \varphi]_1 x$. \square

(1.12) *The product case $E = F \times B$, p = projection.* In this case $g: D_g \rightarrow F \times B$ has the form $g(y, b) = (\gamma(y, b), b)$ with $\gamma: D_g \rightarrow F$. The two maps (γ, φ) combine to a map $(\gamma, \varphi): D \rightarrow F \times B$, where $D (= D_g \cap D_\varphi)$ is an open subset of $F \times B$, and $\text{Fix}(\gamma, \varphi) = \text{Fix}(g) \cap \text{Coinc}(\varphi, p)$. In order to obtain the c.f.p.-index $J(g, \varphi)$ one can use $Y = F$ and the neighborhood retraction $i = r = \text{identity-map}$ of $Y \times B$. The definition 1.9 then shows that

$$J(g, \varphi) = I(\gamma, \varphi);$$

i.e. *in the product case the c.f.p.-index of (g, φ) is simply the Hopf-index of $(y, b) \mapsto (\gamma(y, b), \varphi(y, b))$.*

The procedure 1.6-1.9 in the general case, on the other hand, can be considered as a reduction to the product case.

(1.13) *General properties of $J(g, \varphi)$* follow from corresponding properties of the Hopf-index. For instance, $J(g, \varphi)$ is additive with respect to topological-sum decompositions of $\text{Fix}(g) \cap \text{Coinc}(g, \varphi)$, it is invariant under deformations such that $\cup_{0 \leq t \leq 1} \text{Fix}(g_t) \cap \text{Coinc}(\varphi_t, p)$ is compact, it

depends only on the germ of (g, φ) around $\text{Fix}(g) \cap \text{Coinc}(\varphi, p)$ — in particular, $J(g, \varphi) = 0$ if $\text{Fix}(g) \cap \text{Coinc}(\varphi, p) = \emptyset$, etc. These details are left to the reader. Lefschetz-trace formulas for $J(g, \varphi)$ can be found in 2.1 and 3.5.