

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 24 (1978)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE GELFAND-FUKS COHOMOLOGY  
**Autor:** Haefliger, André  
**Kapitel:** 1. Definitions  
**DOI:** <https://doi.org/10.5169/seals-49696>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# ON THE GELFAND-FUKS COHOMOLOGY <sup>1</sup>

by André HAEFLIGER

In this talk, we would like to report on the work of Gelfand and Fuks on the cohomology of the Lie algebra  $L_M$  of smooth vector fields on a manifold  $M$ , as well as on more recent developments, some of them obtained in collaboration with Raoul Bott.

## 1. DEFINITIONS

*Gelfand-Fuks cohomology.*

$L_M$  will denote the Lie algebra of smooth vector fields on  $M$ , with the topology of uniform convergence of all derivatives on compact sets. For  $M$  compact,  $L_M$  can be thought as the Lie algebra of the group  $\text{Diff}_M$  of diffeomorphisms of  $M$ .

Gelfand and Fuks [7], have considered the differential graded algebra  $C^*(L_M)$  of *continuous* multilinear alternate forms on  $L_M$  with values in  $R$ , the differential of a  $k$ -form  $f$  being the  $(k+1)$ -form  $df$  defined by

$$df(v_0, \dots, v_k) = \sum_{0 \leq r < s \leq k} (-1)^{r+s} f([v_r, v_s], v_0, \dots, \hat{v}_r, \dots, \hat{v}_s, \dots, v_k)$$

where the  $v_i$ 's are vector fields on  $M$ . So those cochains are like distributions.

Suppose that  $G$  is a Lie group acting smoothly and effectively on  $M$ . Then the Lie algebra  $\mathfrak{g}$  of  $G$  is identified with a subalgebra of  $L_M$ . We shall denote by  $C^*(L_M; G)$  the subalgebra of  $C^*(L_M)$  of  $G$ -basic cochains, namely cochains invariant by  $G$  and which vanish if one of the argument  $v_i$  belongs to  $\mathfrak{g}$ .

The cohomology of  $C^*(L_M)$  (resp.  $C^*(L_M; G)$ ) will be denoted by  $H^*(L_M)$  (resp.  $H^*(L_M; G)$ ), and will be called the Gelfand-Fuks cohomology of  $M$  (resp. of  $M$  rel. to  $G$ ).

<sup>1</sup>) Presented at the Colloquium on Topology and Algebra, April 1977, Zurich

*Models (Sullivan Theory)* (cf. [18]).

$C^*(L_M)$  and  $C^*(L_M; G)$  are examples of differential graded commutative (in the graded sense) algebras over  $\mathbf{R}$ , abbreviated *DG*-algebras.

Among *DG*-algebras, we consider the equivalence relation generated by " $A \sim B$ " if there is a morphism  $\varphi: A \rightarrow B$  of *DG*-algebras inducing an isomorphism on cohomology. We say that  $M$  is a *model* for  $A$  if  $M$  is equivalent to  $A$  under this equivalence relation. Following the terminology of Sullivan, we say that  $M$  is a *minimal model* for  $A$  (assuming  $H^0(A) = \mathbf{R}$  and  $H^1(A) = 0$ ) if  $M$  is a *free algebra* (namely the tensor product of a polynomial algebra on even dimensional generators by an exterior algebra on odd dimensional generators), the differential of each generator being decomposable (we also assume that generators are of degree bigger than one). The free algebra on a set of generators  $x_\alpha$  will be denoted by  $A(x_\alpha)$ .

There is a contravariant functor from the category of topological spaces to the category of *DG*-algebras associating to the space  $X$  the *DG*-algebra  $A^*(X)$  of real polynomial forms on its singular complex. If one takes instead rational polynomial forms, this functor induces an equivalence between rational homotopy types of 1-connected spaces with finite dimensional cohomology and equivalence classes of 1-connected *DG* algebras over  $\mathbf{Q}$  with finite dimensional cohomology. A minimal model corresponds to a Postnikov decomposition. In particular the vector space of generators in the minimal model is the dual of the graded vector space  $\pi_*(X) \otimes \mathbf{R}$ , where  $\pi_i(X)$  is the  $i$ -th homotopy group of  $X$ .

We shall say that a *DG*-algebra  $A$  is a model for the space  $X$  if it is a model for the *DG*-algebra  $A^*(X)$ .

The main problem is to find good models for the *DG*-algebras  $C^*(L_M)$  or  $C^*(L_M; G)$ , if possible finite dimensional in each degree.

As an example computed by Gelfand and Fuks [6], consider the case of the circle  $S^1$ . Then  $H^*(L_{S^1})$  is the free algebra on generators  $u$  and  $v$  of degree 2 and 3 represented by the cocycles

$$u(f, g) = \int_0^1 \begin{vmatrix} f' & f'' \\ g' & g'' \end{vmatrix} dx \quad \text{and} \quad v(f, g, h) = \int \begin{vmatrix} f & f' & f'' \\ g & g' & g'' \\ h & h' & h'' \end{vmatrix} dx$$

where the vector fields on  $S^1$  are identified with functions of period 1 on  $\mathbf{R}$ . This is also a model for  $C^*(L_{S^1})$ .

If  $G$  is the group  $SO_2$  of rotations of  $S^1$ , then  $H(L_{S^1}; SO_2)$  is a model for  $C^*(L_{S^1}; SO_2)$ . It is generated by  $u$  and by an element  $e$  of degree 2 represented by

$$e(f, g) = \int_0^1 \begin{vmatrix} f & g \\ f' & g' \end{vmatrix} dx$$

The only relation is  $eu = 0$ .

## 2. CONNECTION WITH FOLIATIONS

Let me indicate very briefly the relation with characteristic classes of flat bundles (cf. [12]).

$H^*(L_M, G)$  could also be interpreted as the differentiable cohomology of a suitable differentiable category (for more informations see [4] and [15]).

We consider on the product  $X \times M$  of a smooth manifold  $X$  with  $M$  a smooth foliation  $F$  whose leaves have the same dimension as  $X$  and cut each fibers  $\{x\} \times M$  transversally.

To such a foliation is naturally associated a continuous  $DG$ -algebra map

$$\chi_F : C^*(L_M) \rightarrow \Omega_X$$

where  $\Omega_X$  is the  $DG$ -algebra of differential forms on  $X$ . In fact there is a bijection between such morphisms and foliations  $F$  as above.

Passing to cohomology, we get the characteristic map

$$H^*(L_M) \rightarrow H^*(X; R)$$

If we replace the trivial bundle by a bundle  $E$  with fiber  $M$ , base space  $X$  and structural group  $G$ , then for a foliation  $F$  on  $E$  complementary to the fibers, we still get a morphism

$$\chi_F : C^*(L_M; G) \rightarrow \Omega_X$$

hence a characteristic homomorphism

$$H^*(L_M, G) \rightarrow H^*(X; R)$$

Denoting by  $BG$  the classifying space for  $G$ -bundles, we also have the usual characteristic map  $H^*(BG; R) \rightarrow H^*(X; R)$ . This map factorizes