

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 24 (1978)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE GELFAND-FUKS COHOMOLOGY  
**Autor:** Haefliger, André  
**Kapitel:** 2. Connection with foliations  
**DOI:** <https://doi.org/10.5169/seals-49696>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

If  $G$  is the group  $SO_2$  of rotations of  $S^1$ , then  $H(L_{S^1}; SO_2)$  is a model for  $C^*(L_{S^1}; SO_2)$ . It is generated by  $u$  and by an element  $e$  of degree 2 represented by

$$e(f, g) = \int_0^1 \left| \begin{array}{cc} f, & g \\ f', & g' \end{array} \right| dx$$

The only relation is  $e u = 0$ .

## 2. CONNECTION WITH FOLIATIONS

Let me indicate very briefly the relation with characteristic classes of flat bundles (cf. [12]).

$H^*(L_M, G)$  could also be interpreted as the differentiable cohomology of a suitable differentiable category (for more informations see [4] and [15]).

We consider on the product  $X \times M$  of a smooth manifold  $X$  with  $M$  a smooth foliation  $F$  whose leaves have the same dimension as  $X$  and cut each fibers  $\{x\} \times M$  transversally.

To such a foliation is naturally associated a continuous  $DG$ -algebra map

$$\chi_F: C^*(L_M) \rightarrow \Omega_X$$

where  $\Omega_X$  is the  $DG$ -algebra of differential forms on  $X$ . In fact there is a bijection between such morphisms and foliations  $F$  as above.

Passing to cohomology, we get the characteristic map

$$H^*(L_M) \rightarrow H^*(X; R)$$

If we replace the trivial bundle by a bundle  $E$  with fiber  $M$ , base space  $X$  and structural group  $G$ , then for a foliation  $F$  on  $E$  complementary to the fibers, we still get a morphism

$$\chi_F: C^*(L_M; G) \rightarrow \Omega_X$$

hence a characteristic homomorphism

$$H^*(L_M, G) \rightarrow H^*(X; R)$$

Denoting by  $BG$  the classifying space for  $G$ -bundles, we also have the usual characteristic map  $H^*(BG; R) \rightarrow H^*(X; R)$ . This map factorizes

through a map  $H^*(BG; R) \rightarrow H^*(L_M; G)$  so that we get a commutative diagram

$$\begin{array}{ccc} & H^*(L_M; G) & \\ H^*(BG; R) & \swarrow & \downarrow \\ & H^*(X; R) & \end{array}$$

So it is important to compute the map  $H^*(BG; R) \rightarrow H^*(L_M; G)$ . When  $G$  is a compact connected Lie group, then  $H^*(BG; R)$  is the algebra  $I(G)$  of invariant polynomials on the Lie algebra of  $G$ , and the map from  $I(G)$  to  $C^*(L_M; G)$  is given by a  $G$ -connexion in  $C^*(L_M)$  (cf. [5]).

In the example above, namely  $M = S^1$  and  $G = SO_2$ , then  $H^*(BSO_2)$  is a polynomial algebra in a generator of degree 2, the Euler class, which is mapped on a non zero multiple of  $e$ .

### 3. THE FORMAL VECTOR FIELDS AND THE DIAGONAL COMPLEX

Given a point  $x$  on  $M$ , we can consider the Lie algebra  $L_M^x$  of infinite jets at  $x$  of vector fields on  $M$  with the quotient topology. It is isomorphic to the Lie algebra  $\mathfrak{a}_n$  of formal vector fields  $\sum v_i(x) \partial/\partial x^i$  in  $R^n$ , where the  $v_i(x)$  are formal power series in the coordinates  $x^1, \dots, x^n$ .

The natural map  $L_M \rightarrow L_M^x$  associating to a vector field its jet at  $x$  gives a  $DG$ -algebra morphism

$$C^*(L_M^x) \rightarrow C^*(L_M)$$

where  $C^*(L_M^x)$  is the algebra of multilinear alternate forms on  $L_M^x$  depending only on finite order jets.

The first and most important step in the work of Gelfand-Fuks was the complete determination of the cohomology  $H^*(\mathfrak{a}_n)$  of the topological Lie algebra of formal vector fields on  $R^n$ .

**THEOREM 1.** (Gelfand-Fuks [8], [9]). *Let  $E(h_1, \dots, h_n)$  be the exterior algebra on generators  $h_i$  of degree  $2i-1$  and let  $R[c_1, \dots, c_n]_{2n}^\wedge$  be the quotient of the polynomial algebra in generators  $c_i$  of degree  $2i$  by the ideal of elements of degree  $> 2n$ .*