Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GELFAND-FUKS COHOMOLOGY
Autor: Haefliger, André
Kapitel: 2. Connection with foliations
DOI: https://doi.org/10.5169/seals-49696

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

If G is the group SO_2 of rotations of S^1, then $H(L_{S^1}; SO_2)$ is a model for $C^*(L_{S^1}; SO_2)$. It is generated by u and by an element e of degree 2 represented by

$$e(f, g) = \int_0^1 \begin{vmatrix} f, & g \\ f', & g' \end{vmatrix} dx$$

The only relation is $e\,u = 0$.

2. Connection with foliations

Let me indicate very briefly the relation with characteristic classes of flat bundles (cf. [12]).

$H^*(L_M, G)$ could also be interpreted as the differentiable cohomology of a suitable differentiable category (for more informations see [4] and [15]).

We consider on the product $X \times M$ of a smooth manifold X with M a smooth foliation F whose leaves have the same dimension as X and cut each fibers $\{x\} \times M$ transversally.

To such a foliation is naturally associated a continuous DG-algebra map

$$\chi_F : C^*(L_M) \to \Omega_X$$

where Ω_X is the DG-algebra of differential forms on X. In fact there is a bijection between such morphisms and foliations F as above.

Passing to cohomology, we get the characteristic map

$$H^*(L_M) \to H^*(X; R)$$

If we replace the trivial bundle by a bundle E with fiber M, base space X and structural group G, then for a foliation F on E complementary to the fibers, we still get a morphism

$$\chi_F : C^*(L_M; G) \to \Omega_X$$

hence a characteristic homomorphism

$$H^*(L_M, G) \to H^*(X; R)$$

Denoting by BG the classifying space for G-bundles, we also have the usual characteristic map $H^*(BG; R) \to H^*(X; R)$. This map factorizes

through a map $H^*(BG; R) \to H^*(L_M; G)$ so that we get a commutative diagram

$$\begin{tikzcd} & H^*(L_M; G) \arrow[d] \\ H^*(BG; R) \arrow[ur] \arrow[r] & H^*(X; R) \end{tikzcd}$$

So it is important to compute the map $H^*(BG; R) \to H^*(L_M; G)$. When G is a compact connected Lie group, then $H^*(BG; R)$ is the algebra $I(G)$ of invariant polynomials on the Lie algebra of G, and the map from $I(G)$ to $C^*(L_M; G)$ is given by a G-connexion in $C^*(L_M)$ (cf. [5]).

In the example above, namely $M = S^1$ and $G = SO_2$, then $H^*(BSO_2)$ is a polynomial algebra in a generator of degree 2, the Euler class, which is mapped on a non zero multiple of e.

3. The formal vector fields and the diagonal complex

Given a point x on M, we can consider the Lie algebra L_M^x of infinite jets at x of vector fields on M with the quotient topology. It is isomorphic to the Lie algebra $\mathfrak{a}_n$ of formal vector fields $\sum v_i(x)\, \partial/\partial x^i$ in R^n, where the $v_i(x)$ are formal power series in the coordinates $x^1, \ldots, x^n$.

The natural map $L_M \to L_M^x$ associating to a vector field its jet at x gives a DG-algebra morphism

$$C^*(L_M^x) \to C^*(L_M)$$

where $C^*(L_M^x)$ is the algebra of multilinear alternate forms on L_M^X depending only on finite order jets.

The first and most important step in the work of Gelfand-Fuks was the complete determination of the cohomology $H^*(\mathfrak{a}_n)$ of the topological Lie algebra of formal vector fields on R^n.

Theorem 1. (Gelfand-Fuks [8], [9]). *Let $E(h_1, \ldots, h_n)$ be the exterior algebra on generators h_i of degree $2i-1$ and let $R[c_1, \ldots, c_n]_{2n}^{\wedge}$ be the quotient of the polynomial algebra in generators c_i of degree $2i$ by the ideal of elements of degree $> 2n$.*