

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GELFAND-FUKS COHOMOLOGY
Autor: Haefliger, André
Kapitel: 4. Main theorem
DOI: <https://doi.org/10.5169/seals-49696>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THEOREM 3. (Guillemin [10], Losik [17]). $C_{\Delta}^*(L_M, \Omega_M)$ is a model for a bundle E with fiber F_n , base space M , associated to the tangent bundle of M .

More precisely, a model for $C_{\Delta}^*(L_M, \Omega_M)$ is the DG-algebra $\Omega_M \otimes WU_n$ over Ω_M , where

$$d(1 \otimes c_i) = 0 \quad d(1 \otimes h_i) = 1 \otimes c_i - p_{i/2} \otimes 1$$

where $p_{i/2}$ is zero if i is odd and is a form representing the Pontrjagin class of M of degree $2i$ if i is even.

Note that if a foliation F on $X \times M$ transverse to the fibers $\{x\} \times M$ is given, one has a characteristic homomorphism

$$C^*(L_M, \Omega_M) \rightarrow \Omega_{X \times M}$$

One has also a morphism

$$WO_n \rightarrow C_{\Delta}^*(L_M, \Omega_M)$$

(or $WU_n \rightarrow C^*(L_M, \Omega_M)$ in case M has trivial Pontrjagin classes) whose composition with the previous one is the usual characteristic homomorphism for the foliation F (cf. [3], [12]).

4. MAIN THEOREM

THEOREM 1. $C^*(L_M)$ is a model for the space Γ of continuous sections of the bundle E described in the theorem above.

This result, first conjectured by Bott (and also Fuks), has been proved by several people (Bott-Segal¹), Fuks-Segal, Haefliger [13], Ph. Trauber, and others).

Suppose that G is a compact connected Lie group acting on M . Then it also acts on the bundle E and on its space of sections. Let us denote by Γ_G the total space of the bundle with fiber Γ associated to the universal G -bundle with base space BG .

THEOREM 1'. $C^*(L_M; G)$ is a model for the space Γ_G .

The way I proved theorem 1 was to construct first a tentative algebraic model A for Γ following ideas of R. Thom [20] and D. Sullivan [18], and

¹) Added on proof: *Topology* 16 (1977), pp. 285-298.

a morphism of A in $C^*(L_M)$. Then one proves directly that it induces an isomorphism in cohomology. The fact that A is also a model for Γ was proved in a similar way (cf. [14]).

When M has a finite dimensional model, one can construct a model for Γ which is finite dimensional in each degree, and with it one can make explicit calculations.

Note that the inclusion $C_{\Delta}^*(L_M, \Omega_M) \rightarrow C^*(L_M, \Omega_M)$ is a model for the evaluation map $\Gamma \times M \rightarrow E$ associating to a section s and a point x of M the element $s(x)$ of E .

For computations along the lines of the spectral sequence of Gelfand-Fuks, see Cohen and Taylor [22].

The proof of theorem 1' is very similar to the proof of theorem 1. In the next paragraph, we explain the construction of an algebraic model for Γ_G suitable for computations. In § 6, we indicate briefly why this is a model for Γ_G .

5. CONSTRUCTION OF AN ALGEBRAIC MODEL FOR THE SPACE OF SECTIONS OF A FIBER BUNDLE ([20], [18], [13]).

As a guide, consider first the geometric situation. Let $p: E \rightarrow M$ be a fiber bundle with base space M , fiber F and let Γ be the space of continuous sections of E .

We have the commutative diagramm

$$\begin{array}{ccc}
 & & e \\
 & & \longrightarrow \\
 & M \times \Gamma & \longrightarrow E \\
 & \downarrow & \searrow \\
 & & M \\
 & \downarrow & \downarrow \\
 & \Gamma & \\
 & \searrow & \downarrow \\
 & & *
 \end{array}$$

1)

where e is the evaluation map associating to the point x of M and the section s the point $s(x)$ of E . The other maps are natural projections (* is a point).