

6. Sketch of the proof of the main theorem and applications

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **24 (1978)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

is the free commutative algebra $\Lambda(s, e)$, where $\deg s = 1$ and $ds = e$. As a B -module, it is free with basis 1 and s . A model for E_G is just $A \otimes \Lambda(x_\alpha)$.

As model for Γ_G , we take $R[e] \otimes \Lambda(x_\alpha, \bar{x}_\alpha)$, where $\deg \bar{x}_\alpha = \deg x_\alpha - 1$, the image of x_α by ε being $1 \otimes x_\alpha + s \otimes \bar{x}_\alpha$. The differential d is described as follows (compare with Sullivan [18] or [19]). Let h be the derivation of degree -1 of $\Lambda(x_\alpha, \bar{x}_\alpha)$ given by $hx_\alpha = \bar{x}_\alpha$ and $h\bar{x}_\alpha = 0$. Then if d_0 denotes the differential in $\Lambda(x_\alpha)$ identified to a subalgebra of $\Lambda(x_\alpha, \bar{x}_\alpha)$, we have

$$de = 0, dx_\alpha = d_0 x_\alpha - e \bar{x}_\alpha, d\bar{x}_\alpha = -hd_0 x_\alpha$$

Remark. In the case where E is the bundle described in § 4, its minimal model $A \otimes \Lambda(x_\alpha)$ over M_G is complicated, because there is an infinite number of generators x_α (except for $n=1$) labelled by a basis of the rational homotopy of a wedge of spheres, so by a basis of the free graded Lie algebra $L(n)$ generated by the spheres of this wedge (cf. [13]).

6. SKETCH OF THE PROOF OF THE MAIN THEOREM AND APPLICATIONS

We represent the universal principal G -bundle as a limit of finite dimensional bundles P_k and we denote by Ω_P the inverse limit of algebras of forms Ω_{P_k} .

First note that we can replace $C^*(L_M; G)$ by the DG -algebra $C^*(L_M, \Omega_P)_G$ of G -basic elements in $C^*(L_M, \Omega_P)$ (compare with Cartan [5], exposé 20).

A model for E_G will be the algebra $C_\Delta^*(L_M, \Omega_{M \times P})_G = [C_\Delta^*(L_M, \Omega_M \hat{\otimes} \Omega_P)]_G$ and a model for the evaluation map will be the inclusion of this DG -algebra in $C^*(L_M, \Omega_{M \times P})_G$.

In the construction of § 5, we choose $B = \Omega_{BG}$ as model for BG and, instead of taking for A a finite dimensional module over B , we take the DG -algebra $\Omega_{M_G} \approx [\Omega_{M \times P}]_G$ as model for M_G . We have to build the model for Γ_G along the same lines as in § 5, but in more intrinsic terms like in [13]. The minimal model (or Postnikov decomposition of E_G) will be of the form $A \otimes S^*(V)$, where $S^*(V)$ denotes the algebra of symmetric multilinear forms on a graded vector space V (cf. [13]).

As an algebra, the model for Γ_G will be the algebra $S_B^*(A \otimes V, B)$ of continuous symmetric B -multilinear forms on the graded B -module $A \otimes V$. One can construct a map of this model in $C^*(L_M, \Omega_{M \times P})_G$ and prove that it induces an isomorphism in cohomology.

Similarly, one can prove that $S_B^*(A \otimes V, B)$ is effectively a model for the space of sections Γ_G (cf. [14]).

Eventually for computations, one proves that one gets also a model for Γ_G by using instead of Ω_{M_G} a *DG*-algebra A as in § 5 which is a finite dimensional free B -module.

7. EXAMPLE OF A COMPUTATION

Let us consider the case where M is the n -sphere S^n , G the rotation group SO_{n+1} and E the bundle described in § 3. A model for M_G is the *DG*-algebra A defined by

$$A = R[p_1, \dots, p_k, s] / (s^2 - p_k) \quad d \equiv 0 \quad \text{for } n = 2k$$

or $A = R[p_1, \dots, p_{k-1}, \chi] \otimes E(s) \quad ds = \chi \quad \text{for } n = 2k-1$

where $\deg p_i = 4i$ and $\deg s = n$.

A model for E_G is obtained by taking the tensor product of A with WU_n , the differential being defined by

$$dh_i = c_i - p_{i/2} \quad \text{and} \quad dc_i = 0.$$

By the way, WSO_n is also a model for E_G .

We now consider the case $n = 2$. The minimal model of E_G is the *DG*-algebra which begins as

$$A \otimes \Lambda(x_1, x_2, x_3, x_4, x_5, x_{12}, x_{13}, x_{23}, \dots)$$

where

$$\begin{aligned} \deg x_1 &= \deg x_2 = 5, \deg x_3 = 7, \deg x_4 = \deg x_5 = 8, \\ \deg x_{12} &= 9, \deg x_{13} = \deg x_{23} = 11, \end{aligned}$$

etc.

(there is an infinite number of generators).

The differential is defined by

$$\begin{aligned} dx_1 &= dx_2 = 0, dx_3 = -p_1^2, dx_4 = p_1 x_1, dx_5 = p_1 x_2, \\ dx_{12} &= x_1 x_2, dx_{13} = x_1 x_3 - p_1 x_4, dx_{23} = x_2 x_3 - p_1 x_5, \end{aligned}$$

etc.

According to the construction of § 5, a minimal model for the bundle $\Gamma_G \rightarrow B_G$ begins as

$$R[p_1] \otimes \Lambda(\bar{x}_1, \bar{x}_2, x_1, x_2, \bar{x}_3, \bar{x}_4, \bar{x}_5, x_3, \bar{x}_{12}, x_4, x_5, \dots)$$