Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 24 (1978)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GELFAND-FUKS COHOMOLOGY
Autor: Haefliger, André
Kapitel: 6. Sketch of the proof of the main theorem and applications
DOI: https://doi.org/10.5169/seals-49696

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 20.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

is the free commutative algebra $\Lambda(s, e)$, where $\deg s = 1$ and $ds = e$. As a B-module, it is free with basis 1 and s. A model for E_G is just $A \otimes \Lambda(x_\alpha)$.

As model for Γ_G, we take $R[e] \otimes \Lambda(x_\alpha, \bar{x}_\alpha)$, where $\deg \bar{x}_\alpha = \deg x_\alpha - 1$, the image of x_α by ε being $1 \otimes x_\alpha + s \otimes \bar{x}_\alpha$. The differential d is described as follows (compare with Sullivan [18] or [19]). Let h be the derivation of degree -1 of $\Lambda(x_\alpha, \bar{x}_\alpha)$ given by $hx_\alpha = \bar{x}_\alpha$ and $h\bar{x}_\alpha = 0$. Then if d_0 denotes the differential in $\Lambda(x_\alpha)$ identified to a subalgebra of $\Lambda(x_\alpha, \bar{x}_\alpha)$, we have

$$de = 0, dx_\alpha = d_0 x_\alpha - e\,\bar{x}_\alpha, d\bar{x}_\alpha = -hd_0 x_\alpha$$

Remark. In the case where E is the bundle described in § 4, its minimal model $A \otimes \Lambda(x_\alpha)$ over M_G is complicated, because there is an infinite number of generators x_α (except for $n=1$) labelled by a basis of the rational homotopy of a wedge of spheres, so by a basis of the free graded Lie algebra $L(n)$ generated by the spheres of this wedge (cf. [13]).

6. Sketch of the proof of the main theorem and applications

We represent the universal principal G-bundle as a limit of finite dimensional bundles P_k and we denote by Ω_P the inverse limit of algebras of forms Ω_{P_k}.

First note that we can replace $C^*(L_M; G)$ by the DG-algebra $C^*(L_M, \Omega_P)_G$ of G-basic elements in $C^*(L_M, \Omega_P)$ (compare with Cartan [5], exposé 20).

A model for E_G will be the algebra $C^*_\Delta(L_M, \Omega_{M\times P})_G = [C^*_\Delta(L_M, \Omega_M \hat{\otimes} \Omega_P]_G$ and a model for the evaluation map will be the inclusion of this DG-algebra in $C^*(L_M, \Omega_{M\times P})_G$.

In the construction of § 5, we choose $B = \Omega_{BG}$ as model for BG and, instead of taking for A a finite dimensional module over B, we take the DG-algebra $\Omega_{M_G} \approx [\Omega_{M\times P}]_G$ as model for M_G. We have to build the model for Γ_G along the same lines as in § 5, but in more intrinsic terms like in [13]. The minimal model (or Postnikov decomposition of E_G) will be of the form $A \otimes S^*(V)$, where $S^*(V)$ denotes the algebra of symmetric multilinear forms on a graded vector space V (cf. [13]).

As an algebra, the model for Γ_G will be the algebra $S^*_B(A \otimes V, B)$ of continuous symmetric B-multilinear forms on the graded B-module $A \otimes V$. One can construct a map of this model in $C^*(L_M, \Omega_{M\times P})_G$ and prove that it induces an isomorphism in cohomology.

Similarly, one can prove that $S_B^*(A \otimes V, B)$ is effectively a model for the space of sections Γ_G (cf. [14]).

Eventually for computations, one proves that one gets also a model for Γ_G by using instead of Ω_{M_G} a DG-algebra A as in § 5 which is a finite dimensional free B-module.

7. Example of a computation

Let us consider the case where M is the n-sphere S^n, G the rotation group SO_{n+1} and E the bundle described in § 3. A model for M_G is the DG-algebra A defined by

$$A = R[p_1, \ldots, p_k, s]/(s^2 - p_k) \quad d \equiv 0 \quad \text{for} \quad n = 2k$$

$$\text{or} \quad A = R[p_1, \ldots, p_{k-1}, \chi] \otimes E(s) \quad ds = \chi \quad \text{for} \quad n = 2k-1$$

where $\deg p_i = 4i$ and $\deg s = n$.

A model for E_G is obtained by taking the tensor product of A with WU_n, the differential being defined by

$$dh_i = c_i - p_{i/2} \quad \text{and} \quad dc_i = 0 .$$

By the way, WSO_n is also a model for E_G.

We now consider the case $n = 2$. The minimal model of E_G is the DG-algebra which begins as

$$A \otimes \Lambda(x_1, x_2, x_3, x_4, x_5, x_{12}, x_{13}, x_{23}, \ldots)$$

where

$$\deg x_1 = \deg x_2 = 5, \deg x_3 = 7, \deg x_4 = \deg x_5 = 8 ,$$

$$\deg x_{12} = 9, \deg x_{13} = \deg x_{23} = 11 ,$$

etc.

(there is an infinite number of generators).

The differential is defined by

$$dx_1 = dx_2 = 0, dx_3 = -p_l^2, dx_4 = p_1 x_1, dx_5 = p_1 x_2 ,$$

$$dx_{12} = x_1 x_2, dx_{13} = x_1 x_3 - p_1 x_4, dx_{23} = x_2 x_3 - p_1 x_5 ,$$

etc.

According to the construction of § 5, a minimal model for the bundle $\Gamma_G \to B_G$ begins as

$$R[p_1] \otimes \Lambda(\bar{x}_1, \bar{x}_2, x_1, x_2, \bar{x}_3, \bar{x}_4, \bar{x}_5, x_3, \bar{x}_{12}, x_4, x_5, \ldots)$$